

Pacific Journal of Mathematics

0-PRIMITIVE ORDERED PERMUTATION GROUPS

STEPHEN H. MCCLEARY

O-PRIMITIVE ORDERED PERMUTATION GROUPS

STEPHEN H. MCCLEARY

Let G be a transitive l -subgroup of the lattice-ordered group $A(\Omega)$ of all order-preserving permutations of a chain Ω . (In fact, many of the results are generalized to partially ordered sets Ω and transitive groups G such that $\beta < \gamma$ implies $\beta g = \gamma$ for some positive $g \in G$, thus encompassing some results on non-ordered permutation groups.) The orbits of any stabilizer subgroup G_α , $\alpha \in \Omega$, are convex and thus can be totally ordered in a natural way. The usual pairing $\Delta \longleftrightarrow \Delta' = \{\alpha g \mid \alpha \in \Delta g\}$ establishes an o -anti-isomorphism between the set of "positive" orbits and the set of "negative" orbits. If Δ is an o -block (convex block) of G for which $\Delta G_\alpha = \Delta$, then Δ' is also an o -block. If G_α has a greatest orbit Γ , then $\{\beta \in \Omega \mid \Gamma' < \beta < \Gamma\}$ constitutes an o -block of G . A correspondence is established between the centralizer $Z_{A(\Omega)}G$ and a certain subset of the fixed points of G_α .

The main theorem states that every o -primitive group (G, Ω) which is not o -2-transitive or regular looks strikingly like the only previously known example, in which Ω is the reals and $G = \{f \in A(\Omega) \mid (\beta + 1)f = \beta f + 1 \text{ for all } \beta \in \Omega\}$. The "configuration" of orbits of G_α must consist of a set o -isomorphic to the integers of "long" (infinite) orbits with some fixed points interspersed; and there must be a "period" $z \in Z_{A(\bar{\Omega})}G$ ($\bar{\Omega}$ the Dedekind completion of Ω) analogous to the map $\beta z = \beta + 1$ in the example. Periodic groups are shown to be l -simple, and more examples of them are constructed.

Transitivity guarantees that the "configuration" of orbits of G_α is independent of α , so that we may speak of the *configuration* of G (defined more precisely later). There is appreciable interplay between this configuration and other properties of G . For example, o -2-transitive groups are characterized by having only one positive orbit, and regular groups by having configurations consisting entirely of fixed points.

For periodically o -primitive groups, the period z is the unique o -permutation of $\bar{\Omega}$ such that for every $\beta \in \Omega$, βz is the sup of the first positive orbit of G_β . $(\beta z)g = (\beta g)z$ for all $\beta \in \Omega$, $g \in G$, and in fact z generates $Z_{A(\bar{\Omega})}G$. This periodicity is of paramount importance. For example, it guarantees that the action of $g \in G$ on any long orbit of G_α determines its action on all of Ω .

Transitive l -subgroups of $A(\Omega)$ have been studied from a lattice-ordered group (l -group) orientation by Holland [5, 6, 7], Lloyd [10, 11], Sik [15], and McCleary [12, 13]. Holland showed that every l -group

is l -isomorphic to a subdirect product of transitive l -permutation groups [5]. A nonlattice point of view has been taken by Holland and McCleary [8, 14], where it was shown that every transitive ordered permutation group can be embedded in the generalized ordered wreath product of its o -primitive "components" (an important motivation for the present paper); and by G. Higman [4] and Wielandt [17, §6]. The concept of configuration is a refinement of the concept of rank in [3].

The generalization to partially ordered Ω requires very little additional work, but it is less intuitive than the totally ordered case and the reader will not lose much if he assumes that Ω is totally ordered, or even that G is an l -permutation group, as we have done in this introduction.

2. **Coherent o -permutation groups.** Let Ω be a partially ordered set (po -set) containing more than one point. Points of Ω will be denoted by lower case Greek letters; subsets, by upper case Greek letters; and permutations, by lower case Roman letters. The image of $\beta \in \Omega$ under the permutation f will be denoted by βf , so that if g is also a permutation, $\beta(fg) = (\beta f)g$.

An *order-preserving permutation* (o -permutation, *automorphism*) of Ω is a permutation f such that for $\beta, \gamma \in \Omega$, $\beta < \gamma$ iff $\beta f < \gamma f$. We define $f \leq g$ iff $\beta f \leq \beta g$ for all $\beta \in \Omega$, making the group $A(\Omega)$ of all permutations of Ω into a partially ordered group (po -group). If Ω is totally ordered, f is an o -permutation provided only that $\beta < \gamma$ implies $\beta f < \gamma f$. In this case $A(\Omega)$ is an l -group, with $\beta(f \vee g) = \max \{\beta f, \beta g\}$ and $\beta(f \wedge g) = \min \{\beta f, \beta g\}$; and G is said to be an l -permutation group if it is an l -subgroup of $A(\Omega)$, i.e., a subgroup which is also a sublattice. Standard results about po -groups and l -groups can be found in [2], but we shall make minimal use of them.

Our o -permutation group G will always be assumed to be a transitive subgroup of $A(\Omega)$ (i.e., $\beta, \gamma \in \Omega$ implies $\beta g = \gamma$ for some $g \in G$). Thus Ω must be homogeneous; and if ordered nontrivially ($\beta < \gamma$ for some $\beta, \gamma \in \Omega$), it must be infinite. Furthermore, we shall always assume that if $\beta < \gamma \in \Omega$, there exists $1 < g \in G$ such that $\beta g = \gamma$. (This property implies its dual, which states that if $\beta > \gamma$, there exists $1 > g \in G$ such that $\beta g = \gamma$; and implies that if $\beta f < \gamma$, $f \in G$, then there exists $g \in G$ such that $\beta g = \gamma$ and $g > f$). Transitive groups that satisfy this property will be called *coherent*. Of course, if Ω is totally ordered, transitivity need not be separately assumed. Transitive l -permutation groups are coherent, for if $\beta < \gamma$ and $\beta g = \gamma$, then also $\beta(g \vee 1) = \gamma$. However, the group in Example 7 is not coherent. If Ω is trivially ordered, $A(\Omega)$ is just the symmetric group $S(\Omega)$, and is itself trivially ordered; and its transitive subgroups are automatically coherent.

B is a convex subset (segment) of a po -set A if $b_1 \leq a \leq b_2, b_1, b_2 \in B, a \in A$ implies $a \in B$. If C and D are any subsets of A , we define $C \leq D$ iff $c \leq d$ for some $c \in C, d \in D$. If A is totally ordered, and C and D are nonvoid disjoint segments of Ω , then $C < D$ iff $c < d$ for all $c \in C, d \in D$.

If (G, Ω) is a transitive (but not necessarily coherent) o -permutation group, let $R(G_\alpha)$ designate $\{G_\alpha g \mid g \in G\}$, ordered as above to give the usual partial ordering on the collection of right cosets of a convex subgroup of a po -group. As with nonordered transitive permutation groups, we make G act faithfully on $R(G_\alpha)$ by defining $(G_\alpha g) = G_\alpha(gk), g, k \in G$. Here we obtain an o -permutation group.

An o -isomorphism from one o -permutation group (G, Ω) onto another (K, Σ) consists of a po -set isomorphism θ_Ω from Ω onto Σ and a po -group isomorphism θ_G from G onto K such that for all $\omega \in \Omega, g \in G, (\omega g)\theta_\Omega = (\omega\theta_\Omega)(g\theta_G)$. The importance of coherence is explained by

THEOREM 1. *Let (G, Ω) be a transitive o -permutation group and let $\alpha \in \Omega$. Then G is coherent if and only if the correspondence $\alpha g \leftrightarrow G_\alpha g$ between Ω and $R(G_\alpha)$ and the identity map on G furnish an o -isomorphism between (G, Ω) and $(G, R(G_\alpha))$.*

Proof. Suppose that G is coherent. $\alpha g_1 = \alpha g_2$ iff $g_1 g_2^{-1} \in G_\alpha$ iff $G_\alpha g_1 = G_\alpha g_2$, so we have a one-to-one correspondence between Ω and $R(G_\alpha)$. $\alpha g_1 \leq \alpha g_2$ iff $\alpha g_1 k = \alpha g_2$ for some $1 \leq k \in G$ (by coherence) iff $G_\alpha g_1 k = G_\alpha g_2$ (for some $1 \leq k \in G$) iff $G_\alpha g_1 \leq G_\alpha g_2$, so the correspondence is an o -isomorphism. For $h \in G, (\alpha g)h = \alpha(gh) \leftrightarrow G_\alpha(gh) = (G_\alpha g)h$. This establishes the o -permutation group isomorphism. The converse is clear.

G is regular if it is transitive and $G_\alpha = \{1\}$.

COROLLARY 2. *Let G be regular. Then G is coherent if and only if (G, Ω) is o -isomorphic to the right regular representation of G . In particular, the right regular representation of G is coherent.*

3. The configuration of an o -permutation group. There will usually be one (arbitrary) point α in Ω on which our attention will be especially focused. The orbit of G_α which contains δ is $\delta G_\alpha = \{\delta h \mid h \in G_\alpha\}$. $\alpha G_\alpha = \{\alpha\}$. If δG_α is not trivially ordered, it is infinite. The orbits of G_α partition Ω . In general, the orbits of G_α need not be convex (Examples 3 and 6), although of course they are convex if Ω is trivially ordered. We also have

PROPOSITION 3. *If G is a transitive l -subgroup of $A(\Omega), \Omega$ totally ordered, then the orbits of G_α are convex.*

Proof. Suppose $\beta \leq \gamma \leq \delta$ and $\beta h = \delta$ for some $h \in G_\alpha$. By transitivity, $\beta g = \gamma$ for some $g \in G$. Let $f = (h \vee 1) \wedge (g \vee 1)$. Then $\beta f = \gamma$. Since $1 \leq f \leq h \vee 1 \in G_\alpha$, the convexity of G_α implies that $f \in G_\alpha$.

To escape having to assume that the orbits of G_α are convex, we shall "enlarge" them to convex sets. The *convexification* $\text{Conv}(\Delta)$ of $\Delta \subseteq \Omega$ is $\{\xi \in \Omega \mid \delta_1 \leq \xi \leq \delta_2 \text{ for some } \delta_1, \delta_2 \in \Delta\}$. If Δ is an orbit of G_α , we shall call $\text{Conv}(\Delta)$ an *orbital* of G_α . Of course, if the orbits of G_α are convex, the concepts of "orbital" and "orbit" coincide. If Γ is an orbital of G and $\gamma \in \Gamma$, then the orbital $\text{Conv}(\gamma G_\alpha)$ of G_α determined by γ is Γ . The orbitals of G_α partition Ω into convex subsets. The set of orbitals of G_α is partially ordered; and is totally ordered if Ω is totally ordered. Two orbits in different orbitals are related as are their orbitals; and two orbits in the same orbital are of course each less than or equal to the other.

Those orbitals of G_α which are strictly greater than $\{\alpha\}$ will be called *positive*; those strictly less than $\{\alpha\}$, *negative*. All points in a positive (negative) orbital are strictly greater than (less than) α . No orbital is both positive and negative; and if Ω is totally ordered, every orbital except $\{\alpha\}$ is one or the other. These remarks apply also to orbits of G_α .

We define for each orbit Δ a *paired orbit* $\Delta' = \Delta'^\alpha = \{\alpha g \mid \alpha \in \Delta g\}$. (The notation Δ' will always refer to pairings with respect to the point denoted by the letter α). It is shown in [18, § 16] that Δ' is indeed an orbit of G_α ; that the map $\Delta \rightarrow \Delta'$ is one-to-one from the set of orbits of G_α onto itself; and that $\Delta'' = \Delta$. $\alpha g \in \Delta'$ iff $\alpha \in \Delta g$, and if $\alpha \in \Delta g$, then $\Delta' = (\alpha g)G_\alpha$.

PROPOSITION 4. *Let (G, Ω) be a coherent o-permutation group. The map $\Delta \rightarrow \Delta'$ is an o-anti-automorphism of the set of orbits of G_α . Since $\{\alpha\}$ is self-paired, the appropriate restriction provides an o-anti-isomorphism from the set of positive orbits of G_α onto the set of negative orbits. If Ω is totally ordered, only $\{\alpha\}$ is self-paired.*

Proof. Use coherence.

A subset Δ of Ω will be called α -full if it contains each orbit of G_α that it meets, i.e., if it is a union of orbits of G_α . Thus the α -full sets are precisely those sets Δ such that $\Delta h = \Delta$ for each $h \in G_\alpha$. We obtain a canonical correspondence between the α -full subsets of Ω and the subsets of the set of orbits of G_α by letting the α -full set Δ correspond to the set of orbits contained in Δ . We shall frequently make the tempting identification and refer to α -full sets as being subsets of the set of orbits of G_α . A convex α -full set Δ is a union of orbitals and is a convex subset of the *po*-set of orbitals of G_α .

Now we extend the concept of pairings to α -full sets. If Δ is α -full, we define Δ' to be $\{\alpha g \mid \alpha \in \Delta g\} = \cup \{\Gamma' \mid \Gamma \text{ is an orbit of } G_\alpha \text{ and } \Gamma \subseteq \Delta\}$. If $\{\Delta_i \mid i \in I\}$ is any family of α -full sets, then $\cup \{\Delta_i \mid i \in I\}$ is α -full and is paired with $\cup \{\Delta'_i \mid i \in I\}$; and similarly for intersections. If $\Delta'^\alpha = \Delta$, we say Δ is *symmetric with respect to* α .

PROPOSITION 5. *If Δ is an α -full set, then $\text{Conv}(\Delta)$ is α -full and $[\text{Conv}(\Delta)]' = \text{Conv}(\Delta')$. If Δ is already convex, so is Δ' . If Δ is symmetric with respect to α , so is $\text{Conv}(\Delta)$.*

Proof. $\Delta \rightarrow \Delta'$ is an α -anti-automorphism.

Since an orbital Δ of G_α is always α -full, the last proposition implies that Δ' is also an orbital, and that it contains precisely those orbits paired with orbits contained in Δ .

THEOREM 6. *Proposition 4 holds for orbitals of G_α .*

If $\beta G_\alpha = \{\beta\}$, β is said to be a *fixed point* of G_α . If not, βG_α is a *long orbit* of G_α and $\text{Conv}(\beta G_\alpha)$ a *long orbital*. Unless it is trivially ordered, a long orbit(al) must be infinite. We make six definitions:

$FxG_\alpha = \{\beta \in \Omega \mid \beta \text{ is a fixed point of } G_\alpha\}$.

$SFxG_\alpha = \{\beta \in \Omega \mid \beta, \beta' \in FxG_\alpha\}$.

$WFxG_\alpha = \{\beta \in \Omega \mid \beta \in FxG_\alpha, \text{ but } \beta' \text{ is a long orbit}\}$.

$LnG_\alpha = \{\Delta \subseteq \Omega \mid \Delta \text{ is a long orbit of } G_\alpha\}$.

$SLnG_\alpha = \{\Delta \subseteq \Omega \mid \Delta, \Delta' \in LnG_\alpha\}$.

$WLnG_\alpha = \{\Delta \subseteq \Omega \mid \Delta \in LnG_\alpha, \text{ but } \Delta' \text{ is a fixed point}\}$.

Points in $SFxG_\alpha$ will be called *strongly fixed*; points in $WFxG_\alpha$, *weakly fixed*; orbits in $SLnG_\alpha$, *strongly long*; and orbits in $WLnG_\alpha$, *weakly long*. XG_α will be a variable which can take on as values each of these six sets. Each XG_α is α -full and thus may be thought of either as a subset of the set of orbits of G_α or as a subset of Ω . Clearly Ω is partitioned by FxG_α and LnG_α . In turn, FxG_α is partitioned by $SFxG_\alpha$ and $WFxG_\alpha$; and LnG_α , by $SLnG_\alpha$ and $WLnG_\alpha$. $SFxG_\alpha$ and $SLnG_\alpha$ are self-paired; and $WFxG_\alpha$ is paired with $WLnG_\alpha$.

PROPOSITION 7. $\beta \in SFxG_\alpha$ iff $G_\beta = G_\alpha$.
 $\beta \in WFxG_\alpha$ iff $G_\beta \supset G_\alpha$.
 $\beta \in WLnG_\alpha$ iff $G_\beta \subset G_\alpha$.
 $\beta \in SLnG_\alpha$ iff $G_\beta \not\subseteq G_\alpha$ and $G_\beta \not\supseteq G_\alpha$.

Proof. Clearly $\beta \in FxG_\alpha$ iff $G_\beta \cong G_\alpha$. Pick $g \in G$ such that $\beta g = \alpha$ and thus $\alpha g \in (\beta G_\alpha)'$. Then $G_\beta \subseteq G_\alpha$ iff $\alpha \in FxG_\beta$ iff $\alpha g \in FxG_\alpha$ iff $(\beta G_\alpha)'$ is a fixed point of G_α . The proposition follows.

We shall say that G is *balanced* if $WFxG_\alpha$ is the empty set \square (iff $WLnG_\alpha = \square$ iff $SFxG_\alpha = FxG_\alpha$ iff $SLnG_\alpha = LnG_\alpha$). By Proposition 7, G fails to be balanced iff G_α is properly contained in one of its conjugates. It follows that finite groups are balanced; in fact, paired orbits have equal cardinalities [18, Theorem 16.3]. Examples can be constructed of l -permutation groups (G, Ω) , Ω totally ordered, which are not balanced.

Proposition 5 yields

PROPOSITION 8. *Any orbit of G_α which is not strongly long is convex. Hence if two different orbits of G_α lie in the same orbital of G_α , both are strongly long.*

We now apply the XG_α terminology to *orbitals* of G_α , being assured that an orbital $\text{Conv}(\Delta)$ is contained in that XG_α containing the orbit Δ .

The α -*configuration* of G is defined to be the po -set (o -set if Ω is totally ordered) of orbitals of G_α , partitioned into $SFxG_\alpha$, $WFxG_\alpha$, $SLnG_\alpha$, and $WLnG_\alpha$, with the point α distinguished; together with the involution $\Delta \rightarrow \Delta'$. α is called the *origin*. (Actually, the α -configuration is completely determined by the po -set of orbitals, the subset of fixed points, the origin, and the involution.) We want to show that this configuration is actually independent of α . By an o -isomorphism from the α -configuration of (G, Ω) onto the β -configuration of (K, Σ) , we mean a po -set isomorphism ψ from the po -set orbitals of G_α onto that of K_β such that $(XG_\alpha)\psi = XK_\beta$ for each XG_α , $\{\alpha\}\psi = \{\beta\}$, and $(\Delta\psi)'\beta = (\Delta'\alpha)\psi$ for all orbitals Δ of G_α . When there is such an o -isomorphism, we shall say that the two configurations are "the same configuration".

For any $f \in G$, an o -automorphism of (G, Ω) is provided by θ_α , defined by $\omega\theta_\alpha = \omega f$, and θ_α , defined by $g\theta_\alpha = f^{-1}gf$. Hence the map $\Delta \rightarrow \Delta f$ is an o -isomorphism from the α -configuration onto the β -configuration. Moreover, if $\alpha f_1 = \alpha f_2$, with $f_1, f_2 \in G$, then $f_1 f_2^{-1} \in G_\alpha$, so for each α -full set Δ , $\Delta f_1 f_2^{-1} = \Delta$ and thus $\Delta f_1 = \Delta f_2$. This proves the fundamental

THEOREM 9. *Let G be a coherent subgroup of $A(\Omega)$. Let $\alpha, \beta \in \Omega$ and pick $f \in G$ such that $\alpha f = \beta$. Then $\Delta \rightarrow \Delta f$ furnishes a canonical o -isomorphism (independent of the choice of f) from the α -configuration onto the β -configuration. The canonical o -isomorphism from the α -configuration onto the β -configuration, followed by that from the β -configuration onto the γ -configuration, yields the canonical o -isomorphism from the α -configuration onto the γ -configuration.*

Hence we may speak of the *configuration* of G without reference to a particular point of Ω . Obviously if two o -permutation groups are o -isomorphic, they have the same configuration. Of course we can state a similar definition of configuration in terms of orbits rather than orbitals. Two groups having the same orbit configurations necessarily have the same orbital configurations; but not conversely (Examples 2 and 3). However, the orbit configuration is determined by the orbital configuration together with the number of orbits in each orbital. When we speak of configurations, we shall mean *orbital* configurations unless specified otherwise.

Two distinct points $\beta < \gamma$ of Ω have three possible relationships: $\beta < \gamma$, $\beta > \gamma$, and β incomparable with γ . G is o -2-transitive if for any $\beta, \gamma, \sigma, \tau \in \Omega$ such that β and γ are related in the same way as are σ and τ , there exists $g \in G$ such that $\beta g = \sigma$ and $\gamma g = \tau$. If G is o -2-transitive, G must have precisely one positive orbit and precisely one negative orbit (unless Ω is trivially ordered); and precisely one incomparable orbit (unless Ω is totally ordered). Conversely, it is easy to see that if G has such a configuration, G is o -2-transitive. Thus o -2-transitive groups can be characterized in terms of *orbit* configurations; though not in terms of orbital configurations (Example 3), except among the class of l -permutation groups.

We shall be interested also in those groups whose *orbital* configurations are the same as the *orbit* configurations described above for o -2-transitive groups. These groups are characterized by the property that for any $\beta, \gamma, \sigma, \tau \in \Omega$ such that β and γ are related as are σ and τ , there exists $g_1 \in G$ such that $\beta g_1 = \sigma$ and $\gamma g_1 \leq \tau$; and $g_2 \in G$ such that $\beta g_2 = \sigma$ and $\gamma g_2 \geq \tau$. Such groups will be called o -2-semitransitive. An o -2-semitransitive l -permutation group is automatically o -2-transitive.

The regular groups can of course be characterized as those whose configurations consist entirely of (strongly) fixed points.

Groups lying between the extremes of o -2-transitivity and regularity can be found among the examples at the end of the paper. See especially Examples 5 and 8. When Ω is totally ordered, the o -anti-isomorphism $\Delta \rightarrow \Delta'$ reduce the problem of determining the o -set of all orbitals to that of determining the o -set of positive orbitals. It can be shown that every o -set occurs as the o -set of positive orbitals for some transitive $(A(\Omega), \Omega)$.

If Δ is an orbit of G_α , the canonically corresponding orbit of G_β will be denoted by Δ_β . In particular, $\Delta_\alpha = \Delta$. Δ_β is to be thought of as "the Δ orbit of G_β ". Of course, $(\Delta_\alpha)f = \Delta_{\alpha f}$. Since $\Delta \rightarrow \Delta f$ also yields a canonical isomorphism from the set of α -full sets onto the set of (αf) -full sets, we may apply the same notation to α -full sets Δ_α , and in particular to orbitals of G_α .

PROPOSITION 10. *If $\alpha g \in SF\alpha G_\alpha$, $g \in G$, then for each orbit(al) Δ of G_α , Δg is another orbit(al) of G_α , and it lies in the same XG_α as Δ .*

Proof. Proposition 7.

4. **O-blocks.** By *o-block* of an *o-permutation group* (G, Ω) , we mean a *convex* subset $\square \neq \Delta \subseteq \Omega$ having the property that for any $g \in G$, $\Delta g = \Delta$ or $\Delta g \cap \Delta = \square$. If the convexity requirement is removed, one has simply a *block* as defined in [18, §6]. Of course, these two concepts coincide when Ω is trivially ordered. The intersection of any collection of *o-blocks* is an *o-block* (provided it is not empty) and the union of any tower of *o-blocks* is an *o-block*. If Δ is an *o-block*, the *o-block system* $\tilde{\Delta}$ is the *po-set* (*o-set* if Ω is totally ordered) of translates Δg ($g \in G$) of Δ . Since G is transitive, the *o-block systems* of G correspond to the convex G -congruences, where a G -congruence is said to be *convex* if its congruence classes are convex.

We partially order the blocks containing α by inclusion, obtaining a complete lattice, of which the *o-blocks* containing α form a complete sublattice; and similarly for the subgroups of G containing G_α .

THEOREM 11. *Let (G, Ω) be a coherent o-permutation group. In the well known o-correspondence $\Delta \rightarrow \{g \in G \mid \Delta g = \Delta\}$ and $C \rightarrow \alpha C$ between the lattice of blocks containing α and the lattice of subgroups containing G_α , the convex subgroups C correspond precisely to the o-blocks Δ .*

Proof. Clearly if Δ is convex, $\{g \in G \mid \Delta g = \Delta\}$ is convex. Now assume that C is convex. Suppose $\alpha c \leq \beta \leq \alpha d$, $c, d \in C$. Pick $f \in G$ such that $\alpha f = \beta$. Use coherence to pick $s \in G$ such that $\alpha s = \alpha d$ and $f \leq s$. Since $d \in C$ and $sd^{-1} \in G_\alpha \subseteq C$, $s \in C$. Similarly, pick $t \in C$ such that $t \leq f$. Since C is convex, $t \leq f \leq s$ implies $f \in C$, so that $\beta = \alpha f \in \alpha C$. Therefore αC is convex. This result fails without coherence (Example 7).

We may make a complete lattice of the set of block systems of G by defining $\tilde{\Gamma} \leq \tilde{\Delta}$ iff $\Gamma \subseteq \Delta$, where Γ and Δ are the blocks in $\tilde{\Gamma}$ and $\tilde{\Delta}$ which contain α . Obviously the definition is independent of the choice of α . The set of *o-block systems* forms a complete sublattice. It is proved in [8, Theorem 3] that if Ω is totally ordered, the lattice of *o-block systems* is also totally ordered. Thus Theorem 11 gives us

COROLLARY 12. *The convex subgroups of G which contain G_α are totally ordered under inclusion.*

For the special case of *l-permutation groups*, this was proved by Holland [5]. His result mentioned only the convex prime *l-subgroups*

containing G_α , but since G_α is prime, every subgroup containing it must automatically be a prime l -subgroup, and thus the two results coincide.

PROPOSITION 13. *A block Δ of G which contains α must be α -full and symmetric with respect to α .*

THEOREM 14. *Let G be a coherent subgroup of $A(\Omega)$, and let $\Delta = \Delta_\alpha$ be a convex α -full set. Then $\Gamma = \{\beta \in \Omega \mid \Delta_\beta = \Delta_\alpha\}$ is a (symmetric) o -block of G .*

Proof. $C = \{g \in G \mid \Delta g = \Delta\}$ is a convex subgroup of G containing G_α . But $\Gamma = \alpha C$, which is an o -block of G by Theorem 11.

It is immediate from the proof of Theorem 14 that even if Δ is not convex, Γ is still a block of G . This can also be deduced from the statement of the theorem. For if we throw away the order on Ω , leaving Ω trivially ordered and G coherent, then Δ becomes convex, so by the theorem, Γ is a block of G . Similar remarks apply to many of the theorems to come.

THEOREM 15. *Let G be a coherent subgroup of $A(\Omega)$. If Δ is an α -full o -block of G , then Δ' is also an (α -full) o -block of G , and $\{\beta \in \Omega \mid \Delta_\beta = \Delta_\alpha\}$ is the translate of Δ' which contains α .*

Proof. Let Γ be the o -block $\{\beta \in \Omega \mid \Delta_\beta = \Delta_\alpha\}$. Pick $f \in G$ such that $\alpha \in \Delta f$. Then Γf , also an o -block, is equal to $\{\eta \in \Omega \mid \Delta_\eta = \Delta f\} = \{\eta \in \Omega \mid \alpha \in \Delta_\eta\}$ (because Δ is a block) $= \{\alpha g \mid \alpha \in \Delta_{\alpha g} = \Delta_\alpha g\} = \Delta'$.

COROLLARY 16. *Let Δ be a weakly long orbit of G_α . Then Δ is an o -block of G . Indeed, if $\alpha g \neq \alpha, g \in G$, then $\Delta g \cap \Delta = \square$.*

Proof. Theorems 15 and 14. Thus for an α -full o -block Δ , Δ' need not lie in the same o -block system as Δ .

When Ω is totally ordered, we may complete Ω by Dedekind cuts and consider Ω to be a subset of its Dedekind completion $\bar{\Omega}$ (without end points). Each $f \in A(\Omega)$ can be extended to $f \in A(\bar{\Omega})$ by defining $\bar{\omega}f$ to be $\sup\{\beta f \mid \beta \in \Omega, \beta \leq \bar{\omega}\}$. $A(\Omega)$ is an l -subgroup of $A(\bar{\Omega})$, but in general is not transitive even on $\bar{\Omega} \setminus \Omega$. A point $\bar{\omega} \in \bar{\Omega}$ is α -full if it is fixed by G_α . Equivalently, $\bar{\omega}$ is α -full if it is the sup (inf) of an α -full segment of Ω . If $\bar{\omega} \in \Omega$, then $\bar{\omega}$ is α -full iff $\bar{\omega} \in \text{Fix } G_\alpha$. For any α -full point $\bar{\omega}_\alpha$, and for any $g \in G, \bar{\omega}_{\alpha g} = \bar{\omega}_\alpha g$ is the (αg) -full point canonically corresponding to $\bar{\omega}_\alpha$.

PROPOSITION 17. *Suppose that Ω is totally ordered and that $\bar{\omega}_\alpha$ is*

an α -full point. Then $\{\beta \in \Omega \mid \bar{\omega}_\beta = \bar{\omega}_\alpha\}$ is an α -block of G .

Proof. $\{\eta \in \Omega \mid \eta \leq \bar{\omega}_\alpha\}$ is an α -full segment of Ω . Apply Theorem 14.

LEMMA 18. Suppose Ω is totally ordered. Let Δ be an α -full set. If $\alpha g \geq \alpha$, then $(\inf \Delta)g \geq \inf \Delta$ and $(\sup \Delta)g \geq \sup \Delta$.

Proof. Pick $1 \leq k \in G$ such that $\alpha k = \alpha g$. Since Δ is α -full, $\Delta g = \Delta k$.

It is easily checked that

LEMMA 19 ([7, Lemma 3]). Let $\alpha \in \Delta \subseteq \Omega$. Suppose that $\Delta g = \Delta$ for each $g \in G$ such that $\alpha g \in \Delta$. Then Δ is a block of G .

LEMMA 20. Suppose that $\alpha \in \Delta \subseteq \Omega$, Ω totally ordered, and that Δ is convex, α -full, and symmetric with respect to α . Let Π be any cofinal subset of Δ . Then Δ is an α -block of G provided only that $\alpha g \in \Pi, g \in G$, implies $\inf \Delta g \succ \inf \Delta$ and $\sup \Delta g \succ \sup \Delta$.

Proof. By the first lemma, we see first that $\Delta g = \Delta$ when $\alpha \leq \alpha g \in \Pi$; and next that $\Delta g = \Delta$ when $\alpha \leq \alpha g \in \Delta$. In view of the second lemma, the conclusion follows from the symmetry of Δ .

THEOREM 21. Let G be a coherent subgroup of $A(\Omega)$, Ω totally ordered. Suppose G has a (long) orbital Δ cofinal with Ω , so that Δ' is a (long) orbital coinitial with Ω . Then $\{\beta \in \Omega \mid \Delta' < \beta < \Delta\}$ is an α -block of G .

Proof. By transitivity, terminal orbitals must be long. Now let Π be the α -full set $\Gamma = \{\beta \in \Omega \mid \Delta' < \beta < \Delta\}$ and let $\bar{\sigma} = \sup \Gamma$. We show first that if $\alpha < \alpha g \in \Gamma, g \in G$, then $\bar{\sigma} g \succ \bar{\sigma}$. For suppose $\bar{\sigma} g > \bar{\sigma}$. Pick $h \in G$ such that $\bar{\sigma} h < \alpha$. Since Δ is cofinal with Ω , we can pick $\delta \in \Delta$ such that $\delta h > \bar{\sigma}$. Now pick $k \in G_\alpha$ such that $(\bar{\sigma} g)k > \delta$. Since $k \in G_\alpha$ and Γ is α -full, $(\alpha g)k \in \Gamma$, so that $\alpha g k \leq \bar{\sigma}$. Since $(\alpha g k)h \leq \bar{\sigma} h < \alpha$, we can use coherence to pick $h \leq f \in G$ such that $(\alpha g k)f = \alpha$. But $\bar{\sigma} g k f \geq \bar{\sigma} g k h > \delta h > \bar{\sigma}$, contradicting the fact that $\bar{\sigma}$ is α -full. Therefore $\bar{\sigma} g \succ \bar{\sigma}$ when $\alpha < \alpha g \in \Gamma$. Similarly, $(\inf \Gamma)f \prec \inf \Gamma$ when $\alpha > \alpha f \in \Gamma$, and thus since Γ is symmetric, $(\inf \Gamma)g \succ \inf \Gamma$ when $\alpha < \alpha g \in \Gamma$. By the last lemma, Γ is an α -block of G .

In generalizations of theorems about finite permutation groups, FxG_α often must be expressed as $SFxG_\alpha$ ($= FxG_\alpha$ if G is finite). For example:

THEOREM 22. *Let (G, Ω) be a coherent o -permutation group. Then $SFxG_\alpha$ is a block of G .*

Proof. $SFxG_\alpha$ is α -full, so $(SFxG_\alpha)g = SFxG_{\alpha g}$. In view of Proposition 7, this says that $\{\beta \in \Omega \mid G_\beta = G_\alpha\}g = \{\gamma \in \Omega \mid G_\gamma = G_{\alpha g}\}$, which is equal to $SFxG_\alpha$ if $G_{\alpha g} = G_\alpha$, and does not meet $SFxG_\alpha$ otherwise.

5. **O-primitive groups.** Following Holland's definition for l -groups [7], we define a coherent subgroup G of $A(\Omega)$, Ω partially ordered, to be *o -primitive* if G has no o -blocks except Ω and the singletons $\{\omega\}$. Theorem 11 establishes Holland's result (obtained in essentially the same way) that G is o -primitive if and only if G_α is a maximal proper convex subgroup of G . O -permutation groups which are primitive are *a fortiori* o -primitive. On the other hand, $A(I)$, I the integers, is o -primitive, but not primitive.

PROPOSITION 23. *Let (G, Ω) be a coherent o -permutation group, Ω totally ordered. If G is o -2-semitransitive, it is o -primitive. If G is o -2-transitive, it is primitive.*

An o -group K is *Archimedean* if for any $1 < k, f \in K, f < k^n$ for some positive integer n ; i.e., if K contains no proper convex subgroups. K is Archimedean iff K is isomorphic as an o -group to an o -subgroup of the additive reals [2, p. 45].

PROPOSITION 24. *Suppose that (G, Ω) is regular, with Ω totally ordered. Then (G, Ω) is o -primitive iff G is Archimedean.*

Proof. By Theorem 11, since $G_\alpha = \{1\}$.

This proposition almost characterizes the o -primitive regular groups in terms of their configurations. Unfortunately, it is possible for an Archimedean o -group (the rationals) to be isomorphic as an o -set to a non-Archimedean o -group ($\overleftarrow{Q} \times I, Q$ the rationals, I the integers). This is the reason for the word "almost".

Among o -primitive groups on totally ordered sets Ω , there are thus two classes which lie at opposite extremes in terms of the amount of movement possible within G_α : the Archimedean regular groups, which we have almost characterized in terms of their configurations; and the o -2-semitransitive groups, which we have completely characterized in terms of their configurations. The remaining o -primitive groups will be discussed in detail in §7. For now, we apply §4 to o -primitive groups in general.

If $\Delta \subseteq \Omega$ and $\beta, \gamma \in \Omega$, we say that β and γ can be *separated* by

Δ if some translate $\Delta g (g \in G)$ of Δ contains precisely one of β and γ . An orbit $\bar{\omega}G$ of G is dense in $\bar{\Omega}$ if it meets every nontrivial segment of $\bar{\Omega}$. Of course, $\bar{\omega}G = \Omega$ if $\bar{\omega} \in \Omega$, and $\bar{\omega}G \cap \Omega = \square$ if $\bar{\omega} \in \bar{\Omega} \setminus \Omega$.

THEOREM 25. *Let (G, Ω) be a coherent o -permutation group. The following are equivalent (except that if Ω is not totally ordered, only the first three make sense):*

- (i) G is o -primitive.
- (ii) For every segment $\square \neq \Delta \subset \Omega$, any $\beta \neq \gamma \in \Delta$ can be separated by Δ .
- (iii) For every α -full segment $\square \neq \Delta_\alpha \subset \Omega$, $\Delta_\beta \neq \Delta_\gamma$ for $\beta \neq \gamma$ ($\alpha, \beta, \gamma \in \Omega$).
- (iv) For every α -full point $\bar{\omega}_\alpha \in \bar{\Omega}$, $\bar{\omega}_\beta \neq \bar{\omega}_\gamma$ for $\beta \neq \gamma$ ($\alpha, \beta, \gamma \in \Omega$).
- (v) For every $\bar{\omega} \in \bar{\Omega}$, $\bar{\omega}G$ is dense in $\bar{\Omega}$.

Proof. It is clear that each of these conditions implies (i). Now suppose that G is o -primitive. If Δ is a segment, $\square \neq \Delta \subset \Omega$, then a convex G -congruence is given by the relation $\beta \equiv \gamma$ iff β and γ cannot be separated by Δ ; and since some pairs $\beta \neq \gamma \in \Omega$ can be separated by Δ , every pair can, so that (ii) holds. For (v), if $\bar{\Gamma}$ were a nontrivial segment of $\bar{\Omega}$ which did not meet $\bar{\omega}G$, then for $\beta \neq \gamma \in \bar{\Gamma} \cap \Omega$ and $\Delta = \{\omega \in \Omega \mid \omega < \bar{\omega}\}$, β and γ could not be separated by Δ . For (iii), we use Theorem 14; and for (iv), Proposition 17. For Ω totally ordered and G an l -subgroup of $A(\Omega)$, the equivalence of (i), (ii), and (v) was shown by Holland [7, Theorem 2]. For Ω trivially ordered, the equivalence of (i) and (ii) was shown by Wielandt [17, Theorem 7.12].

THEOREM 26. *Let (G, Ω) be o -primitive. Then G is balanced and FxG_α is a block of G .*

Proof. Since weakly long orbits are o -blocks, G is balanced, so $FxG_\alpha = SFxG_\alpha$ is a block.

6. Centralizers. In Example 8, the map $z: \Omega \rightarrow \Omega$ given by $\beta z = \beta + 1$ lies in the centralizer $Z_{A(\Omega)}G$ of G in $A(\Omega)$. This phenomenon will be of paramount importance in the study of o -primitive groups. Accordingly, we devote this section to the study of centralizers.

When Ω is totally ordered, we shall be interested also in the centralizer of G in $A(\bar{\Omega})$. We define $\bar{F}xG_\alpha = \{\bar{\omega} \in \bar{\Omega} \mid \bar{\omega}G_\alpha = \bar{\omega}\} = \{\bar{\omega} \in \bar{\Omega} \mid G_{\bar{\omega}} \cong G_\alpha\}$ and $\bar{S}\bar{F}xG_\alpha = \{\bar{\omega} \in \bar{\Omega} \mid \bar{\omega}G_\alpha = \bar{\omega} \text{ and } \alpha G_{\bar{\omega}} = \alpha\} = \{\bar{\omega} \in \bar{\Omega} \mid G_{\bar{\omega}} = G_\alpha\}$. Points in these two sets are α -full. By Proposition 7, $\bar{F}xG_\alpha \cap \Omega = FxG_\alpha$ and $\bar{S}\bar{F}xG_\alpha \cap \Omega = SFxG_\alpha$. In the two lemmas which follow, if Ω is not totally ordered, one replaces $\bar{\Omega}$ by Ω , $\bar{F}xG_\alpha$ by FxG_α ,

and $\bar{S}FxG_\alpha$ by $SFxG_\alpha$.

LEMMA 27. *Let $z: \Omega \rightarrow \bar{\Omega}$ be a function which centralizes G , and let $\bar{\omega}_\alpha = \alpha z$. Then $\bar{\omega}_\alpha \in \bar{F}xG_\alpha$, and for all $\beta \in \Omega$, $\beta z = \bar{\omega}_\beta$. If z is one-to-one, $\bar{\omega}_\alpha \in \bar{S}FxG_\alpha$.*

Proof. For any $g \in G$, $\alpha z g = \alpha g z$; so that $\alpha z \in \bar{F}xG_\alpha$, and $\alpha z \in \bar{S}'FxG_\alpha$ if z is one-to-one. Now let $\beta \in \Omega$ and pick $k \in G$ such that $\alpha k = \beta$. Then $\beta z = \alpha k z = \alpha z k = \omega_\alpha k = \omega_{\alpha k} = \omega_\beta$.

COROLLARY 28. $Z_{S(\Omega)}G = Z_{A(\Omega)}G$, where $S(\Omega)$ is the symmetric group on Ω .

Proof. If $\bar{\omega}_\alpha \in \bar{F}xG_\alpha$, then for any $\alpha \leq \beta \in \Omega$, $\bar{\omega}_\alpha \leq \bar{\omega}_\beta$ by coherence.

LEMMA 29. *Let $\bar{\omega}_\alpha \in \bar{F}xG_\alpha$. Define $z: \bar{\Omega} \rightarrow \bar{\Omega}$ by setting $\beta z = \bar{\omega}_\beta$ for $\beta \in \Omega$, and $\bar{\gamma} z = \sup\{\beta z \mid \beta \leq \bar{\gamma}\}$ for $\bar{\gamma} \in \bar{\Omega}$. Then z centralizes G . If $\bar{\omega}_\alpha \in \bar{S}FxG_\alpha$, z is one-to-one.*

Proof. For $g \in G$, $\beta \in \Omega$, $\beta g z = \bar{\omega}_{\beta g} = \bar{\omega}_\beta g = \beta z g$. It follows that $\bar{\gamma} g z = \bar{\gamma} z g$ for $\bar{\gamma} \in \bar{\Omega}$. If $\bar{\omega}_\alpha \in \bar{S}FxG_\alpha$, z is one-to-one on Ω and hence on $\bar{\Omega}$.

For finite permutation groups, Kuhn [9] established a correspondence between $Z_{S(\Omega)}G$ and FxG_α . Again FxG_α must be expressed as $SFxG_\alpha$.

THEOREM 30. *Let G be a coherent subgroup of $A(\Omega)$ and let $Z = Z_{A(\Omega)}G = Z_{S(\Omega)}G$. If $z \in Z$ and if $\omega_\alpha = \alpha z \in SFxG_\alpha$, then $\beta z = \omega_\beta$ for all $\beta \in \Omega$. Conversely, if $\omega_\alpha \in SFxG_\alpha$ and if $z: \Omega \rightarrow \Omega$ is defined by setting $\beta z = \omega_\beta$ for $\beta \in \Omega$, then $z \in Z$. Z is a po-group and $z \mapsto \alpha z$ gives an o-isomorphism between the po-set Z and the po-set $SFxG_\alpha$.*

COROLLARY 31. *The po-sets which occur as $SFxG_\alpha$ for coherent o-permutation groups (G, Ω) are precisely those po-sets which are carriers of po-groups. The o-sets which occur in this way with Ω totally ordered are those which are carriers of o-groups.*

Proof. Theorem 30 and Corollary 2.

THEOREM 32. *Let G be a coherent subgroup of $A(\Omega)$, Ω totally ordered. Let $\alpha < \omega_\alpha \in SFxG_\alpha$ and let $z \in Z_{A(\Omega)}G$ be defined by $\beta z = \omega_\beta$, $\beta \in \Omega$. For $\gamma \in \Omega$, $B(\gamma, \omega_\gamma) = \text{Conv}\{\gamma z^i \mid i \in I\}$, I the integers, is the smallest o-block of G containing γ and ω_γ , and the collection of $B(\gamma, \omega_\gamma)$'s forms an o-block system of G . Since $(\delta z)g = (\delta g)z$ for*

$g \in G, \delta \in \Omega$, the action of g on $B(\gamma, \omega_\gamma)$ is determined by its action on (γ, ω_γ) , and we shall say that z is a period of G .

Proof. If $g \in G$ is such that $\gamma g = \gamma z^i$ for some i , then for any j , $(\gamma z^j)g = \gamma g z^j = \gamma z^{j+i}$. Apply Lemma 20 to show that $B(\gamma, \omega_\gamma)$ is an o -block of G . The rest is clear.

THEOREM 33. *Let (G, Ω) be o -primitive, Ω totally ordered, and let $Z = Z_{A(\bar{\Omega})}G$. Let $z \in Z$ and let $\bar{\omega}_\alpha = \alpha z \in \bar{F}xG_\alpha = \bar{S}FxG_\alpha$. Then for $\beta \in \Omega, \beta z = \bar{\omega}_\beta$; and for $\bar{\gamma} \in \bar{\Omega}, \bar{\gamma}z = \sup\{\beta z \mid \beta \in \Omega, \beta \leq \bar{\gamma}\}$. Conversely, if $\bar{\omega}_\alpha \in \bar{F}xG_\alpha$ and if z is defined by $\beta z = \bar{\omega}_\beta$ for $\beta \in \Omega$ and $\bar{\gamma}z = \sup\{\beta z \mid \beta \in \Omega, \beta \leq \bar{\gamma}\}$ for $\bar{\gamma} \in \bar{\Omega}$, then $z \in Z$. Z is an o -group and $z \mapsto \alpha z$ gives an o -isomorphism between the o -set Z and the o -set $\bar{F}xG_\alpha$.*

Proof. $\bar{F}xG_\alpha = \bar{S}FxG_\alpha$ because G_α is a maximal proper convex subgroup of G . If $z \in Z$, then Ωz is a dense subset of $\bar{\Omega}$ by Theorem 25, so since z preserves order, $\bar{\gamma}z = \sup\{\beta z \mid \beta \in \Omega, \beta \leq \bar{\gamma}\}$ for $\bar{\gamma} \in \bar{\Omega}$. Conversely, $\beta z = \bar{\omega}_\beta$ maps Ω one-to-one onto a dense subset of $\bar{\Omega}$, so $\bar{\gamma}z = \sup\{\beta z \mid \beta \in \Omega, \beta \leq \bar{\gamma}\}$ extends z to an o -permutation of $\bar{\Omega}$.

COROLLARY 34. *If G is o -2-semitransitive, $Z_{A(\bar{\Omega})}G$ is trivial. If G is o -primitive and regular, $Z_{A(\bar{\Omega})}G$ is isomorphic as an o -group to the integers or the reals.*

Proof. Use the theorem. In the regular case, G is the regular representation of a subgroup of the reals, and every proper Dedekind complete subgroup of the reals is discrete. In the next section we shall deal with the remaining o -primitive groups.

PROPOSITION 35. *For any totally ordered Ω and any subset F of $A(\Omega)$, $Z_{A(\Omega)}F$ is a (not necessarily transitive) l -subgroup of $A(\Omega)$.*

Proof. Since an l -group is a distributive lattice, if z_1 and z_2 commute with $f \in F$, then $(z_1 \vee z_2)f = z_1 f \vee z_2 f = f z_1 \vee f z_2 = f(z_1 \vee z_2)$.

7. Periodically o -primitive groups. We assume from now on that Ω is totally ordered. Earlier we noted that o -2-semitransitive groups and Archimedean regular groups are o -primitive. Now we assume that G is one of the remaining o -primitive groups and prove that it looks strikingly like the group in Example 8.

LEMMA 36. G_α has a first positive long orbital Δ_1 . α is the only point between Δ'_1 and Δ_1 .

Proof. Since G is not regular, G_α has a long orbital Δ . Since G is balanced, Δ may be assumed negative and thus not cofinal with Ω , so that $\bar{\mu} = \sup \Delta \in \bar{\Omega}$. Pick $g \in G$ such that $\alpha \in \Delta g$ and let $\Delta_1 = \text{Conv}((\bar{\mu}g)G_\alpha)$. Pick an arbitrary $\beta \in \Omega$ such that $\alpha < \beta < \bar{\mu}g$. Since $\bar{\mu}G$ is dense in $\bar{\Omega}$ by Theorem 25, we may pick $h \in G$ such that $\alpha < \bar{\mu}h \leq \beta$ and $h \leq g$. $\alpha \in \Delta h$ and thus $\alpha h^{-1} \in \Delta$. Since also $\alpha g^{-1} \in \Delta$, we may pick $k \in G_\alpha$ such that $(\alpha g^{-1})k \geq \alpha h^{-1}$. Now $\alpha(g^{-1}kh) \geq \alpha$, but $(\bar{\mu}g)g^{-1}kh \leq \bar{\mu}kh = \bar{\mu}h$ (since $\bar{\mu}$ is α -full) $\leq \beta$. Finally, we pick $1 \geq m \in G$ such that $(\alpha g^{-1}kh)m = \alpha$. Letting $n = g^{-1}khm$, we have $an = \alpha$ and $(\bar{\mu}g)n \leq \beta$. Since β was arbitrary, there are no points between α and Δ_1 , and Δ_1 is thus the first positive orbital. In view of the definition of Δ_1 , this implies that Δ_1 is long.

Let us define $\bar{\omega} = \bar{\omega}_\alpha \in \bar{F}xG_\alpha$ to be $\sup \Delta_1$. (Δ_1 is bounded above in Ω because G is not o -2-semitransitive.) Let $z \in Z_{A(\bar{\omega})}G$ be the o -permutation of $\bar{\Omega}$ associated with $\bar{\omega}_\alpha$ by Theorem 33. For each integer k , we define $\bar{\omega}_k$ to be αz^k . In particular, $\bar{\omega}_0 = \alpha$ and $\bar{\omega}_1 = \bar{\omega}$. We define Δ_k to be $(\bar{\omega}_{k-1}, \bar{\omega}_k) \subseteq \Omega$, so that $\bar{\Delta}_k = \bar{\Delta}_1 z^{k-1}$. ($\bar{\Delta}_k$ does not include $\bar{\omega}_{k-1}$ or $\bar{\omega}_k$.) The new definition of $\bar{\Delta}_1$ agrees with the old. Since G has period z and since the orbitals of G_α are convex, the fact that Δ_1 is an orbital of G_α implies that each Δ_k is an orbital of G_α . Thus for $k > 0$, Δ_k is the k^{th} positive long orbital; and Δ_{-k} is the $k + 1^{\text{st}}$ long orbital to the left of α . Since G is balanced, Δ_k is paired with Δ_{-k+1} . Between Δ_k and Δ_{k+1} lies precisely one point of $\bar{\Omega}$, namely $\bar{\omega}_k$. If $\bar{\omega}_k \in \Omega$, then $\bar{\omega}_k \in FxG_\alpha (= SFxG_\alpha)$.

LEMMA 37. *For any integers n and k and any $g \in G$, $\alpha g \in \Delta_n$ implies $\bar{\omega}_k g \in \bar{\Delta}_{k+n}$.*

Proof. $\bar{\omega}_k g = \alpha z^k g = \alpha g z^k \in \bar{\Delta}_n z^k = \bar{\Delta}_{k+n}$.

COROLLARY 38. $\text{Conv} \{ \Delta_k \mid k \text{ an integer} \} = \Omega$.


Proof. By Lemma 20, this set is an o -block of the o -primitive group G .

LEMMA 39. *Suppose that some $\bar{\omega}_i \in \Omega (i \neq 0)$. Let n be the least positive integer such that $\bar{\omega}_n \in \Omega$. Then $\bar{\omega}_k \in \Omega$ iff k is a multiple of n .*

Proof. $\bar{\omega}_n$ is the least positive point in the symmetric set $SFxG_\alpha$. Proposition 10 guarantees first that if k is a multiple of n , $\bar{\omega}_k \in \Omega$; and then the converse.

Recapitulating, the (strongly) long orbitals Δ_k of G_α form a set

α -isomorphic to the integers; and denoting $\sup \Delta_k$ by $\bar{\omega}_k$, so that $\bar{\omega}_0 = \alpha$, either the (strongly) fixed points of G_α are precisely those $\bar{\omega}_k$'s such that k is a multiple of some fixed positive integer n , in which case we say that G has *Config* (n), or α is the only fixed point of G_α , in which case we say that G has *Config* (∞).



MAIN THEOREM 40. *Suppose that G is a coherent subgroup of $A(\Omega)$, Ω totally ordered, and that G is o -primitive, but not o -2-semitransitive or regular. Then for some $n = 1, 2, \dots, \infty$, G has *Config* (n). $Z_{A(\bar{\Omega})}G$ is cyclic, having as a generator the o -permutation z of $\bar{\Omega}$ defined by $\beta z = (\bar{\omega}_1)_\beta$ for $\beta \in \Omega$ and $\bar{\gamma}z = \sup \{\beta z \mid \beta \in \Omega, \beta \leq \bar{\gamma}\}$ for $\bar{\gamma} \in \bar{\Omega}$. We shall say that z is the period of G and that G is periodically o -primitive. Δ_{k+1} is "one period up" from Δ_k in the sense that $\bar{\Delta}_k z = \bar{\Delta}_{k+1}$. If G has *Config*(n) for some finite n , $Z_{A(\Omega)}G$ is cyclic, having as a generator the o -permutation \hat{z} of Ω defined by $\beta \hat{z} = (\bar{\omega}_n)_\beta, \beta \in \Omega$; and if G has *Config* (∞), $Z_{A(\Omega)}G$ is trivial.*

A few comments on this theorem are in order. z generates $Z_{A(\bar{\Omega})}G$ by Theorem 33. The fact that $(\bar{\delta}z)g = (\bar{\delta}g)z$ for $g \in G, \bar{\delta} \in \bar{\Omega}$, means that the action of G on Ω is determined by its action on any interval $(\bar{\gamma}, \bar{\gamma}z)$, and in particular on any Δ_k . z is analogous to the function $z: \beta \rightarrow \beta + 1$ of Example 8. If G has *Config*(n) for some finite n and if \hat{z} is the period associated with $\bar{\omega}_n$, then \hat{z} is nicer than z in that it is in $A(\Omega)$ rather than merely in $A(\bar{\Omega})$, but it suffers the disadvantage of being a larger and ultimately less useful period. In the next section, we shall construct examples of o -primitive groups having all of these configurations. Unfortunately, o -imprimitive groups can also have all of these configurations except *Config* (1). What o -blocks might there be containing α ?

PROPOSITION 41. *If an o -imprimitive group G has *Config*(n), n finite, then for some integer $p, 1 \leq p \leq n/2$, the nontrivial o -blocks of G containing α are precisely the sets $\text{Conv} (\Delta'_k \cup \Delta_k), k = 1, \dots, p$. If G has *Config* (∞), this result holds for some $p \geq 1$; or else every $\text{Conv} (\Delta'_k \cup \Delta_k)$ is an o -block.*

Proof. Every nontrivial o -block containing α is symmetric and thus must be of the form $\text{Conv} (\Delta'_k \cup \Delta_k)$ for some $k \geq 1$. If $\text{Conv} (\Delta'_p \cup \Delta_p)$ is an o -block, successive applications of Theorem 21 show that $\text{Conv} (\Delta'_k \cup \Delta_k)$ is an o -block for $k = p - 1, p - 2, \dots, 1$. By Proposition 10, if n is finite, $\text{Conv} (\Delta'_p \cup \Delta_p)$ cannot be an o -block unless

$p \leq n/2$. All of the possibilities not excluded in the proposition do in fact occur for o -imprimitive l -permutation groups (G, Ω) .

COROLLARY 42. *If G has Config (1), G is o -primitive.*

COROLLARY 43. *Suppose G has Config(n) for some $n=1, 2, \dots, \infty$. Then G is o -imprimitive iff $\text{Conv}(\Delta'_1 \cup \Delta_1)$ is an o -block of G .*

This corollary says that whether G is periodically o -primitive is determined by its configuration and knowledge of whether $\text{Conv}(\Delta'_1 \cup \Delta_1)$ is an o -block.

We now investigate the consequences of periodicity. By the support of $g \in \Omega$ we mean $\{\beta \in \Omega \mid \beta g \neq \beta\}$.

COROLLARY 44. (Holland, [7]). *If G is o -primitive, but not o -2-semitransitive, then any $1 \neq g \in G$ has support bounded neither above nor below.*

COROLLARY 45. (Lloyd, [10]). *If $A(\Omega)$ is o -primitive, then it is either o -2-transitive or the regular representation of an Archimedean o -group.*

Proof. Clearly $A(\Omega)$ is not periodic; and the orbits of $A(\Omega)_\alpha$ are automatically convex.

An l -group is l -simple if it has no proper l -ideals.

COROLLARY 46. *An o -primitive l -subgroup G of $A(\Omega)$ is l -simple unless it is o -2-transitive and contains elements of unbounded support.*

Proof. Suppose G is periodically o -primitive. If $1 < g \in G$, then every $\bar{\beta} \in \bar{\Omega}$ is contained in the support of some conjugate of g by Theorem 25. Using periodicity, we apply the argument given at the end of [6] to show that G is l -simple. If G is regular, it is an Archimedean o -group, so it is l -simple. If G is o -2-transitive and contains only elements of bounded support, then G is l -simple by the proof of Theorem 6 of [5]. Note that if Ω is the reals, $A(\Omega)$ is o -2-transitive, but the elements of bounded support form a proper l -ideal.

An o -ideal of a po -group is a normal convex subgroup which is directed. The proof of Corollary 46 also yields

COROLLARY 47. *Suppose that G is an o -primitive subgroup of $A(\Omega)$, Ω totally ordered. Then G lacks proper o -ideals unless it is o -2-semitransitive and contains elements of unbounded support.*

PROPOSITION 48. *Suppose G (not necessarily o -primitive) has $\text{Config}(n)$, n finite. Then any two orbits Δ_j and Δ_k whose subscripts are equal modulo n are o -isomorphic.*

Proof. Proposition 10.

PROPOSITION 49. *Suppose G is periodically o -primitive. Then all long orbitals of G_α have the same cardinality.*

Proof. Let Δ_k be any long orbital of G_α . All proper segments of Δ_k which are cointial with Δ_k have the same cardinality \aleph_I ; and all which are cofinal have the same cardinality \aleph_F . Furthermore, these cardinalities are independent of k . The proposition follows.

COROLLARY 50. *Suppose that G is periodically o -primitive and that some long orbital of G_α is countable. Then all long orbitals of G_α are o -isomorphic to the rationals and so is Ω .*

We can also deduce analogs of several theorems about nonordered permutation groups. For example, if G is a primitive permutation group, $FxG_\alpha = \{\alpha\}$ unless G is regular and $|\Omega|$ is prime [17, Theorem 7.14]. By Theorem 40, this is almost true if G is an o -primitive o -permutation group. Wielandt [17, Theorem 10.13] shows that if a permutation group G is primitive (and if $|\Omega| > \aleph_0$), then for every orbit $\Delta \neq \{\alpha\}$ of G_α , $|\Delta| + |\Delta'| = |\Omega|$. The proof fails for o -primitive groups, but almost all of the conclusion is given by

COROLLARY 51. *Let G be an o -primitive group. Then for every long orbital Δ of G_α , $|\Delta| + |\Delta'| = |\Omega|$. Except when G is o -2-semitransitive, we can strengthen this to $|\Delta| = |\Omega|$.*

Proof. If G is periodically o -primitive, use Proposition 48 and the fact that G has $\text{Config}(n)$. If G is o -2-semitransitive or regular, the conclusion is trivial. It is possible for an o -2-transitive group to have positive and negative orbits of different cardinalities (Example 4).

Wielandt [17, Theorem 10.15] also shows that under somewhat stronger hypotheses, $|\Delta'| = |\Delta|$. This conclusion is given by

COROLLARY 52. *Let G be o -primitive, but not o -2-semitransitive. Then for every orbital Δ of G_α , $|\Delta'| = |\Delta|$.*

8. Full periodically o -primitive groups. For any periodically o -primitive group G , $G \cong Z_{A(\bar{v})}z \cap A(\Omega)$. We shall say that G is full if equality obtains. By Proposition 35, a full periodically o -primitive

group G is automatically an l -subgroup of $A(\Omega)$ and hence the orbits of G_α are convex.

PROPOSITION 53. *Every periodically o-primitive (G, Ω) is contained in a full group (W, Ω) having the same period z .*

Proof. Take $W = Z_{A(\bar{\Omega})}z \cap A(\Omega)$.

In order to construct groups having $\text{Config}(n)$, we characterize those o -sets which occur as Δ_i 's for periodically o -primitive groups G for which the orbits of G_α are convex. Let $I_n = \{1, \dots, n\}$ if n is finite; and let I_n be the integers if $n = \infty$. Let $\Sigma_i = \Delta_i z^{-(i-1)} \subseteq \bar{\Delta}_i, i \in I_n$. The Σ_i 's are pairwise disjoint because $\Omega z^k \cap \Omega = \square$ for $k = 1, \dots, n - 1$ (all k if $n = \infty$). Thus

(a) $\bar{\Delta}_1$ has a collection $\{\Sigma_i | i \in I_n\}$ of dense pairwise disjoint subsets, with $\Sigma_1 = \Delta_1$.

Since for any $h \in G_\alpha, i \in I_n, \Sigma_i h = \Delta_i z^{-(i-1)} h = \Delta_i h z^{-(i-1)} = \Delta_i z^{-(i-1)} = \Sigma_i$, we have

(b) $\{f \in A(\Delta_1) | \Sigma_i f = \Sigma_i \text{ for all } i \in I_n\}$ is transitive on Δ_1 .

For $\bar{\eta} \in \bar{\Delta}_1$, let $L(\bar{\eta}) = \{\bar{\delta} \in \bar{\Delta}_1 | \bar{\delta} < \bar{\eta}\}$ and $R(\bar{\eta}) = \{\bar{\delta} \in \bar{\Delta}_1 | \bar{\delta} > \bar{\eta}\}$. Suppose $\alpha g \in \Delta_k, g \in G, k \in I_n$. Let $\bar{\mu} = \alpha g z^{-(k-1)} \in \Sigma_k$. Let $\bar{\nu} = \bar{\omega}_k g^{-1} (= \bar{\omega}_n z^{k-n} g^{-1} = \bar{\omega}_n g^{-1} z^{k-n} \in \Sigma_{n-(k-1)}$ if n finite). Since $g z^{-(k-1)}$ maps $L(\bar{\nu})$ onto $R(\bar{\mu})$ and $g z^{-k}$ maps $R(\bar{\nu})$ onto $L(\bar{\mu})$, we obtain

(c) For any $\bar{\mu}$ in any $\Sigma_k, k \in I_n$, there exists $\bar{\nu} (\bar{\nu} \in \Sigma_{n-(k-1)}$ if n finite, and $\bar{\nu} \in \bar{\Delta}_1 \setminus \cup \{\Sigma_i\}$ if $n = \infty$) such that there exists an o -isomorphism $s(\bar{\mu}, \bar{\nu})$ of $L(\bar{\nu})$ onto $R(\bar{\mu})$ with $(L(\bar{\nu}) \cap \Sigma_j) s(\bar{\mu}, \bar{\nu}) = R(\bar{\mu}) \cap \Sigma_p$, where $p = j + k - 1 \pmod n$ if n finite), and there exists an o -isomorphism $t(\bar{\mu}, \bar{\nu})$ of $R(\bar{\nu})$ onto $L(\bar{\mu})$ with $(R(\bar{\nu}) \cap \Sigma_j) t(\bar{\mu}, \bar{\nu}) = L(\bar{\mu}) \cap \Sigma_q$, where $q = j + k \pmod n$ if n finite).

Sets Δ_1 satisfying these conditions will be discussed in the corollaries of the following theorem. When $n = 1$, these conditions state simply that $A(\Delta_1)$ is transitive and that for $\delta \in \Delta_1, \{\beta \in \Delta_1 | \beta < \delta\}$ is o -isomorphic to $\{\beta \in \Delta_1 | \beta > \delta\}$; or equivalently, that Δ_1 is an open interval of some chain Ω for which $A(\Omega)$ is o -2-transitive.

THEOREM 54. *The o -sets which occur as first positive orbits in periodically o -primitive groups G which have $\text{Config}(n)$ and for which the orbits of G_α are convex are precisely those o -sets Δ_1 satisfying conditions (a), (b), and (c).*

Proof. We construct, for any o -set Δ_1 satisfying these conditions, a full periodically o -primitive group (G, Ω) having Δ_1 as the first positive orbit of G_α . As the construction for $n = \infty$ is similar to and simpler than the construction for finite n , we shall assume that n is

finite and leave the case $n = \infty$ to the reader.

Let $\Delta_i (= \Sigma_i), \dots, \Delta_n$ be pairwise disjoint copies of $\Sigma_1, \dots, \Sigma_n$, and let A be the ordinal sum $\Delta_1 + \dots + \Delta_n$ with a point α adjoined at the bottom. Let Ω be $\overleftarrow{A} \times I, I$ the integers. For each $i \in I$, let $\Delta_i = \{(\sigma, a) \mid \sigma \in \Delta_0\}$, where $i = an + b$ ($1 \leq b \leq n$). This identifies A with $\{(\lambda, 0) \mid \lambda \in A\}$. Let $\bar{\omega}_i = \sup \bar{\Delta}_i$. $\bar{\omega}_i \in \Omega$ iff i is a multiple of n . Define $\hat{z} \in A(\Omega)$ by $(\lambda, a)\hat{z} = (\lambda, a + 1)$. Now pick an o -isomorphism w_i from Σ_i onto $\Delta_i, i = 1, \dots, n$, with w_1 the identity map on Δ_1 . Since Σ_i is a dense subset of $\bar{\Delta}_1$, we can extend w_i to an o -isomorphism of $\bar{\Delta}_1$ onto $\bar{\Sigma}_i$. We define $z \in A(\bar{\Omega})$ as follows: For $\bar{\beta} \in \bar{\Delta}_i, i = 1, \dots, n - 1, \bar{\beta}z = \bar{\beta}w_i^{-1}w_{i+1}$, and for $\beta \in \bar{\Delta}_n, \bar{\beta}z = \bar{\omega}_n^{-1}\hat{z}$. $\bar{\omega}_i z = \bar{\omega}_{i+1}, i = 0, \dots, n - 1$. This defines z on $\bar{A} = [\alpha, \bar{\omega}_n)$, and we extend it to $\bar{\Omega}$ so that it has \hat{z} as a period, i.e., we define $(\beta\hat{z}^j)z = (\beta z)\hat{z}^j$ for all $\beta \in [a, \bar{\omega}_n), j \in I$.

We define G to be $Z_{A(\bar{\Omega})} \cap A(\bar{\Omega})$, an l -subgroup of $A(\bar{\Omega})$. First we show that G is transitive on Ω . It suffices to show that for each $\alpha \neq \lambda \in A$, there exists $g \in G$ such that $\alpha g = \lambda$. $\lambda \in \Delta_k$ for some $k \in I_n$, so that $\bar{\mu} = \lambda w_k^{-1} \in \Sigma_k$. Pick $\bar{\nu} \in \Sigma_{n-(k-1)}, s(\bar{\mu}, \bar{\nu})$, and $t(\bar{\mu}, \bar{\nu})$ as in (c). Now we define $g \in G$ as follows: $\alpha g = \lambda$ and $(\bar{\nu} w_{n-(k-1)})g = \bar{\omega}_n$. For $\beta \in (L(\bar{\nu}) \cap \Sigma_j)w_j, \beta g = \beta w_j^{-1}s(\bar{\mu}, \bar{\nu})w_{j+(k-1)} \in \Delta_{j+(k-1)}$, where if $j + (k-1) > n, w_{j+(k-1)} = w_{j+(k-1)-n}\hat{z}$. For $\beta \in (R(\bar{\nu}) \cap \Sigma_j)w_j, \beta g = \beta w_j^{-1}t(\bar{\mu}, \bar{\nu})w_{j+k} \in \Delta_{j+k}$. This defines g on $A = [\alpha, \bar{\omega}_n)$, and we extend it to Ω by defining $(\beta\hat{z}^j)g = (\beta g)\hat{z}^j$ for all $\beta \in [a, \bar{\omega}_n), j \in I$. Since $w_i^{-1}w_{i+1} = z$ and $z^n = \hat{z}$, we have $g \in G$, establishing the transitivity of G .

Each $\bar{\omega}_j$ is fixed by G_α because for $h \in G_\alpha, \bar{\omega}_j h = \alpha z^j h = \alpha h z^j = \alpha z^j = \bar{\omega}_j$. By (b), the first positive orbit of G_α is Δ_1 , and since G has period z , the j^{th} positive long orbit of G_α is Δ_j , so that G has $\text{Config}(n)$. By periodicity, no $\text{Conv}(\Delta'_j \cup \Delta_j)$ is an o -block of G , so G is o -primitive, and by construction, it is full.

COROLLARY 55. *For each $n = 1, 2, \dots, \infty$, there is a full periodically o -primitive group on the rationals (which are the only countable candidate) having $\text{Config}(n)$.*

Proof. Let Δ_i be the rationals, which satisfy conditions (a), (b), and (c). (Take the Σ_i 's to be distinct cosets of the rationals in the reals). By Corollary 50, Ω is o -isomorphic to the rationals.

COROLLARY 56. *Suppose that Ω is Dedekind complete and that G is a coherent subgroup of $A(\Omega)$. (Do not assume that G is o -primitive). Then*

- (1) G is the regular representation of the integers or the reals,
 - or (2) G is o -2-semitransitive and $|\Omega| = 2^{\aleph_0}$,
 - or (3) G is periodically o -primitive with $\text{Config}(1)$ and $|\Omega| = 2^{\aleph_0}$.
- $A(\Omega)$ is o -2-transitive for uncountably many nonisomorphic Dedekind

complete Ω 's; and uncountably many nonisomorphic Dedekind complete Ω 's support full periodically o -primitive groups having $\text{Config}(1)$.

Proof. Since Ω is Dedekind complete and nontrivial o -blocks of G have no sups in Ω , G must in fact be o -primitive. If g is regular, it is Archimedean, so since Ω is Dedekind complete, G must be isomorphic as an o -permutation group to the regular representation of the integers or the reals. If G has $\text{Config}(n)$ for some n , then $n = 1$ because Ω is Dedekind complete.

For the statements about cardinality, we appeal to some interesting results of Babcock [1]. Babcock's Theorem 22 states that a Dedekind complete chain, not the integers, which is homogeneous (and thus in its order topology satisfies the first countability axiom by [16, Theorem 1]) has cardinality 2^{\aleph_0} . This finishes (2) and (3). When Ω is Dedekind complete, the $\text{Config}(1)$ conditions on Δ_1 state precisely that Δ_1^* (Δ_1 with end points) is Dedekind complete and that any two nontrivial closed subintervals of Δ_1^* are o -isomorphic. Babcock constructs uncountably many o -sets satisfying these conditions [1, p. 2]. Moreover, it can be verified that in this special case, Δ_1 is o -isomorphic to Ω , so we get uncountably many nonisomorphic Dedekind complete Ω 's supporting full periodically o -primitive groups having $\text{Config}(1)$. Of course, for each of these Ω 's, $A(\Omega)$ is o -2-transitive.

9. Locally o -primitive groups. Following Holland [7], we say when Ω is totally ordered that G is *locally o -primitive* if in the totally ordered set (Theorem 12) of o -block systems of G , there is a minimal nontrivial system $\tilde{\Delta}$. Certainly o -primitive groups are locally o -primitive. The o -blocks in $\tilde{\Delta}$ are called the *primitive segments* of G . If Γ is a primitive segment, let $G|_{\Gamma}$ denote the restriction of G to Γ , i.e., $\{g|_{\Gamma} : g \in G \text{ and } \Gamma g = \Gamma\}$. Then $(G|_{\Gamma}, \Gamma)$ is o -primitive. As noted in the introduction, every l -group can be embedded in a subdirect product of o -permutation groups (G_i, Ω_i) , with each Ω_i totally ordered and G_i a transitive l -subgroup of $A(\Omega_i)$. It can be further arranged that each G_i is locally o -primitive [7].

If for some (and hence each) primitive segment Γ , $G|_{\Gamma}$ is o -2-semitransitive (regular, periodically o -primitive), we shall say that G is *locally o -2-semitransitive (regular, periodically o -primitive)*. For example, the o -imprimitive groups of Proposition 41 are locally o -2-semitransitive; and if Ω is discrete, G is locally regular with primitive segments o -isomorphic to the integers.

THEOREM 57. *Every locally o -primitive group is locally o -2-semitransitive, locally regular, or locally periodically o -primitive.*

We almost characterize locally o -primitive groups by their configurations with

THEOREM 58. *If G_α has a first positive orbital, then G is locally o -primitive. Conversely, if G is locally o -primitive, then G_α has a first positive orbital (unless G is locally regular and Ω is not discrete).*

Proof. Suppose that G_α has a first positive orbital Δ . By Proposition 13, every o -block $\neq \{\alpha\}$ of G which contains α must contain Δ . Let Γ be the intersection of all such o -blocks. Since $\{\alpha\} \neq \Gamma$, Γ must be a primitive segment of G . Therefore G is locally o -primitive. The converse follows from the previous theorem.

10. Examples.

EXAMPLE 1. Let Ω be the reals and let G be the set of o -permutations of Ω having everywhere a strictly positive derivative. G is an o -2-transitive coherent subgroup of $A(\Omega)$, but it is not an l -subgroup.

EXAMPLE 2. Let Ω be the reals and let G be the linear group $\{\alpha x + b \mid \alpha, b \text{ real}, \alpha > 0\}$. $\alpha x + b$ is positive iff $\alpha = 1$ and $b \geq 0$. Again G is coherent and o -2-transitive, but not an l -permutation group.

EXAMPLE 3. In Example 2, let H be the coherent subgroup of elements $\alpha x + b$ of G for which α is rational. H is not o -2-transitive, but is o -2-semitransitive. Although H is o -primitive, it is not primitive because the rationals form a block of H .

EXAMPLE 4. Let ω_1 be the first uncountable ordinal; let Σ be the rationals with the usual order; and let Ω be the lexicographic product $\overleftarrow{\Sigma} \times \omega_1$, ordered from the right, i.e., $(\sigma_1, \gamma_1) \leq (\sigma_2, \gamma_2)$ iff $\gamma_1 < \gamma_2$, or $\gamma_1 = \gamma_2$ and $\sigma_1 \leq \sigma_2$. $A(\Omega)$ is o -2-transitive. The negative orbit of $A(\Omega)_\alpha$ is countable; the positive orbit is not.

EXAMPLE 5. Let I be the integers with the usual order. $A(I)$ is isomorphic as an o -group to the integers. Let (G, Ω) be the ordered wreath product of $(A(I), I)$ with itself, i.e., $\Omega = \overleftarrow{I} \times I$ and each $g \in G$ is given by $(m, n)g = (m + k_n, n + k)$, where k depends only on g , but k_n depends on n as well as g . In fact, $G = A(\Omega)$, and the configuration of G can be obtained by starting with I , replacing one integer by a set of strongly fixed points o -isomorphic to I , replacing each other integer by a strongly long orbit, and establishing the obvious pairings.

EXAMPLE 6. Let $A(\Omega)$ be as in Example 5. Let G be the coherent subgroup of elements of $A(\Omega)$ which satisfy

$$(1) \quad k_n = k_p \quad \text{if } n \equiv p \pmod{2}$$

and

$$(2) \quad k_n \equiv k_p \pmod{2} \quad \text{even if } n \not\equiv p \pmod{2}.$$

None of the long orbits of G_α is convex; indeed, each long orbital of G_α contains precisely two long orbits. The configuration of G consists of alternating strongly long orbitals and o -blocks (each o -isomorphic to the integers) of strongly fixed points.

EXAMPLE 7. In Example 6, replace (2) by (2') $k_n = -k_p$ if $n \equiv p \pmod{2}$. Then G is not coherent; indeed no point can be moved to its successor by a positive $g \in G$. (G, Ω) is regular, but not o -isomorphic to the right regular representation of G . $\Delta \rightarrow \Delta'$ is not an o -anti-automorphism of the totally ordered set of orbit(al)s of G_α . $\{(i, 0) \mid i \text{ even}\}$ is a block Δ of G which is not convex; but $\{g \in G \mid (0, 0)g \in \Delta\}$ is trivially ordered and hence is a convex subgroup of G .

EXAMPLE 8 (Holland, [6]). The only previously known example of an o -primitive group which is neither o -2-semitransitive nor regular was as follows: Let Ω be the reals and let $G = \{f \in A(\Omega) \mid f \text{ has period } 1, \text{ i.e., } (\beta + 1)f = \beta f + 1 \text{ for all } \beta \in \Omega\}$. The map $z: \Omega \rightarrow \Omega$, given by $\beta z = \beta + 1$, lies in the centralizer $Z_{A(\Omega)}G$ of G in $A(\Omega)$, and indeed $G = \{f \in A(\Omega) \mid zf = fz\}$. G is a full periodically o -primitive group having Config(1). (See §7). It is shown in [6] that G is l -simple.

EXAMPLE 9. Let G be the full periodically o -primitive group of Example 8. Let $G^{(m)}$ consist of those elements of G which have m^{th} derivatives and whose first derivatives are positive everywhere. Then $G \supset G^{(1)} \supset G^{(2)} \supset \dots$. Each $G^{(m)}$ is periodically o -primitive with period 1. The $G^{(m)}$'s are not l -subgroups of $A(\Omega)$ and of course are not full.

REFERENCES

1. W. W. Babcock, *On linearly ordered topological spaces*, Dissertation, Tulane University, 1964.
2. L. Fuchs, *Partially ordered algebraic systems*, Addison-Wesley, Reading, Mass., 1963.
3. D. G. Higman, *Finite permutation groups of rank 3*, Math. Zeit., **86** (1964), 145-156.
4. G. Higman, *On infinite simple groups*, Publ. Math. Debrecen, **3** (1954), 221-226.
5. C. Holland, *The lattice-ordered group of automorphisms of an ordered set*, Michigan Math. J., **10** (1963), 399-408.
6. _____, *A class of simple lattice-ordered groups*, Proc. Amer. Math. Soc., **16** (1965), 326-329.
7. _____, *Transitive lattice-ordered permutation groups*, Math. Zeit., **87** (1965), 420-433.
8. C. Holland and S. H. McCleary, *Wreath products of ordered permutation groups*, Pacific J. Math., **31** (1969), 703-716.

9. Harry Waldo Kuhn, *On imprimitive substitution groups*, Amer. J. Math., **26** (1904), 45-102.
10. J. T. Lloyd, *Lattice-ordered groups and σ -permutation groups*, Dissertation, Tulane University, 1964.
11. ———, *Complete Distributivity in Certain Infinite Permutation Groups*, Mich. Math. J., **14** (1967), 393-400.
12. S. H. McCleary, *Pointwise suprema of order-preserving permutations*, Illinois J. Math., **16** (1972), 69-75.
13. ———, *The closed prime subgroups of certain ordered permutation groups*, Pacific J. Math., **31** (1969), 745-753.
14. ———, *Generalized wreath products viewed as sets with valuation*, Journal of Algebra, **16** (1970), 163-182.
15. F. Sik, *Automorphismen geordneter Mengen*, Casopis Pest. Mat., **83** (1958), 1-22.
16. L. B. Treybig, *Concerning homogeneity in totally ordered, connected topological spaces*, Pacific J. Math., **13** (1963), 1417-1421.
17. H. Wielandt, *Unendliche Permutationsgruppen*, Lecture Notes, University of Tübingen, 1960.
18. ———, *Finite permutation groups*, Academic Press, New York, N. Y., 1964.

Received February 10, 1970 and in revised form March 11, 1971. This paper grew out of the author's doctoral dissertation, written under the very helpful supervision of Charles Holland and supported in part by the National Science Foundation.

UNIVERSITY OF GEORGIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 40, No. 2

October, 1972

Louis I. Alpert and L. V. Toralballa, <i>An elementary definition of surface area in E^{n+1} for smooth surfaces</i>	261
Eamon Boyd Barrett, <i>A three point condition for surfaces of constant mean curvature</i>	269
Jan-Erik Björk, <i>On the spectral radius formula in Banach algebras</i>	279
Peter Botta, <i>Matrix inequalities and kernels of linear transformations</i>	285
Bennett Eisenberg, <i>Baxter's theorem and Varberg's conjecture</i>	291
Heinrich W. Guggenheimer, <i>Approximation of curves</i>	301
A. Hedayat, <i>An algebraic property of the totally symmetric loops associated with Kirkman-Steiner triple systems</i>	305
Richard Howard Herman and Michael Charles Reed, <i>Covariant representations of infinite tensor product algebras</i>	311
Domingo Antonio Herrero, <i>Analytic continuation of inner function-operators</i>	327
Franklin Lowenthal, <i>Uniform finite generation of the affine group</i>	341
Stephen H. McCleary, <i>0-primitive ordered permutation groups</i>	349
Malcolm Jay Sherman, <i>Disjoint maximal invariant subspaces</i>	373
Mitsuru Nakai, <i>Radon-Nikodým densities and Jacobians</i>	375
Mitsuru Nakai, <i>Royden algebras and quasi-isometries of Riemannian manifolds</i>	397
Russell Daniel Rupp, Jr., <i>A new type of variational theory sufficiency theorem</i>	415
Helga Schirmer, <i>Fixed point and coincidence sets of biconnected multifunctions on trees</i>	445
Murray Silver, <i>On extremal figures admissible relative to rectangular lattices</i>	451
James DeWitt Stein, <i>The open mapping theorem for spaces with unique segments</i>	459
Arne Stray, <i>Approximation and interpolation</i>	463
Donald Curtis Taylor, <i>A general Phillips theorem for C^*-algebras and some applications</i>	477
Florian Vasilescu, <i>On the operator $M(Y) = TYS^{-1}$ in locally convex algebras</i>	489
Philip William Walker, <i>Asymptotics for a class of weighted eigenvalue problems</i>	501
Kenneth S. Williams, <i>Exponential sums over $GF(2^n)$</i>	511