

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Developing Continuous Variable Composites for Rorschach Measures of Thought Problems, Vigilance, and Suicide Risk**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1508195> since 2016-06-30T17:56:06Z

*Published version:*

DOI:10.1177/1073191112446963

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Viglione D; Giromini L; Gustafson ML; Meyer G. Developing Continuous Variable Composites for Rorschach Measures of Thought Problems, Vigilance, and Suicide Risk. *ASSESSMENT*. 21 (1) pp: 42-49.  
DOI: 10.1177/1073191112446963

The publisher's version is available at:

<http://asm.sagepub.com/cgi/doi/10.1177/1073191112446963>

When citing, please refer to the published version.


Link to this full text:

<http://hdl.handle.net/2318/1508195>



**Developing Continuous Variable Composites  
for Rorschach Measures of Thought Problems,**

Journal:	<i>Assessment</i>
Manuscript ID:	ASMNT-12-0027
Manuscript Type:	Original Manuscript
Keywords:	Rorschach, Schizophrenia, Psychosis, Thought Disorder , Vigilance , Suicide, Composites



## Abstract

Using a multiple regression approach with a large developmental sample (N = 460) of Rorschach protocols from psychiatric, forensic, and nonclinical control groups, we created continuous multi-variable Composite scores corresponding to the Comprehensive System (CS, Exner, 2003) Perceptual-Thinking Index (PTI), Hypervigilance Index (HVI), and Suicide Constellation (S-CON). Within a validation sample (N=230), these three new scores, called the Thought and Perception Composite (TP-Comp), Vigilance Composite (V-Comp), and Suicide risk-Composite (SC-Comp) were strongly associated with the three original CS Indices. Additional analyses suggest that the new Composite scores were more reliable than and at least as valid as the original Indices. Interpretive guidelines are offered.

## Developing Continuous Variable Composites

## for Rorschach Measures of Thought Problems, Vigilance, and Suicide Risk

To date, Rorschach systems have not taken full advantage of computers to perform complex calculations. In the Rorschach Comprehensive System (CS; Exner, 2003), for example, clinical Indices are derived from a series of dichotomous cut-off items that then lead to a step-based integer scale rather than a fully dimensional scale. Although these Indices can be calculated by hand, such integer-based scales, relative to dimensional, continuous scales, suffer from distributional and reliability problems because they assume only a limited number of values and fail to use all the available variance. For example, the Perceptual Thinking Index (PTI) is calculated with five individual algebraic statements or “steps” based on cut-off scores, each generating a value of 0 or 1, which are summed to create the total PTI score. One of the underlying steps is  $M- > 1$  or  $X-\% > .40$ ; if either is true, one point is added to the PTI value. Thus, the index can only assume integer values from zero to five. Decimal or intermediate values, such as 3.5 are not possible. One might describe this integer-based, discontinuous structure as having “cliffs.” These cliffs limit the reliability of the index in that small changes in a few responses could have a large impact on the index. As an extreme example, mis-coding a single human movement form quality minus response (M-) might add as much as three points to the PTI total by changing the outcomes for three steps that relate to form quality (i.e., using  $XA\%$  and  $WDA\%$  in Step 1,  $X-\%$  in Step 2, and  $X-\%$  or  $M-$  in Step 3). Theoretically, one might expect limitations in the validity of the scale because cliffs cause it to be less highly correlated with the latent construct that it measures, thought and perceptual disturbance, which would be a demonstration of how reliability can limit validity. Support for this hypothesis also comes from a

1  
2  
3 recent review indicating that continuous measures of psychopathology are more valid and  
4  
5 reliable than discrete measures (Markon, Chmielewski & Miller, 2011). Thus, a fully  
6  
7 dimensional PTI could improve reliability and validity and thus also improve interpretive  
8  
9 accuracy in applied work.  
10  
11

12  
13 The current study attempts to develop continuous variables corresponding to three of the  
14  
15 CS clinical Indices and to demonstrate basic reliability and validity for using these variables.  
16  
17 Based on available research findings (Meyer & Archer, 2001; Meyer, Hsiao, Viglione, Mihura,  
18  
19 & Abraham, 2012; Mihura, Meyer, Dumitrascu, & Bombel 2012; Viglione, 1999; Viglione &  
20  
21 Hilsenroth, 2001; Viglione & Meyer, 2008), we chose two Indices with the most empirical  
22  
23 support – the PTI and Suicide Constellation (S-CON) – and one Index that has some support and  
24  
25 a reasonable inference base related to coded behaviors that transparently generalize to extratest  
26  
27 behaviors – the Hypervigilance Index (HVI). By convention within the CS, the S-CON is  
28  
29 interpreted dichotomously as an indicator of suicide risk. The HVI is likewise dichotomously  
30  
31 interpreted as a focused, effortful, detail oriented, and vigilant cognitive style. However, research  
32  
33 (Lindgren & Carlsson, 2002; Meyer, 1997, 1999) has supported using the S-CON integer score  
34  
35 as a dimensional indicator of distress and the HVI integer score for wariness and paranoid states.  
36  
37 Thus, there is support for using them dimensionally as we do in this study rather than only as a  
38  
39 dichotomous indicator. Indeed, the latent constructs of thought and perceptual disturbance,  
40  
41 vigilance, and intensity of suicidal risk are dimensional in nature and thus are appropriately  
42  
43 measured on a continuous scale. To develop continuous variables corresponding to the PTI, HVI,  
44  
45 and S-CON, we substitute their dichotomous cut-scores and integer results with a regression-  
46  
47 based model that produces continuous scores. To distinguish from the CS Indices, we call these  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 new scores “Composites;” i.e., the Thought and Perception Composite (TP-Comp), Vigilance  
4 Composite (V-Comp), and Suicide Concern Composite (SC-Comp).  
5  
6

#### 7 8 Method 9

10 To produce the new Composites, the individual variables included in the steps of the  
11 original CS Indices were used as predictors in multiple regression with the applicable CS clinical  
12 Index as the criterion. For example, we used all of the variables found in the original five PTI  
13 steps, X-%, M-, etc. as individual predictors with the PTI total score as the criterion variable.  
14  
15 The raw score (b weight) prediction equations were then used to calculate the Composites, which  
16 were then tested for generalizability with an independent validation sample. The goal was to  
17 maximize the association between the original CS Index and its corresponding dimensional  
18 Composite score in the validation sample.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

#### 29 *Variable Selection* 30

31 Several modifications to the predictor variables from the original Indices were necessary.  
32 Some of the original CS steps use a deviation score rather than a simple frequency. For instance,  
33 S-CON Step 3 is positive if the Egocentricity Index is less than .31 or greater than .44. To create  
34 a dimensional variant of this variable in the model, we used the absolute value of the deviation  
35 from the mean of these two cut-points (i.e.,  $|X - .375|$ ). Similarly, for the S-CON criterion of  $P <$   
36  $3$  or  $P > 8$ , we used the absolute value of the deviation from the mean of these two cut-points  
37 (i.e.,  $|X - 5.5|$ ). The Zd score has such limited empirical support (Meyer, Viglione, Mihura,  
38 Erdberg, & Erard., 2011; Mihura et al., 2012) that its use is not advised. Given, that it is likely  
39 not to be used consistently in the future, we replaced with a logical proxy based on the relative  
40 degree of complexity evident in selecting and synthetically linking inkblot locations. More  
41 specifically, we used the Location-Developmental Quality Complexity/R Index (Loc-DQ/R).  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Location and Developmental Quality are the central variables in producing Zd, so that a low Zd  
4  
5 score would correspond to a low Loc-DQ/R score. The fifth S-CON criterion contains a  
6  
7 deviation algorithm,  $Zd < -3.5$  or  $Zd > 3.5$ ; consequently, the absolute value of the difference  
8  
9 between Loc-DQ/R and its mean value in our developmental sample was used ( $ILQ-compl/R -$   
10  
11 1.373).

12  
13  
14  
15 In addition, some predictor variables were transformed to reduce their skew and kurtosis.  
16  
17 For example, distributions of FAB2, SumT, and MOR substantially diverged from a normal  
18  
19 distribution so transformations were applied to prevent the final formulae of the Composites  
20  
21 from being affected by nonnormality, excessive variability, and extreme outliers in any of the  
22  
23 individual variables. Said differently, we wanted to avoid allowing a single extreme value in any  
24  
25 of the individual variables to have an excessively large impact on the final score. In fact,  
26  
27 ultimately, we found that the congruence between the integer Indices and continuous Composites  
28  
29 at the interpretively important higher values was greatest when we took a very conservative  
30  
31 approach of transforming variables with skew values greater than one. In all but two cases the  
32  
33 square root transformation reduced the predictor variable skew sufficiently, but two variables  
34  
35 required an inverse reciprocal to the second power transformation [ $-1/(x+1)^2$ ; see Behrens, 1997].  
36  
37 Eventually, all variables included in the analysis had absolute skew values  $\leq 1$  and absolute  
38  
39 kurtosis values  $\leq 1.5$ , except for the transformed FAB2 variable (skew = 2.17, kurtosis = 2.83).  
40  
41  
42  
43  
44

#### 45 *Sample*

46  
47  
48 We compiled a large sample of computerized Rorschach protocols from our available  
49  
50 records. To fit with contemporary recommendations and data concerning administration to  
51  
52 optimize the range of responses obtained (Dean et al., 2007; Meyer et al., 2011), we used the  
53  
54 same procedures as those used to generate the R-PAS norms to statistically model the number  
55  
56  
57  
58  
59  
60

1  
2  
3 and distribution of responses in these protocols so they would closely approximate the  
4  
5 distribution of responses observed on each of the ten cards when using the R-Optimized (R-Opt)  
6  
7 administration method<sup>1</sup>. This method encourages examinees to give 2 or 3 responses per card,  
8  
9 prompts for another response if only one is given to any card, and stops collecting responses  
10  
11 after the fourth to any card (see Meyer et al., 2011). Applying these R-Opt modeling procedures  
12  
13 to our pool of protocols left 1350 records. This sample included a substantial proportion of  
14  
15 records from non-clinical controls and non-patients or children. However, we wished to  
16  
17 maximize predictions for adults and clinical patients, given that the three Indices address clinical  
18  
19 issues and the S-CON is only calculated for adults. Consequently, all child records and most of  
20  
21 the control and nonpatient records were removed. The final sample included 690 records, 100 of  
22  
23 which were randomly selected from our pool of nonclinical controls or nonpatients. As expected,  
24  
25 the distribution for R in this sample ( $M = 24.1$ ,  $SD = 4.6$ ) closely approximated the normative  
26  
27 distribution for R ( $M = 24.2$ ,  $SD = 4.7$ ; Meyer et al., 2011).  
28  
29  
30  
31  
32  
33

34       Formulae for the Composites were developed within a randomly selected developmental  
35  
36 sample ( $N = 460$ ) and tested with a validation sample ( $N = 230$ ). These 690 adult Rorschach  
37  
38 records came from eight nonoverlapping subgroups, identified by patient status, evaluation  
39  
40 context, diagnosis, or presenting problem. 30.0% were from a mixed sample of inpatients and  
41  
42 outpatients evaluated for psychiatric reasons, 28.7% were from offenders, 14.5% from control or  
43  
44 non-patients, 9.1% from trauma patients, 4.1% from patients with schizophrenia or other  
45  
46  
47  
48  
49

---

50  
51 <sup>1</sup> The R-Opt modeling procedure is described in detail in Meyer et al. (2011). Briefly, it allows  
52  
53 one to select suitable records collected using standard CS guidelines and randomly prune among  
54  
55 excess responses so as to exactly match the response level parameters observed in a sample of  
56  
57 123 target records collected by experienced examiners using R-Opt guidelines. Because R-Opt  
58  
59 greatly limits the number of cards with only one response, records with an insufficient number of  
60  
61 cards having more than one responses are dropped, such that the final pool of modeled records is  
62  
63 smaller than the original pool of available records.

1  
2  
3 psychotic disorders, 3.5% from patients with depression or substance abuse, 1.9% from forensic  
4  
5 psychiatric patients, and 8.3% from patients with other conditions. The same proportions of these  
6  
7 subgroups was retained in the developmental and validation samples.  
8  
9

10 Men comprise 56.6% of the sample and the mean age is 35.04 (SD = 11.54)<sup>2</sup>. More than  
11  
12 50 examiners contributed protocols to the sample. Ethnicity is not well documented in some of  
13  
14 these archival, computerized files. However, in the largest subsample of mixed inpatients and  
15  
16 outpatients, almost one third of these individuals are African-American.  
17  
18

### 19 Results

20 We used the same strategy and method for calculating the three Composites. At first,  
21  
22 multiple regression equations were established within the developmental sample using the  
23  
24 original CS target Index as the dependent variable and fully dimensional versions of its  
25  
26 component scores as predictors. Non-significant predictors were removed and a new model was  
27  
28 tested. The prediction equation derived from the raw score b values produces a fully dimensional  
29  
30 total score to replace the original CS integer score. Each prediction equation was then used to  
31  
32 calculate this new dimensional scale in the derivation sample and subsequently in the  
33  
34 independent validation sample.  
35  
36  
37  
38  
39

40 Descriptive data in Table 1 reveal that the distributions of the Composites closely  
41  
42 resemble the original Indices. Standard deviations are smaller for the all three Composites as a  
43  
44 result of regression to the mean. However, this reduction in variability also may be related to less  
45  
46 error in the Composites from replacing the individual integer steps with dimensional scores.  
47  
48

49 Because of the smaller SDs interpretive ranges to designate high, clinically relevant values will  
50  
51 be slightly lower for the Composite variables in contrast to the original CS Indices.  
52  
53

---

54  
55 <sup>2</sup> Six records were missing gender information and 34 were missing ages, but are known to be  
56  
57 adults.  
58  
59  
60

*TP-Comp*

Table 2 provides information on the original Indices and their regression-based dimensional alternatives. When predicting the PTI, the transformed variable for Level 2 Special Scores [Lvl2;  $(-1/(Lvl2+1)^2)$ ] was not a significant contributor and it was removed from the analysis. All other variables were significant. Within the developmental sample, the overall model was significant ( $F(6,453) = 250.808, p < .001$ ), with a multiple  $R = .88$ . In the independent validation sample, the correlation between the TP-Comp and the observed PTI was .87. Distribution data in Table 1 reveal that the original PTI and new TP-Comp are highly similar. A supplemental table available from the first author provides the M, SD, and confidence interval for the Composite scores at each value of the original CS Index for both the developmental and the validation samples. These results illustrate the regression to the mean noted before. For instance, PTI values of 2, 3, 4, and 5 equate to lower TP-Comp values, with corresponding means of about 1.8, 2.5, 3.5, and 4.5.

*V-Comp*

All V-Comp predictor variables were significant and the model was significant ( $F(8,451) = 171.372, p < .001$ ) with a multiple  $R = .87$  in the developmental sample. The correlation between the V-Comp and the observed HVI within the validation sample was .86. As with TP-Comp, a close correspondence between the Index and Composite at all points of the original scale was achieved. Again, at the more extreme high HVI values, the corresponding V-Comp mean values are less, again demonstrating regression to the mean. HVI values of 5, 6, 7, and 8 have corresponding V-Comp means of approximately 4.5, 5.5, 6.0, and 7.0.

*SC-Comp*

1  
2  
3 R was not significant and was excluded from the regression analysis. Within the  
4  
5 developmental sample, the model was significant ( $F(11,448) = 94.054, p < .001$ ) with a multiple  
6  
7 R of .84. The correlation with the observed S-CON in the validation sample was .79. Descriptive  
8  
9 data for the validation sample in Table 1 reveal nearly identical means. Because of regression to  
10  
11 the mean, S-CON values of 6, 7, 8, and 9 have corresponding SC-Comp means of roughly 5.7,  
12  
13 6.5, 7.0, and 8.0.  
14  
15

### 16 17 *Interrater Reliability*

18  
19 Initial inter-rater reliability data for the Composites was recently investigated in a sample  
20  
21 of 50 Rorschach records collected with R-Opt administration (Viglione, Blume-Marcovici,  
22  
23 Miller, Giromini & Meyer, 2012). This sample was diverse in terms of age, sex, ethnicity,  
24  
25 educational background and patient status, and 16 examiners were involved. The exact  
26  
27 agreement intraclass correlations for a single rater (ICCs) for the Composites and Indices are  
28  
29 shown in Table 3 (upper part). To determine whether the interrater reliability for the new  
30  
31 Composites exceeded that of the corresponding original Indices we used procedures to compare  
32  
33 coefficients derived from the same sample that do not share overlapping variables (Raghunathan,  
34  
35 Rosenthal, & Rubin, 1996). The statistical test compares coefficients using the Pearson-Filon  
36  
37 statistic after they are transformed to Fisher's Zr (ZPF). To obtain an effect size associated with  
38  
39 these comparisons, the difference in the Zr coefficients was converted back from the Zr metric to  
40  
41 the r metric. The resulting ZPF and r difference (r Diff) values shown in Table 3 indicate that the  
42  
43 reliability of the TP-Comp and V-Comp is superior than the reliability of the original Indices  
44  
45 (Table 3, upper part).  
46  
47  
48  
49  
50  
51

52  
53 To test further the expected reliability gains with the Composites over the Indices, we  
54  
55 also re-analyzed two previously published data sets. In 2003, Viglione and Taylor reported  
56  
57  
58  
59  
60

1  
2  
3 strong inter-rater reliabilities for CS variables in a mixed clinical and non-patient sample of 84  
4  
5 individuals. According to our new analyses, the interrater reliability for each of the Composite  
6  
7 scores falls in the excellent range and is significantly higher than the Index reliability (Table 3,  
8  
9 middle part). In 2002, Meyer et al. described interrater reliability for CS variables in several  
10  
11 samples collected by students (N = 66) and clinicians (N = 153). Given that the entire Meyer et  
12  
13 al.' sample size was large enough to select a subset of records with the R-Opt modeling  
14  
15 procedure, we computed the analyses both within the original and within the R-Opt modeled  
16  
17 samples (Table 3, lower part). Compared to the CS Indices, the Composites demonstrated higher  
18  
19 ICCs in all 18 comparisons, with significant differences ( $p < .05$ ) in 13. Taken together, these  
20  
21 results indicate interrater reliability will be higher using the dimensional Composites rather than  
22  
23 the original CS Indices.  
24  
25  
26  
27  
28

### 29 *Validation*

30  
31 To investigate validity, we calculated the Indices and new Composites in previously  
32  
33 published studies. Dean et al. (2007) investigated the PTI among 61 residential care respondents,  
34  
35 31 of whom had psychotic disorders. These individuals were randomly assigned to either a CS  
36  
37 Rorschach administration or an alternative administration designed to control the number of  
38  
39 responses, similar to the R-Opt procedure. In the original study the PTI correlated at .381 ( $p <$   
40  
41  $.01$ ) with a synthetic measure of thought disorder based on semi-structured interview, speech  
42  
43 sample, and self-report. The correlation between the thought disorder measure and the new TP-  
44  
45 Comp was .434 ( $p < .01$ ). The difference between the two correlations was tested using  
46  
47 procedures for comparing correlations obtained from the same sample that share a common  
48  
49 variable (Meng, Rosenthal, & Rubin, 1992). This comparison was not statistically significant,  $Z$   
50  
51  $= 0.866$ ,  $p = 0.386$ , but in the expected direction. In addition, when only the R-Opt like  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 alternative administration records ( $n = 31$ ) were selected and the validity correlations re-  
4  
5 calculated, the TP-Comp seemed to be more valid than the PTI. Indeed, the PTI correlated .435  
6  
7 ( $p < .05$ ), while the TP-Comp correlated .615 ( $p < .001$ ); the difference between the two  
8  
9 correlations is statistically significant,  $Z = 2.360$ ,  $p = .018$ , and of medium size,  $r_{difference} =$   
10  
11 .246. In contrast, when only the standard CS administered records ( $n = 30$ ) were selected, neither  
12  
13 the PTI ( $r = .33$ ,  $p > .05$ ) nor the TP-Comp ( $r = .28$ ,  $p > .05$ ) significantly correlated with the  
14  
15 measure of thought disorder. These results suggest that response distributions consistent with R-  
16  
17 Opt administration increases validity over standard CS administration, and within that context  
18  
19 TP-Comp has better validity than the PTI.  
20  
21  
22  
23

24  
25 Another set of analyses for the PTI and TP-Comp was derived from a sample of 432  
26  
27 consecutive evaluations in a hospital-based assessment service in Chicago (Meyer, 2002). Data  
28  
29 has previously been reported on two proxy measures of psychotic functioning using diagnostic  
30  
31 information obtained from billing records before the psychological testing was initiated: (a)  
32  
33 diagnosis of a psychotic disorder and (b) severity of disturbance based on 5-point continuum  
34  
35 derived from the most severe diagnosis assigned (see Dawes, 1999; Meyer, 2002; Meyer &  
36  
37 Resnick, 1996; Viglione & Hilsenroth, 2001). Table 4 contains the correlations with these two  
38  
39 proxy criterion measures and shows both the correlations within the whole sample and within the  
40  
41 subset of R-Opt modeled records. In one of the four validity coefficient comparisons, the TP-  
42  
43 Comp was superior to the PTI, though with a small effect size.  
44  
45  
46  
47

48  
49 Meyer (1997, 1999; Meyer, Riethmiller, Brooks, Benoit & Handler, 2000) reported on 87  
50  
51 clients selected from this same sample on the basis of concordant response style on the  
52  
53 Rorschach and MMPI-2. When test-taking response styles on the two tests was similar, as  
54  
55 measured by markers of each test's first factor, the CS Indices were substantially correlated with  
56  
57  
58  
59  
60

1  
2  
3 corresponding MMPI-2 Indices. When response styles differed between the two tests,  
4  
5 correlations were either not significant or negative. Using patients with concordant test-taking  
6  
7 styles, Table 5 provides the correlation of the CS Indices with their corresponding MMPI-2  
8  
9 criteria reported in Meyer (1999), as well as, correlations for the corresponding Composites from  
10  
11 the current study, along with significance tests contrasting the two correlations<sup>3</sup>. Overall, these  
12  
13 correlations suggest equal validity for the clinical Indices and Composites. As explained in the  
14  
15 original articles, readers should keep in mind that these are artificially inflated coefficients using  
16  
17 just a subset of the original data after aligning method variance.  
18  
19  
20  
21

## 22 Discussion

23  
24 Our aim was to develop continuous versions of the PTI, HVI, and S-CON in order to gain  
25  
26 reliability and precision over the cut-off-based integers used by the CS Indices. We used  
27  
28 dimensional versions of all the individual variables contained in the Indices and multiple  
29  
30 regression to create the continuous Composite variables. Using a relatively large developmental  
31  
32 sample (N = 460), we created the TP-Comp, V-Comp, and SC-Comp from the raw regression  
33  
34 coefficients applied to the individual predictor variables. These three continuous Composites  
35  
36 were tested in a validation sample (N=230), where the correlation between the original and new  
37  
38 score and the overall distribution parameters revealed a high degree of similarity. According to  
39  
40 these results, TP-Comp, V-Comp and SC-Comp are sufficiently identical to the three original CS  
41  
42 Indices that they can be substituted for them. Overall, our results suggest that the three  
43  
44 Composites have superior interrater reliability and are at least as valid as the original Indices. At  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

---

55 <sup>3</sup> In the original study, the Schizophrenia Index, a precursor of the PTI was reported, but not the  
56 PTI itself. Accordingly, the PTI correlations, in addition to the Composite correlations were  
57 calculated anew for this paper  
58  
59  
60

1  
2  
3 the same time, however, it should be pointed out that these new Composites are statistically  
4 complex and cannot be calculated by hand<sup>4</sup>, whereas their CS predecessors were not so limited.  
5  
6

7  
8 As would be expected with regression toward the mean, extreme low values in the  
9 original CS Indices correspond to slightly higher new Composite values, and extreme high  
10 values in the original CS Indices correspond to slightly lower new Composite values. In terms of  
11 clinical interpretations, however, this should not pose a problem as the results presented here  
12 show that one should merely adjust the interpretive ranges from the original Indices to fit the  
13 new, slightly narrower Composite distributions. Drawing on data from both the development and  
14 validation samples in order to stabilize results coming from the tail ends of the distributions  
15 where observations are not numerous, we can offer interpretive guidelines for the new  
16 Composites by linking them to their equivalent scores on the original Index. In all instances, we  
17 would expect Composite scores to be lower than Index scores. For instance, PTI values of 3, 4,  
18 and 5 roughly correspond to TP-Comp values of 2.5, 3.5, and 4. Thus, a TP-Comp score of 3.5  
19 suggests considerable concern about disturbed thinking and distorted perceptions of reality.  
20 Previously, a PTI score of 3 has been interpreted as evidence of unusual thinking and perception.  
21 TP-Comp scores approaching values of 2.5 should be interpreted in a similar fashion.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 As noted earlier, the HVI in the CS is traditionally interpreted as dichotomous variable,  
42 though Meyer (1999) and Lindgren and Carlsson (2002) found support for its use as a  
43 dimensional variable. Moreover, it measures a focused, effortful, detail oriented, and vigilant  
44 cognitive style. It thus requires some cognitive sophistication and psychological differentiation.  
45 HVI values of 5, 6, 7, and 8 have corresponding V-Comp means of about 4.5, 5.5, 6.0, and 7.0.  
46 From Table 1 it can be seen that a V-Comp score of 4.5 is almost one standard deviation above  
47  
48  
49  
50  
51  
52  
53  
54

55  
56  
57  
58 <sup>4</sup> A file to calculate the Composites is freely available online (see Meyer et al., 2011).  
59  
60

1  
2  
3 the mean. Taking these data together, a V-Comp score exceeding 4.5 suggests a modest degree  
4  
5 of a focused, effortful, detail-oriented, and vigilant cognitive processing. A score of 5.5 suggests  
6  
7 a more extreme, inflexible, and vigilant cognitive style.  
8  
9

10 For the SC-Comp, a similar approach to anchoring interpretation is recommended.  
11  
12 From the literature an S-CON score of 7 suggests possible Suicide Concern that might or might  
13  
14 not be evident in self-report and a score of 8 indicates probable Suicide Concern (Fowler et al.,  
15  
16 2001; Viglione & Hilsenroth, 2001). S-CON values of 6, 7, 8, and 9 have corresponding SC-  
17  
18 Comp means of roughly 5.7, 6.5, 7.0, and 8.0. Thus, an SC-Comp score of 6.5 or above serves as  
19  
20 a suitable replacement for the S-CON score of 7, and would, therefore, suggest possible suicide  
21  
22 risk. We also recommend that a SC-Comp score of 7 be interpreted as suggesting a possible  
23  
24 Suicide Concern that is worthy of being evaluated, though with recognition of the risk for false  
25  
26 positive scores. A SC-Comp score of 8 should be interpreted as indicating a more worrisome risk  
27  
28 suicide.  
29  
30  
31  
32  
33

34 The high similarity between the Indices and Composites indicates they are nearly  
35  
36 identical and should have the same network of relationships with important criteria.  
37  
38 Nevertheless, the validity findings reported here suggest the new TP-Comp is likely more valid  
39  
40 than the PTI, suggesting the continuous, dimensional distribution of the TP-Comp offers an  
41  
42 advantage in capturing more valid variance. The validity data for the V-Comp and SC-Comp  
43  
44 relied on more limited criteria and produced less conclusive results, suggesting that the validity  
45  
46 of these Composites approximates that of the original Indices. All three Composites clearly have  
47  
48 superior interrater reliability. Because of this better reliability, which comes from being  
49  
50 continuous rather than integer based, these Composites will likely be less affected by scoring  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 errors when used in the field, thus improving assessment accuracy in clinical and forensic  
4  
5 evaluations.  
6

7  
8 A limitation of this research is that the great majority of the records used across the  
9  
10 analyses were administered by CS methods and statistically modeled to approximate R-Opt  
11  
12 administration. However, our reliability sample obtained using R-Opt procedures performed just  
13  
14 like the modeled samples, suggesting that the findings should generalize across modes of  
15  
16 administration. This point is reinforced by seeing how similar the validity results are in Table 4  
17  
18 for the original Chicago protocols and for the protocols that remained after R-Opt modeling.  
19  
20 Finally, the strong validity support for the early R-Opt administration method used in the Dean et  
21  
22 al. (2007) data suggests that the TP-Comp findings should generalize to the R-Opt administration  
23  
24 procedure in R-PAS. Another limitation of this study concerns the type of psychopathology and  
25  
26 the patient status represented in these records, such that some populations may be over-  
27  
28 represented (e.g., forensic) and others under-represented (e.g., trauma). Accordingly, future  
29  
30 studies should test the validity of these scales with the R-Opt administration method, specific  
31  
32 clinical populations, and appropriate criteria. Also, in future research it would be worthwhile to  
33  
34 explore how simple z-score sums of variables might perform instead of using regression  
35  
36 weighted variables when creating new versions of these Indices.  
37  
38  
39  
40  
41  
42

43  
44 Another avenue is opened by placing the V-Comp and SC-Comp on continuous scales so  
45  
46 that research can address the validity of these two Composites with dimensional criteria. For the  
47  
48 V-Comp, one can identify potential correlates associated with vigilance, paranoid processing,  
49  
50 and other cognitive and interpersonal phenomena. For the SC-Comp one could address  
51  
52 psychological issues such as suicide intentionality or despair. Also, determining whether there  
53  
54 are correlates to low scores on all three scales would be useful.  
55  
56  
57  
58  
59  
60

## References

- 1  
2  
3  
4  
5  
6 Behrens, J. T. (1997). Principles and procedures of exploratory data analysis. *Psychological*  
7  
8 *Methods*, 2, 131-160.  
9  
10 Dawes, R. M. (1999). Two methods for studying incremental validity of a Rorschach variable.  
11  
12 *Psychological Assessment*, 11, 297-302.  
13  
14  
15 Dean, K., Viglione D.J., Perry, W., & Meyer G.J. (2007). A method to optimize response range  
16  
17 while maintaining Rorschach Comprehensive System validity. *Journal of Personality*  
18  
19 *Assessment*, 89(2), 149-161.  
20  
21  
22 Exner, J. E. (2003). *The Rorschach: A comprehensive system*. (4<sup>th</sup> ed., Vol. 1: Basic foundations  
23  
24 and principles of interpretation). New York: Wiley.  
25  
26  
27 Fowler, J. C., Piers, C., Hilsenroth, M. J., Holdwick, D. J., & Padawer, J. R. (2001). The  
28  
29 Rorschach Suicide Constellation: Assessing various degrees of lethality. *Journal of*  
30  
31 *Personality Assessment*, 76, 333-351.  
32  
33  
34 Lindgren, T., & Carlsson, A. M. (2002). Relationship Between the Rorschach and the MMPI-2  
35  
36 in a Swedish Population: A Replication Study of the Effects of First-Factor Related Test-  
37  
38 Interaction Styles. *Journal of Personality Assessment*, 79, 357-370.  
39  
40  
41 Markon, K. E., Chmielewski, M., & Miller, C. J. (2011). The reliability and validity of discrete  
42  
43 and continuous measures of psychopathology: A quantitative review. *Psychological*  
44  
45 *Bulletin*, 137(5), 856-879.  
46  
47  
48 Meng, X. L., Rosenthal, R., & Rubin, D. B. (1992). Comparing correlated correlation  
49  
50 coefficients. *Psychological Bulletin*, 111, 172-175.  
51  
52  
53 Meyer, G. J. (1997). On the integration of personality assessment methods: The Rorschach and  
54  
55 MMPI-2. *Journal of Personality Assessment*, 68, 297-330.  
56  
57  
58  
59  
60

- 1  
2  
3 Meyer, G. J. (1999). The convergent validity of MMPI and Rorschach scales: An extension using  
4  
5 profile scores to define response-character styles on both methods and a re-examination  
6  
7 of simple Rorschach response frequency. *Journal of Personality Assessment*, 72, 1–35.  
8  
9  
10 Meyer, G. J. (2002). Exploring possible ethnic differences and bias in the Rorschach  
11  
12 Comprehensive System. *Journal of Personality Assessment*, 78, 104-129.  
13  
14  
15 Meyer, G. J., & Archer, R. P. (2001). The hard science of Rorschach research: What do we know  
16  
17 and where do we go? *Psychological Assessment*, 13, 486-502.  
18  
19  
20 Meyer, G. J., & Resnick, G. D. (1996). *Assessing ego impairment: Do scoring procedures make*  
21  
22 *a difference?* Paper presented at the 15th International Research Conference, Boston.  
23  
24  
25 Meyer, G. J., Hilsenroth, M. J., Baxter, D., Exner, J. E., Fowler, C. J., Piers, C. C., & Resnick, J.  
26  
27 (2002). An examination of interrater reliability for scoring the Rorschach Comprehensive  
28  
29 System in eight data sets. *Journal of Personality Assessment*, 78(2), 219–274.  
30  
31  
32 Meyer, G. J., Hsiao, W.-C., Viglione, D. J., Mihura, J. L., & Abraham, L. H. (2012). A survey of  
33  
34 applied clinical validity for the Rorschach. Manuscript provisionally accepted for  
35  
36 publication at *Journal of Personality Assessment*.  
37  
38  
39 Meyer, G. J., Riethmiller, R. J., Brooks, G. D., Benoit, W. A., & Handler, L. (2000). A  
40  
41 replication of Rorschach and MMPI-2 convergent validity. *Journal of Personality*  
42  
43 *Assessment*, 74, 175-215.  
44  
45  
46 Meyer, G. J., Viglione, D. J., Mihura, J., Erard, R. E., & Erdberg, P. (2011). *Rorschach*  
47  
48 *Performance Assessment System: Administration, coding, interpretation, and technical*  
49  
50 *manual*. Toledo, OH: Rorschach Performance Assessment System.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Mihura, J. L., Meyer, G. J., Dumitrascu, N., & Bombel, G. (2012). *A systematic review and*  
4  
5 *meta-analysis of the Rorschach Comprehensive System validity literature*. Manuscript re-  
6  
7 submitted for publication.  
8  
9  
10 Raghunathan, Rosenthal, & Rubin (1996). Comparing correlated but nonoverlapping  
11  
12 correlations. *Psychological Methods, 1*, 178-183.  
13  
14  
15 Viglione, D. J., & Hilsenroth, M. (2001). The Rorschach: facts, fiction, and future. *Psychological*  
16  
17 *Assessment, 13*, 452-471.  
18  
19  
20 Viglione, D. J., & Meyer G. J. (2008). An overview of Rorschach psychometrics for forensic  
21  
22 practice. In C.B Gacono & F.B. Evans with N. Kaser-Boyd (Eds.) *Handbook of Forensic*  
23  
24 *Rorschach psychology*. (pp. 22-54). Matwah, NJ: Lawrence Erlbaum Associates.  
25  
26  
27 Viglione, D. J., & Taylor, N. (2003). Empirical support for interrater reliability of the Rorschach  
28  
29 Comprehensive System coding. *Journal of Clinical Psychology, 59*(1), 111-121.  
30  
31  
32 Viglione, D. J., Blume-Marcovici, A., Miller, H., Giromini, L. & Meyer G. J. (2012) An Initial  
33  
34 Inter-Rater Reliability Study for the Rorschach Performance Assessment System. (In  
35  
36 submission).  
37  
38  
39 Viglione, D.J. (1999). A review of recent research addressing the utility of the Rorschach.  
40  
41 *Psychological Assessment, 11*, 251-265.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1

*Descriptive Data for Responses, Original CS Indices, and New Dimensional Composites*

Variable	Developmental (N=460)				Validation (N=230)			
	Mean R = 24.29 (SD = 4.68)				Mean R = 23.67 (SD = 4.51)			
	M	SD	Skew	Kurtosis	M	SD	Skew	Kurtosis
PTI	1.19	1.37	0.95	-0.12	1.26	1.30	0.83	-0.02
PT-Comp	1.19	1.21	0.66	0.52	1.28	1.20	0.68	1.17
HVI	3.33	1.61	0.20	-0.53	3.55	1.73	0.36	-0.62
V-Comp	3.33	1.39	0.17	0.02	3.50	1.50	0.20	0.18
S-CON	4.80	1.64	0.34	0.04	4.83	1.71	0.07	-0.43
SC-Comp	4.81	1.37	0.65	0.46	4.90	1.49	0.73	0.96

Table 2

*Individual Variables Used in the Original CS Indices and in the Multiple Regression Equations to Create Dimensional Composite*

*Alternatives*

Original Steps in the CS Indices	Alternative Variables in the Composite Equations	Final Regression Coefficients		
		Standardized $\beta$	raw b	p
PTI <sup>a</sup>				
<input type="checkbox"/> XA% < .70 & WDA% < .75	<i>Constant</i>		1.939	<.001
<input type="checkbox"/> X-% > .29	WDA%	-.217	-2.478	<.001
<input type="checkbox"/> Level 2 Special Scores > 2 & FAB2 > 0	X-%	.459	5.396	<.001
<input type="checkbox"/> R < 17 & WSum6 > .12 <i>or</i> R > 16 & WSum6 > 17	-1/(FAB2+1) <sup>2</sup>	.119	.586	<.001
<input type="checkbox"/> M- > 1 <i>or</i> X-% > .40	R	-.055	-.016	=.022
	$\sqrt{\text{WSum6}}$	.274	.186	<.001
	$\sqrt{\text{M-}}$	.175	.335	<.001
HVI				
	<i>Constant</i>		.332	.272
<input type="checkbox"/> FT+TF+T = 0	$\sqrt{\text{SumT}}$	-.266	-.631	<.001
<input type="checkbox"/> Zf > 12	Zf	.199	.065	<.001
<input type="checkbox"/> Zd > 3.5	Loc_DQ-Complexity/R	.121	.699	<.001
<input type="checkbox"/> S > 3	S	.323	.224	<.001
<input type="checkbox"/> H+(H)+Hd+(Hd) > 6	$\sqrt{[\text{H}+(\text{H})+\text{Hd}+(\text{Hd})]}$	.286	.652	<.001
<input type="checkbox"/> (H)+(A)+(Hd)+(Ad) > 3	$\sqrt{[(\text{H})+(\text{A})+(\text{Hd})+(\text{Ad})]}$	.182	.388	<.001
<input type="checkbox"/> H+A : Hd+Ad < 4:1	$[\text{H}+(\text{H})+\text{A}+(\text{A})]/[\text{H}+(\text{H})+\text{A}+(\text{A})+\text{Hd}+(\text{Hd})+\text{Ad}+(\text{Ad})]$	-.208	-2.340	<.001
<input type="checkbox"/> Cg > 3	$\sqrt{\text{Cg}}$	.134	.278	<.001
S-CON <sup>b</sup>				

## Rorschach Composites 21

	Constant		2.020	<.001
□ FV+VF+V+FD>2	$\sqrt{(\text{SumV}+\text{FD})}$	.157	.322	<.001
□ C. Sh Blends>0	$\sqrt{\text{C.Sh Blends}}$	.351	.896	<.001
□ Ego<.31 or >.44	$\sqrt{ \text{Ego} - .375 }$	.192	2.123	<.001
□ MOR>3	$\sqrt{\text{MOR}}$	.129	.263	<.001
□ Zd>3.5 or <-3.5	$\sqrt{ \text{Loc\_DQ-Complexity/R} - 1.373 }$	.096	.882	<.001
□ Es>EA	es-EA	.196	.065	<.001
□ CF+C>FC	(CF+C)-FC	.244	.152	<.001
□ X+%<.70	X+%	-.078	-1.031	=.004
□ S>3	S	.251	.177	<.001
□ P<3 or >8	$\sqrt{ \text{P} - 5.5 }$	.158	.589	<.001
□ PureH<2	H	-.156	-.128	<.001
□ R<17				

The final regression equations are derived by multiplying each variable by its raw b value, summing the results, and adding the constant. For instance, the TP-Comp score is obtained from this equation: TP-Comp = 1.939 - 2.478\*(WDA%) + 5.396\*(X-%) + .586\*(-1/(FAB2+1)<sup>2</sup>) - .016\*(R) + .186\*( $\sqrt{\text{WSum6}}$ ) +.335\*( $\sqrt{\text{MQ-}}$ ).

a. -1/(Lvl2+1)<sup>2</sup> was not significant and it was removed from the analysis; XA% was excluded because it is essentially the complement of X-%. In the developmental sample it was correlated with X-% at -.960 (p < .001);

b. R was not significant and it was removed from the analysis.

Table 3

*Intraclass Correlation Coefficients for Interrater Reliability of the Dimensional Composites and Original Indices*

Sample	Variable	ICC	ZPF	p	r Diff
Viglione et al. (2012) Sample; R-Opt Collected Data (N = 50) <sup>a</sup> .	PTI	.819	2.538	.011	.352
	TP-Comp	.909			
	S-CON	.809	.456	.648	.074
	SC-Comp	.833			
Viglione & Taylor (2003) Sample; Non Modeled Data (N = 84) <sup>b</sup> .	HVI	.918	3.289	.001	.462
	V-Comp	.969			
	PTI	.768	3.768	<.001	.381
	TP-Comp	.889			
Meyer et al. (2002) Sample; Non Modeled Data (N = 219) <sup>c</sup> .	S-CON	.876	2.418	.016	.291
	SC-Comp	.930			
	HVI	.947	4.586	<.001	.479
	V-Comp	.981			
Meyer et al. (2002) Sample; R-Opt Modeled Data (N = 93).	PTI	.917	1.105	.269	.074
	TP-Comp	.928			
	S-CON	.844	5.829	<.001	.381
	SC-Comp	.927			
Meyer et al. (2002) Sample; R-Opt Modeled Data (N = 93).	HVI	.910	6.051	<.001	.374
	V-Comp	.958			
	PTI	.885	2.740	.006	.275
	TP-Comp	.933			
Meyer et al. (2002) Sample; R-Opt Modeled Data (N = 93).	S-CON	.846	2.633	.008	.273
	SC-Comp	.909			
	HVI	.884	3.355	<.001	.371
V-Comp	.945				

Note: All individual ICCs are significant at  $p < .001$ .

a. ICCs for Composites were initially reported in Viglione et al. (2012)

b. ICCs for Indices were initially reported in Viglione & Taylor (2003)

c. ICCs for Indices were initially reported in Meyer et al. (2002)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Table 4

*Composite Correlations with Psychosis and Diagnostic Severity in the Chicago Sample*

	All Available Protocols—CS Administration				
	TP-Comp	PTI	Z	P	r Diff
Psychotic Diagnosis (N = 316)	.327	.275	2.026	0.043	0.057
Diagnostic Severity (N = 315)	.319	.283	1.400	0.161	0.040
	R-Opt Modeled Protocols				
	TP-Comp	PTI	Z	P	r Diff
Psychotic Diagnosis (N = 171)	.326	.312	0.404	0.686	0.016
Diagnostic Severity (N = 171)	.316	.309	0.202	0.840	0.008

Note: All correlations significant at  $p < .0005$ . Data derived from Meyer, 2002.

Table 5

Composite Correlations with Corresponding MMPI-2 Scales, N = 85

<i>Psychosis</i>					
<i>MMPI-2 Scales</i>	<i>PTI</i>	<i>TP-Comp</i>	<i>Z</i>	<i>p</i>	<i>r Diff</i>
Scale 8	.515***	.495***	-0.440	0.660	-0.027
BIZ	.444***	.452***	0.169	0.865	0.010
PSY-5-Psy	.453***	.457***	0.085	0.932	0.005
<i>Wariness</i>					
<i>MMPI-2 Scales</i>	<i>HVI</i>	<i>V-Comp</i>	<i>Z</i>	<i>p</i>	<i>r Diff</i>
Scale 6	.456***	.453***	-0.068	0.946	-0.004
CYN	.215*	.140	-1.537	0.124	-0.077
SOD	.386***	.375***	-0.241	0.810	-0.013
TRT2	.444***	.376***	-1.510	0.131	-0.082
<i>Affective Distress</i>					
<i>MMPI-2 Scales</i>	<i>SCON</i>	<i>SC-Comp</i>	<i>Z</i>	<i>p</i>	<i>r Diff</i>
Scale 2	.503***	.525***	0.360	0.718	0.030
Scale 7	.668***	.708***	0.800	0.424	0.076
DEP	.630***	.704***	1.435	0.151	0.133
ANX	.625***	.686***	1.163	0.245	0.107
PSY-5-Neg	.619***	.617***	-0.036	0.971	-0.003

a. Correlations were initially reported in Meyer (1999). However, two records with fewer than 14 responses were dropped so some of the coefficients differ slightly from those in the original publication.

\* $p < .05$ ; \*\*  $p < .01$ , \*\*\*  $p < .001$