



Cochrane
Library

Cochrane Database of Systematic Reviews

Exercise for intermittent claudication (Review)

Lane R, Harwood A, Watson L, Leng GC

Lane R, Harwood A, Watson L, Leng GC.
Exercise for intermittent claudication.
Cochrane Database of Systematic Reviews 2017, Issue 12. Art. No.: CD000990.
DOI: [10.1002/14651858.CD000990.pub4](https://doi.org/10.1002/14651858.CD000990.pub4).

www.cochranelibrary.com

Exercise for intermittent claudication (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	7
OBJECTIVES	8
METHODS	8
RESULTS	10
Figure 1.	10
Figure 2.	12
Figure 3.	13
Figure 4.	18
Figure 5.	19
Figure 6.	20
DISCUSSION	21
AUTHORS' CONCLUSIONS	23
ACKNOWLEDGEMENTS	23
REFERENCES	24
CHARACTERISTICS OF STUDIES	37
DATA AND ANALYSES	88
Analysis 1.1. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 1 Maximum walking distance (m).	89
Analysis 1.2. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 2 Pain-free walking distance (m).	89
Analysis 1.3. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 3 Maximum walking time (min).	90
Analysis 1.4. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 4 Pain-free walking time (min).	90
Analysis 1.5. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 5 Change in MWD/T.	91
Analysis 1.6. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 6 Change in ICD/T.	92
Analysis 1.7. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 7 Ankle brachial index.	92
Analysis 1.8. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 8 Mortality. ...	93
Analysis 1.9. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 9 Amputation.	93
Analysis 1.10. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 10 Peak exercise calf blood flow (mL/100 mL/min).	93
Analysis 2.1. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 1 Maximum walking distance (m).	95
Analysis 2.2. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 2 Pain-free walking distance (m).	95
Analysis 2.3. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 3 Maximum walking time (min).	95
Analysis 2.4. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 4 Pain-free walking time (min).	96
Analysis 2.5. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 5 Ankle brachial index.	96
Analysis 2.6. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 6 Mortality.	97
Analysis 2.7. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 7 Quality of Life SF-36.	97
Analysis 2.8. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 8 Peak exercise calf blood flow (mL/100 mL/min).	98

Analysis 3.1. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 1 Maximum walking distance (m).	100
Analysis 3.2. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 2 Pain-free walking distance (m).	100
Analysis 3.3. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 3 Maximum walking time (min).	100
Analysis 3.4. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 4 Pain-free walking time (min).	101
Analysis 3.5. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 5 Ankle brachial index.	101
Analysis 3.6. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 6 Mortality.	102
Analysis 3.7. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 7 Quality of Life SF-36.	102
Analysis 3.8. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 8 Peak exercise calf blood flow (mL/100 mL/min).	104
Analysis 4.1. Comparison 4 Exercise regimen compared with antiplatelet therapy, Outcome 1 Maximum walking time (min). ...	104
Analysis 4.2. Comparison 4 Exercise regimen compared with antiplatelet therapy, Outcome 2 Ankle brachial index.	105
Analysis 4.3. Comparison 4 Exercise regimen compared with antiplatelet therapy, Outcome 3 Peak exercise calf blood flow (mL/100 mL/min).	105
Analysis 5.1. Comparison 5 Exercise regimen compared with pentoxifylline therapy, Outcome 1 Maximum walking time (min). .	105
Analysis 6.1. Comparison 6 Exercise regimen compared with iloprost therapy, Outcome 1 Maximum walking distance (m).	106
Analysis 6.2. Comparison 6 Exercise regimen compared with iloprost therapy, Outcome 2 Pain-free walking distance (m).	106
Analysis 7.1. Comparison 7 Exercise regimen compared with pneumatic foot and calf compression, Outcome 1 Maximum walking distance (m).	106
Analysis 7.2. Comparison 7 Exercise regimen compared with pneumatic foot and calf compression, Outcome 2 Pain-free walking distance (m).	106
Analysis 7.3. Comparison 7 Exercise regimen compared with pneumatic foot and calf compression, Outcome 3 Mortality.	107
Analysis 8.1. Comparison 8 Exercise regimen compared with vitamin E, Outcome 1 3-monthly maximum walking time (min). ..	107
Analysis 8.2. Comparison 8 Exercise regimen compared with vitamin E, Outcome 2 6-monthly maximum walking time (min). ..	107
Analysis 8.3. Comparison 8 Exercise regimen compared with vitamin E, Outcome 3 3-monthly ABI.	107
Analysis 8.4. Comparison 8 Exercise regimen compared with vitamin E, Outcome 4 6-monthly ABI.	108
ADDITIONAL TABLES	108
APPENDICES	110
WHAT'S NEW	112
HISTORY	112
CONTRIBUTIONS OF AUTHORS	113
DECLARATIONS OF INTEREST	113
SOURCES OF SUPPORT	113
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	113
INDEX TERMS	113

[Intervention Review]

Exercise for intermittent claudication

Risha Lane¹, Amy Harwood¹, Lorna Watson², Gillian C Leng³

¹Vascular Unit, Hull Royal Infirmary, Hull, UK. ²NHS Fife, Leven, UK. ³National Institute for Health and Care Excellence, London, UK

Contact address: Risha Lane, Vascular Unit, Hull Royal Infirmary, Anlaby Road, Hull, HU3 2JZ, UK. risha@doctors.org.uk, risha@doctors.net.uk.

Editorial group: Cochrane Vascular Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 12, 2017.

Citation: Lane R, Harwood A, Watson L, Leng GC. Exercise for intermittent claudication. *Cochrane Database of Systematic Reviews* 2017, Issue 12. Art. No.: CD000990. DOI: [10.1002/14651858.CD000990.pub4](https://doi.org/10.1002/14651858.CD000990.pub4).

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Exercise programmes are a relatively inexpensive, low-risk option compared with other, more invasive therapies for treatment of leg pain on walking (intermittent claudication (IC)). This is the fourth update of a review first published in 1998.

Objectives

Our goal was to determine whether an exercise programme was effective in alleviating symptoms and increasing walking treadmill distances and walking times in people with intermittent claudication. Secondary objectives were to determine whether exercise was effective in preventing deterioration of underlying disease, reducing cardiovascular events, and improving quality of life.

Search methods

For this update, the Cochrane Vascular Information Specialist searched the Specialised Register (last searched 15 November 2016) and the Cochrane Central Register of Controlled Trials (CENTRAL; 2016, Issue 10) via the Cochrane Register of Studies Online, along with trials registries.

Selection criteria

Randomised controlled trials of an exercise regimen versus control or versus medical therapy for people with IC due to peripheral arterial disease (PAD). We included any exercise programme or regimen used for treatment of IC, such as walking, skipping, and running. Inclusion of trials was not affected by duration, frequency, or intensity of the exercise programme. Outcome measures collected included treadmill walking distance (time to onset of pain or pain-free walking distance and maximum walking time or maximum walking distance), ankle brachial index (ABI), quality of life, morbidity, or amputation; if none of these was reported, we did not include the trial in this review.

Data collection and analysis

For this update (2017), RAL and AH selected trials and extracted data independently. We assessed study quality by using the Cochrane 'Risk of bias' tool. We analysed continuous data by determining mean differences (MDs) and 95% confidence intervals (CIs), and dichotomous data by determining risk ratios (RRs) and 95% CIs. We pooled data using a fixed-effect model unless we identified significant heterogeneity, in which case we used a random-effects model. We used the GRADE approach to assess the overall quality of evidence supporting the outcomes assessed in this review.

Main results

We included two new studies in this update and identified additional publications for previously included studies, bringing the total number of studies meeting the inclusion criteria to 32, and involving a total of 1835 participants with stable leg pain. The follow-up period ranged from two weeks to two years. Types of exercise varied from strength training to polestriding and upper or lower limb exercises; supervised sessions were generally held at least twice a week. Most trials used a treadmill walking test for one of the primary outcome

measures. The methodological quality of included trials was moderate, mainly owing to absence of relevant information. Most trials were small and included 20 to 49 participants. Twenty-seven trials compared exercise versus usual care or placebo, and the five remaining trials compared exercise versus medication (pentoxifylline, iloprost, antiplatelet agents, and vitamin E) or pneumatic calf compression; we generally excluded people with various medical conditions or other pre-existing limitations to their exercise capacity.

Meta-analysis from nine studies with 391 participants showed overall improvement in pain-free walking distance in the exercise group compared with the no exercise group (MD 82.11 m, 95% CI 71.73 to 92.48, $P < 0.00001$, high-quality evidence). Data also showed benefit from exercise in improved maximum walking distance (MD 120.36 m, 95% CI 50.79 to 189.92, $P < 0.0007$, high-quality evidence), as revealed by pooling data from 10 studies with 500 participants. Improvements were seen for up to two years.

Exercise did not improve the ABI (MD 0.04, 95% CI 0.00 to 0.08, 13 trials, 570 participants, moderate-quality evidence). Limited data were available for the outcomes of mortality and amputation; trials provided no evidence of an effect of exercise, when compared with placebo or usual care, on mortality (RR 0.92, 95% CI 0.39 to 2.17, 5 trials, 540 participants, moderate-quality evidence) or amputation (RR 0.20, 95% CI 0.01 to 4.15, 1 trial, 177 participants, low-quality evidence).

Researchers measured quality of life using Short Form (SF)-36 at three and six months. At three months, the domains 'physical function', 'vitality', and 'role physical' improved with exercise; however this was a limited finding, as it was reported by only two trials. At six months, meta-analysis showed improvement in 'physical summary score' (MD 2.15, 95% CI 1.26 to 3.04, $P = 0.02$, 5 trials, 429 participants, moderate-quality evidence) and in 'mental summary score' (MD 3.76, 95% CI 2.70 to 4.82, $P < 0.01$, 4 trials, 343 participants, moderate-quality evidence) secondary to exercise. Two trials reported the remaining domains of the SF-36. Data showed improvements secondary to exercise in 'physical function' and 'general health'. The other domains - 'role physical', 'bodily pain', 'vitality', 'social', 'role emotional', and 'mental health' - did not show improvement at six months.

Evidence was generally limited in trials comparing exercise versus antiplatelet therapy, pentoxifylline, iloprost, vitamin E, and pneumatic foot and calf compression owing to small numbers of trials and participants.

Review authors used GRADE to assess the evidence presented in this review and determined that quality was moderate to high. Although results showed significant heterogeneity between trials, populations and outcomes were comparable overall, with findings relevant to the claudicant population. Results were pooled for large sample sizes - over 300 participants for most outcomes - using reproducible methods.

Authors' conclusions

High-quality evidence shows that exercise programmes provided important benefit compared with placebo or usual care in improving both pain-free and maximum walking distance in people with leg pain from IC who were considered to be fit for exercise intervention. Exercise did not improve ABI, and we found no evidence of an effect of exercise on amputation or mortality. Exercise may improve quality of life when compared with placebo or usual care. As time has progressed, the trials undertaken have begun to include exercise versus exercise or other modalities; therefore we can include fewer of the new trials in this update.

PLAIN LANGUAGE SUMMARY

Exercise for reducing intermittent claudication symptoms

Background

Intermittent claudication is a cramping leg pain that develops when walking and is relieved with rest. It is caused by inadequate blood flow to the leg muscles caused by atherosclerosis (fatty deposits restricting blood flow through the arteries). People with mild to moderate claudication are advised to keep walking, stop smoking, and reduce cardiovascular risk factors. Other treatments include antiplatelet therapy, pentoxifylline or cilostazol, angioplasty (inserting a balloon into the artery to open it up), and bypass surgery.

Studies and key results

Review authors identified 32 controlled trials that randomised 1835 adults with stable leg pain to exercise, usual care or placebo, or other interventions (current until November 2016). Researchers measured outcomes at times ranging from two weeks to two years. Types of exercise varied from strength training to polestriding and upper or lower limb exercises; in general, supervised sessions were held at least twice a week. The quality of included trials was moderate, mainly because of absence of relevant information. Ten trials reported that in the exercise groups, pain-free walking distance and the maximum distance that participants could walk was increased. Improvements were seen for up to two years. Exercise did not improve ankle to brachial blood pressure index. No evidence of an effect of exercise was seen on death or need for amputation because data were limited. Researchers assessed quality of life using the SF-36 Questionnaire at three and six months. At three months, indicators of quality of life - 'physical function', 'vitality', and 'role physical' - had all improved with exercise, but these data are limited, as only two trials reported this. Five studies reported improved 'physical summary score' and four studies reported improved 'mental health score' following exercise at six months, with two trials also reporting improvements in 'physical function' and 'general health'. All other domains showed no improvement at six months following exercise.

Comparisons of exercise with antiplatelet therapy, pentoxifylline, iloprost, vitamin E, and pneumatic foot and calf compression were limited because numbers of identified trials and participants were small.

Exercise for intermittent claudication (Review)

Quality of the evidence

The present review shows that exercise programmes appear to improve walking distance for people considered fit for exercise regimens. This benefit appears to be sustained over two years. Evidence presented in this review was of moderate to high quality. Although differences between trials were evident, populations and outcomes were comparable overall, and findings were relevant to people with intermittent claudication. Combined results were derived from large sample sizes - over 300 participants for most outcomes - using reproducible methods.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Is exercise an effective intervention in intermittent claudication?

Exercise compared with no exercise for intermittent claudication

Patient or population: adults with intermittent claudication

Settings: hospital or community-based physical therapy exercise programmes

Intervention: exercise^a

Comparison: no exercise (previously known as usual care)

Outcomes	Illustrative comparative risks* (95% CI) ^b		Relative effect (95% CI)	No. of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No exercise	Exercise ^a				
Pain-free walking distance (m) follow-up: 6 weeks to 24 months	Mean pain-free walking distance ranged across control groups from 63.3 m to 253 m.	Mean pain-free walking distance (m) in exercise groups ranged from 116 m to 413 m. MD was 82.11 m further in the exercise group. (95% CI 71.73 to 92.48)		391 (9 RCTs)	⊕⊕⊕⊕ high^c	4/9 studies (Cucato 2013; Gardner 2002; Jansen 1991 and Mika 2005) reported a clear improvement.
Maximum walking distance (m) follow-up: 6 weeks to 24 months	Mean maximum walking distance ranged across control groups from 122 m to 771 m.	Mean maximum walking distance (m) in exercise groups ranged from 136 m to 1100 m. MD was 120.36 m further in the exercise group. (95% CI 50.79 to 189.92)		500 (10 RCTs)	⊕⊕⊕⊕ high^d	5/10 studies (Cucato 2013; Gardner 2002; Jansen 1991; Tew 2015; Zwierska 2005) reported a clear improvement.
Ankle brachial index follow-up: 3 to 12 months	Mean ABI ranged across control groups from 0.32 to 0.89.	Mean ABI in exercise groups ranged from 0.34 to 0.96. MD was 0.04 higher in the exercise group. (95% CI 0.00 to 0.08)		570 (13 RCTs)	⊕⊕⊕⊖ moderate^e	A change in ABI of 0.04 is of limited clinical significance.

Mortality: all-cause deaths follow-up: 3 to 12 months	A total of 9/273 deaths occurred in the no exercise group.	A total of 8/267 deaths occurred in the exercise group.	RR 0.92 (0.39 to 2.17)	540 (5 RCTs)	⊕⊕⊕⊖ moderate^f
Amputation follow-up: 12 months	A total of 2/88 amputations occurred in the no exercise group.	No amputations occurred in the exercise group (0/89).	RR 0.20 (0.01 to 4.15)	177 (1 RCT)	⊕⊕⊖⊖ low^g
Quality of Life SF-36 Physical Summary score (scale 0 to 100, higher score indicates better quality of life) follow-up: 6 months	Mean physical summary score ranged across control groups from 34.2 to 47.1.	Mean Physical Summary score in exercise groups ranged from 41 to 54.1. MD score was 2.15 higher in the exercise group. (95% CI 1.26 to 3.04)		429 (5 RCTs)	⊕⊕⊕⊖ moderate^h
Quality of Life SF-36 Mental Summary score (scale 0 to 100, higher score indicates better quality of life) follow-up: 6 months	Mean Mental Summary score ranged across control groups from 38.8 to 62.0.	Mean Mental Summary score in exercise groups ranged from 39.6 to 67. MD score was 3.76 higher in the exercise group. (95% CI 2.7 to 4.82)		343 (4 RCTs)	⊕⊕⊕⊖ moderateⁱ

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

ABI: ankle brachial index; CI: confidence interval; MD: mean difference; RCT: randomised controlled trial; RR: risk ratio; SF: Short Form.

GRADE Working Group grades of evidence.

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

^aVariability in type, duration, and frequency of exercise programmes as prescribed. Exercise programmes included a duration of six weeks up to one year, generally twice or three times a week, varying in length between 30 minutes and one hour.

^bWhen possible, the no exercise (control) groups results were measured at the same time point and were used in the meta-analysis to calculate assumed risk.

^cThe level of evidence was high, as most trials were of high quality.

^dThe level of evidence was high, as most trials were of high quality. Heterogeneity was significant ($P = 0.01$, $I^2 = 89\%$), and CIs were broad (50.79 to 189.92). However further research is unlikely to change this.

- ^eWe downgraded by one step owing to significant heterogeneity ($P \leq 0.01$, $I^2 = 64\%$); however CIs were narrow (0.00 to 0.09). Given the size of the data set, it is unlikely that further data will demonstrate a difference between the two groups.
- ^fWe downgraded the evidence by one step, as analysis included only five small studies (of which one was underpowered) and showed wide confidence intervals (imprecision).
- ^gWe downgraded the evidence by two steps, as data are from a single study with wide CIs (imprecision) and events were few overall.
- ^hWe downgraded owing to heterogeneity ($P = 0.02$, $I^2 = 66\%$), narrow CI (1.26 to 3.04), and a symmetrical distribution on funnel plot. Other quality of life measures were used and have been described more fully in the additional [Table 1](#). Improvement was also seen in the following domains: vitality and role physical at three months, physical function, and general health at six months.
- ⁱWe downgraded owing to heterogeneity ($P < 0.01$, $I^2 = 87\%$), narrow CI (2.70 to 4.82), and a symmetrical distribution on funnel plot. Additional improvements in other domains may be seen in the future as more studies report their outcomes for quality of life scores.

BACKGROUND

Peripheral arterial disease is an important cause of morbidity and mortality for people in many Western countries. It is estimated that most adults have some degree of atherosclerosis by the time they reach middle age, and approximately 4% will develop intermittent claudication (Leng 1993). As the population ages, the prevalence of claudication will increase. Risk factors for development of lower limb arterial disease are similar to those for coronary heart disease, and include smoking, raised cholesterol levels, hypertension, and diabetes.

Several epidemiological studies have demonstrated an association between sedentary habits, functional decline, and worsening claudication (McDermott 2006; McDermott 2011). The impact of exercise therapy on physical functional ability, muscle strength, and walking times has drawn increased focus since a review was prepared in 2013 (Parmenter 2013). Over the past few years, further research into calf muscle strength has supported known changes in muscle architecture while enhancing focus on the impact of these changes in relation to walking distance and physical function. Researchers have described reduction in myofibre cross-sectional area, enzymes, and power (Clyne 1985; Dahllof 1974; King 2015; Koutakis 2015), along with alterations in gait (Gommans 2016; King 2012), among patients with intermittent claudication.

Description of the condition

Peripheral arterial disease covers a spectrum ranging from asymptomatic disease through to claudication, critical limb ischaemia, and finally limb loss. Within this spectrum, most people have relatively stable disease, termed 'claudication'. Intermittent claudication occurs secondary to atherosclerosis of the lower limb arteries, resulting from impaired blood flow. Whether at rest or when walking slowly, this reduction may go unnoticed; however during periods of exercise or walking with additional loads, for example, carrying while shopping, a cycle of pain requiring short rests occurs. This muscular, cramp-like tightening of the calf, the buttocks, or the foot on walking is known as 'claudication'.

Description of the intervention

Treatment options for intermittent claudication include bypass surgery, angioplasty, and drug therapy, but the mainstay of treatment for many patients with mild to moderate claudication remains advice to 'stop smoking and keep walking' (Housley 1988; NCGC 2012) while modifying cardiovascular risk factors.

Exercise therapy or programmes usually require a regular weekly commitment, lasting from six weeks to a year. In general, programmes are run twice or three times per week. Duration can vary; usually a minimum of 30 minutes is required per session.

How the intervention might work

Researchers have conducted numerous studies of exercise therapy using various regimens that differ in duration and intensity; many of these studies suggest that exercise can prove beneficial for individuals with intermittent claudication (Ernst 1992). Underlying mechanisms through which exercise may effect improvement include increased and more effective distribution of blood flow to the legs (Ernst 1987a), improved rheological characteristics of the blood (Ernst 1987), less reliance on anaerobic metabolism (Ruell 1984), and greater use of oxygen (Dahllof 1974).

A systematic review of the effect of exercise on lower limb haemodynamics in individuals with mild to moderate claudication identified 33 trials. In these trials, investigators reported no change when comparing effects of control versus exercise therapy on resting ankle brachial index (ABI), postexercise ABI, resting calf arterial blood flow, reactive hyperaemic blood flow post ischaemia, and resting toe systolic pressure (Parmenter 2011). An extensive non-systematic review of the literature focussed on exercise training in people with claudication and physiological changes associated with exercise (Haas 2012). This review discussed how regular exercise improves endothelial flow-mediated dilatation (FMD). Exercise training also improves FMD in those with claudication.

Exercise is proposed to improve walking among claudicants through angiogenesis. Underlying biomechanisms by which this may occur include ischaemia, sheer stress secondary to exercise, and remodelling of skeletal muscle (Hiatt 1994; Regensteiner 1993). Researchers have done extensive work on the changes that occur in skeletal muscle secondary to claudication. These changes can be summarised as a change in capillary density (Clyne 1985), as a ratio of type I to type II fibres (Clyne 1985; Hiatt 1994; Sjöström 1982), or as increases in arteriogenesis and mitochondrial activity. Trialists have described subsequent changes in skeletal muscle that can occur with training in humans (Lundgren 1989a; Terjung 1988; Wang 2009), as well as in animal models with artificial hindlimb stenoses (Yang 1991).

More recently, researchers have explored changes secondary to differing exercise programmes, while focussing on calpain activity, which has been associated with muscle atrophy in animal models. When trial authors explored effects of walking regimens versus strength training combined with walking, they noted that neither was seen to alter calpain activity (Delaney 2014).

The impact of peripheral arterial disease (PAD) on lower limb skeletal muscle becomes apparent when focussing on daily tasks or measuring strength. Individuals with PAD have reduced lower limb strength and ability or endurance for performing lower limb tasks, that is, knee flexion, dorsiflexion, or plantar flexion, when compared with healthy controls (Câmara 2012). In addition, people with PAD and reduced muscle density are more likely to have higher all-cause and cardiovascular disease mortality (McDermott 2012).

Clinicians have identified reduced muscle strength (Câmara 2012; Gohil 2013a; Parmenter 2013a), decreased walking distances (Parmenter 2013a), greater imbalance (Gohil 2013; Mockford 2014), and alterations in gait (King 2012; Koutakis 2010), secondary to PAD.

Why it is important to do this review

A meta-analysis of exercise rehabilitation programmes for claudication pain showed that exercise training to near-maximal pain for at least six months was beneficial in improving claudication (Gardner 1995). That review provided good evidence for the best type of exercise therapy but did not compare findings with non-exercised control groups. The present Cochrane review focusses on randomised controlled trials only and encompasses additional endpoints. The National Institute for Health and Care Excellence (NICE) (NCGC 2012) currently advocates that all patients with PAD should undergo an exercise programme as first-line treatment. Numerous other studies have echoed this advice (Haas 2012; Lauret 2012; Willigendael 2005). However, uptake remains low across the

UK. This review aims to add weight to the current body of available evidence recommending exercise, while demonstrating effects of different durations of exercise on walking distance and quality of life.

OBJECTIVES

Our goal was to determine whether an exercise programme was effective in alleviating symptoms and increasing walking treadmill distances and walking times in people with intermittent claudication. Secondary objectives were to determine whether exercise was effective in preventing deterioration of underlying disease, reducing cardiovascular events and improving quality of life.

METHODS

Criteria for considering studies for this review

Types of studies

We included all randomised controlled trials (RCTs) of an exercise regimen versus control, or versus medical therapy. We excluded trials that used alternation (e.g. allocation by date of birth or days of the week). We included trials that did not perform intention-to-treat analysis, provided all randomised participants were accounted for. We excluded trials for which numerical data were not available in a usable format, despite contact with study authors, but that were otherwise suitable.

We included studies in this review if they focussed on exercise versus usual care or other medical interventions. We excluded studies focussing on exercise compared with other forms of exercise, unsupervised exercise, angioplasty, or surgery, to prevent overlap with other Cochrane reviews ([Antoniou 2017](#); [Fakhry 2013](#); [Fokkenrood 2013](#); [Fowkes 1998](#)). We excluded studies in which usual care included walking advice or suggestions to increase daily exercise, as this advice constitutes unsupervised exercise. Although the exclusion of walking advice as part of usual care may be deemed controversial, this has been undertaken according to what has been reported by the trialists in the study papers.

Types of participants

We included trials involving participants with symptomatic intermittent claudication due to atherosclerotic disease. Intermittent claudication may be diagnosed objectively by an ABI < 0.9 or evidence of PAD on Doppler ultrasound or angiography, or both, or by questionnaire or clinically if objective measures such as ABI or imaging were not used or reported. We excluded studies of participants with asymptomatic lower limb atherosclerosis that was identified by testing.

Types of interventions

We included any exercise programme used for treatment of patients with intermittent claudication, such as walking, skipping, and running, and home-based therapies provided researchers compared treatment against placebo or no therapy. Inclusion of trials was not affected by duration, frequency, or intensity of the exercise programme, but these issues were taken into account in the meta-analysis. This review did not consider supervised versus unsupervised exercise because this is the topic of another Cochrane review ([Fokkenrood 2013](#)). Walking advice provided by consultants in clinic can be seen as the best medical treatment or

control. However, this may also be deemed unsupervised exercise. Therefore, we excluded from this review all studies that provided best medical treatment, which includes walking advice.

To avoid overlap with other Cochrane reviews ([Antoniou 2017](#); [Fakhry 2013](#); [Fowkes 1998](#)), this review excluded all modalities by which exercise can be compared with percutaneous transluminal angioplasty (PTA) or surgery. We therefore reviewed studies originally included in this review comparing exercise versus PTA or surgery, and excluded them when appropriate.

Types of outcome measures

We included only studies that reported one or more of the following outcome measures: treadmill walking distance (time to onset of pain or pain-free walking distance and maximum walking time or maximum walking distance), ABI, quality of life, morbidity, or amputation.

Primary outcomes

- Treadmill walking distance (time to onset of pain or pain-free walking distance and maximum walking time or maximum walking distance)

Secondary outcomes

- ABI
- Mortality
- Amputation
- Quality of life (includes QoL measured by the Short Form (SF)-36 Questionnaire and other validated measurements such as the EuroQoL Group Quality of Life Questionnaire based on five dimensions (EQ-5D), the Vascular Quality of Life Questionnaire (VascuQoL), and the Intermittent Claudication Questionnaire)
- Peak exercise blood flow
- Cardiovascular events

Search methods for identification of studies

Electronic searches

For this update, the Cochrane Vascular Information Specialist (CIS) searched the following databases for relevant trials.

- Cochrane Vascular Specialised Register (15 November 2016).
- Cochrane Central Register of Controlled Trials (CENTRAL; 2016, Issue 10) via the Cochrane Register of Studies Online.

See [Appendix 1](#) for details of the search strategy used to search CENTRAL.

The CIS maintains the Cochrane Vascular Specialised Register, which is constructed from weekly electronic searches of MEDLINE Ovid, Embase, Ovid, the Cumulative Index to Nursing and Allied Health Literature (CINAHL), and the Allied and Complementary Medicine Database (AMED), and through handsearching of relevant journals. The full list of databases, journals, and conference proceedings searched, as well as the search strategies used, is presented in the [Specialised Register](#) section of the Cochrane Vascular Module in the Cochrane Library (www.cochranelibrary.com).

The CIS searched the following trial registries for details of ongoing and unpublished studies.

- ClinicalTrials.gov (www.clinicaltrials.gov).
- World Health Organization International Clinical Trials Registry Platform (www.who.int/trialsearch).
- ISRCTN Register (www.isrctn.com/).

See [Appendix 2](#).

Searching other resources

We checked the reference lists of relevant studies retrieved via electronic searches.

Data collection and analysis

Selection of studies

For this update, one of the review authors (RAL) independently identified relevant trials and determined their eligibility for inclusion in the review; another review author (AH) checked this work. We resolved disagreements by discussion or by consultation with a third review author (GL); however this was not required. As necessary, we sought additional information from authors of all trials that appeared to meet the inclusion criteria.

Data extraction and management

Two review authors (RAL and AH) independently extracted data. Review authors resolved disagreements by discussion and included the final results in the review.

Assessment of risk of bias in included studies

For this update (2017), two review authors (RAL and AH) independently performed risk of bias assessments. We discussed discrepancies, and we planned that if we could not reach agreement, we would ask a third review author (GL) to assess the trial. Risk of bias assessment comprises seven domains: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and the additional option of assessing any other aspects deemed to produce bias.

Measures of treatment effect

When appropriate, we pooled trial results in a statistical meta-analysis using guidelines published by Cochrane Vascular. We analysed continuous data by determining mean differences (MDs) and 95% confidence intervals (CIs) using a fixed-effect model. When significant heterogeneity was present, we used a random-effects model. We analysed dichotomous data by determining risk ratios (RRs) and 95% CIs using a fixed-effect model, unless we suspected heterogeneity. In addition, for the primary outcome measure treadmill walking distance (time to onset of pain or pain-free walking distance and maximum walking time or maximum walking distance), we also analysed percentage change by using the 95% CI.

Unit of analysis issues

We analysed all data by using means and standard deviations. When original study papers did not provide these, we calculated them using the method recommended by the statistician for Cochrane Vascular. The unit of analysis was the individual participant.

Dealing with missing data

When data were not available from the original study paper, we contacted study authors to request the relevant data. If these were not available, we excluded the paper from the review. If data for at least one predefined outcome of the review were available, we included the study and examined available data in the meta-analysis.

Assessment of heterogeneity

We subjectively tested heterogeneity between trial results by using clinical judgement of differences in patient populations, interventions (including type, duration, and intensity of exercise programmes), and outcome assessments. We assessed heterogeneity statistically by using the Chi² test and the I² statistic. We deemed heterogeneity as significant if the P value of the Chi² test was less than 0.01, or if I² was greater than 70%. An I² of 50% to 70% equated to moderate heterogeneity.

Assessment of reporting biases

When we identified sufficient studies (> 10), we assessed publication bias using a funnel plot. For meta-analyses, when the number of studies was less than 10, we did not use funnel plots, as this would have led to reduced power and inability to differentiate artefactual from true asymmetry.

Data synthesis

For this update, two review authors (RAL and AH) independently collected and pooled data, provided agreement was met. We then uploaded data into Review Manager 5 ([RevMan 2014](#)) software for analysis. We performed meta-analysis by using a fixed-effect model unless we detected heterogeneity (Chi² test P < 0.01); we used a random-effects model in the analysis if heterogeneity was present.

Subgroup analysis and investigation of heterogeneity

Owing to the numerous domains associated with assessment of quality of life as per the Medical Outcomes Study (MOS) Short Form (SF)-36, we analysed all domains by performing subgroup analysis. When possible, we also analysed and presented data by subgroups for usual care and placebo.

Sensitivity analysis

Exercise programmes consisted of 12-week interventions or 24-week interventions. In one case, the duration of the programme was two years. In view of the variable length of programmes, we separated and analysed data as two separate analyses - one for 12 weeks and one for 24 weeks.

'Summary of findings' table

For this update we included a table summarising the findings presented by this review for our main comparison of exercise versus no exercise in adults with symptomatic intermittent claudication. The study population continues to be at low risk with regards to their claudication; however, this is only one aspect of a progressive debilitating disease that can lead to limb loss and is accompanied by the coexistent risk of cardiovascular disease. We selected for inclusion in [Summary of findings for the main comparison](#) the most important and clinically relevant outcomes thought to be essential for decision-making. We described these under [Types of outcome measures](#); they include pain-free walking distance, maximum

walking distance, ABI, mortality, amputation, and quality of life. We calculated assumed control intervention risks by using the mean number of events in control groups of selected studies for each outcome. We used the system developed by the GRADE Working Group in grading the quality of evidence as high, moderate, low, or very low, on the basis of within-study risk of bias, directness of evidence, heterogeneity, precision of effects estimates, and risk of publication bias (GRADE 2004). We used GRADEpro GDT (GRADEpro

GD T 2015) software to create [Summary of findings for the main comparison](#).


RESULTS

Description of studies

Results of the search

See [Figure 1](#).

Figure 1. Study flow diagram.



Included studies

See [Characteristics of included studies](#).

We included two additional trials for this update ([McGuigan 2001](#); [Tew 2015](#)). We also included five publications related to the previously included GOALS 2013 trial ([McDermott 2013](#); [McDermott 2013a](#); [McDermott 2014](#); [McDermott 2014a](#); [Rejeski 2014](#)). However only one of these publications provided data (on mortality at six months) that could be included within the results section ([McDermott 2014a](#)). This brings the total number of included studies to 32 ([Arosio 2001](#); [Castro-Sanchez 2013](#); [Ciuffetti 1994](#); [Collins 2005](#); [Crowther 2012](#); [Cucato 2013](#); [Dahllöf 1974](#); [Gardner 2002](#); [Gelin 2001](#); [GOALS 2013](#); [Guidon 2010](#); [Hiatt 1990](#); [Hiatt 1994](#); [Hobbs 2005](#); [Jansen 1991](#); [Kakkos 2005](#); [Larsen 1966](#); [Leicht 2011](#); [Mannarino 1991](#); [McDermott 2008](#); [McGuigan 2001](#); [Mika 2005](#); [Mika 2006](#); [Mika 2011](#); [Sanderson 2006](#); [Schlager 2011](#); [Tew 2009](#); [Tew 2015](#); [Tisi 1997](#); [Tsai 2002](#); [Wood 2006](#); [Zwierska 2005](#)) and the total number of participants to 1835.

Eighteen trials included fewer than 50 participants ([Arosio 2001](#); [Ciuffetti 1994](#); [Crowther 2012](#); [Cucato 2013](#); [Dahllöf 1974](#); [Guidon 2010](#); [Hiatt 1990](#); [Hiatt 1994](#); [Hobbs 2005](#); [Jansen 1991](#); [Kakkos 2005](#); [Larsen 1966](#); [Leicht 2011](#); [Mannarino 1991](#); [McGuigan 2001](#); [Sanderson 2006](#); [Schlager 2011](#); [Wood 2006](#)), ten between 50 and 100 participants ([Castro-Sanchez 2013](#); [Collins 2005](#); [Gardner 2002](#); [Mika 2005](#); [Mika 2006](#); [Mika 2011](#); [Tew 2009](#); [Tew 2015](#); [Tisi 1997](#); [Tsai 2002](#)), three over 100 participants ([GOALS 2013](#); [McDermott 2008](#); [Zwierska 2005](#)), and one over 200 participants ([Gelin 2001](#)).

Researchers compared exercise versus six different modes of treatment, the most common being usual care or placebo. Two early trials compared exercise versus placebo tablets ([Dahllöf 1974](#); [Larsen 1966](#)), but in more recent studies, trial authors used usual care as the control comparator ([Collins 2005](#); [Crowther 2012](#); [Gardner 2002](#); [Gelin 2001](#); [GOALS 2013](#); [Guidon 2010](#); [Hiatt 1990](#); [Hiatt 1994](#); [Jansen 1991](#); [Leicht 2011](#); [McDermott 2008](#); [McGuigan 2001](#); [Mika 2005](#); [Mika 2006](#); [Mika 2011](#); [Sanderson 2006](#); [Schlager 2011](#); [Tew 2009](#); [Tew 2015](#); [Tisi 1997](#); [Tsai 2002](#); [Wood 2006](#); [Zwierska 2005](#)). Investigators compared exercise with the following drug therapies: antiplatelet agents ([Mannarino 1991](#)), pentoxifylline ([Ciuffetti 1994](#)), iloprost ([Arosio 2001](#)), and vitamin E ([Collins 2005](#)). One study compared exercise versus pneumatic foot and calf compression ([Kakkos 2005](#)); one used a placebo treatment protocol of disconnected ultrasound electrotherapy ([Castro-Sanchez 2013](#)); and one used a 'stretching class' as usual care ([Cucato 2013](#)).

Inclusion and exclusion criteria of the included studies varied widely, but usually excluded people with serious comorbidities that would compromise an exercise programme, or would make it impractical.

We noted some variation in the exercise regimens used, although all recommended at least two sessions weekly. All specified some element of supervision, except the earliest trial ([Larsen 1966](#)), in which participants were simply advised to exercise at home and were given some training or walking regimen. Types of exercise varied from strength training to polestriding, cycling, and upper or lower limb exercises. We did not identify studies that included skipping or running. Duration of treatment generally fell within 3 to 12 months. Trialists measured outcomes at times ranging from 14 days to 2 years and reported variable compliance with exercise.

Nearly all trials used a treadmill walking test to assess one of the outcome measures, but results show considerable variation in outcomes. Some reported walking distance, and others reported walking times. Trial authors reported calf blood flow and ABI and often provided haematological and biochemical measures. Trialists provided little information about mortality, amputations, and fatal or non-fatal cardiovascular events. Eleven studies reported quality of life measures ([Collins 2005](#); [Gardner 2002](#); [Gelin 2001](#); [GOALS 2013](#); [Guidon 2010](#); [Kakkos 2005](#); [McDermott 2008](#); [Tew 2015](#); [Tisi 1997](#); [Tsai 2002](#); [Zwierska 2005](#)). As trials increasingly used the Short Form (SF)-36 Questionnaire, we have combined these results for meta-analysis in this review. We did not include in the meta-analysis trials that used an alternative quality of life assessment; we provided individual study findings in [Table 1](#).

We sought additional information from trialists in most of the included studies for the updated version of this review.

Excluded studies

See [Characteristics of excluded studies](#).

For this update (2017), we excluded 18 additional studies ([Aruna 2015](#); [Cucato 2015](#); [Dantas 2016](#); [Gardner 2014](#); [Gardner 2014a](#); [Gibbs 2013](#); [Guidon 2013](#); [Guirro 2015](#); [Kono 2013](#); [LIFE Study](#); [Mays 2015](#); [NCT02075502](#); [NCT02879019](#); [PROPEL study](#); [Rodrigues 2014](#); [Schlager 2011a](#); [Sonaglia 2013](#); [Ventura 1984](#)). We added one publication to a previously excluded study ([EXITPAD 2010](#)).

We excluded a total of 111 studies from the current review.

We excluded studies from this review because they compared percutaneous transluminal angioplasty versus exercise ([CLEVER 2009](#); [Creasy 1990](#); [Greenhalgh 2008](#); [Hobbs 2006](#); [Kruidenier 2011](#); [Mazari 2010](#); [Spronk 2009](#); [SUPER study](#)), surgery versus exercise ([Lundgren 1989](#)), different exercise regimens ([Allen 2010](#); [Andreozzi 2008](#); [Beutel 1985](#); [Buchwalsky 1974](#); [Cachovan 1999](#); [Cheetham 2004](#); [Choi 2012](#); [Collins 2012](#); [Cucato 2011](#); [Cucato 2011a](#); [Cucato 2015](#); [Dedes 2010](#); [Degischer 2002](#); [Fakhry 2011](#); [Gardner 2005](#); [Gardner 2012](#); [Gardner 2014](#); [Gardner 2014a](#); [Gottstein 1987](#); [Jones 1996](#); [Kiesewetter 1987](#); [Labs 1999](#); [Martinez 2009](#); [Nawaz 1999](#); [Nawaz 2001](#); [NCT01241747](#); [NCT02879019](#); [Nicolai 2010](#); [Nielsen 1977](#); [Parr 2009](#); [Patterson 1997](#); [Pinto 1997](#); [Riebe 2001](#); [Ritti-Dias 2010](#); [Rodrigues 2014](#); [Saleem 2011](#); [Savage 2001](#); [Scheffler 1991](#); [Slordahl 2005](#); [Sonaglia 2013](#); [Thomson 1999](#); [Zwierska 2004](#)), and exercise versus walking advice as the form of conservative best medical treatment ([Bronas 2011](#); [Crowther 2008](#); [EXERT 2009](#); [EXITPAD 2010](#); [Gardner 2011](#); [Gardner 2012](#); [Hodges 2008](#); [Mays 2015](#); [Nawaz 1999](#); [NCT02075502](#); [Nordanstig 2011](#); [Parr 2009](#); [Stewart 2008](#); [Wang 2008](#)). We excluded nine studies because relevant suitable numerical data were not available despite attempts to contact study authors ([Collins 2010](#); [Fowler 2002](#); [Gibbellini 2000](#); [Holm 1973](#); [Maejima 2005](#); [Schlager 2011a](#); [Streminski 1992](#); [Tebbutt 2011](#); [Ventura 1984](#)). We excluded the remainder of excluded studies because they did not fit the inclusion criteria (e.g. participants did not have intermittent claudication, not an RCT, insufficient evidence of randomly allocated population, no non-exercise control group) ([Aruna 2015](#); [Brotans 2011](#); [Bulling 1991](#); [Carmeli 2004](#); [Cina 1996](#); [Cunningham 2012](#); [Dahllöf 1976](#); [Dantas 2016](#); [Dittmar 1977](#); [Ericsson 1970](#); [Ernst 1987](#); [Ernst 1990](#); [Fitzgerald 1971](#); [Gibbs 2013](#); [Guidon 2013](#); [Guirro 2015](#); [Kono 2013](#); [Krause 1976](#); [Lee 2007](#); [Leon 2005](#); [Lepantalo 1984](#); [LIFE Study](#); [Mannarino 1988](#); [Mannarino 1989](#); [McDermott 2004](#); [NCT01065740](#);

Necker 2003; Presern-Strukelj 200; PROPEL study; Riccioni 2010; Richardson 1991; Schoneberger 1994; Silvestro 2002; Snabl 1958; Taft 2004; Treat-Jacobson 2012; Walker 2000; Waller 1988; Wang 2008; Winterfeld 1983).

Ongoing studies

We included within the ongoing studies section two studies that are still awaiting data: [NCT01231360](#) (completed 2014) and

[NCT01822457](#) (completed August 2016). See [Characteristics of ongoing studies](#).

Risk of bias in included studies

[Figure 2](#) and [Figure 3](#) provide an overall summary of bias present within each of the included studies (see also [Characteristics of included studies](#)). The high level of unclear bias was due to unclear reporting about sequence generation and allocation. We have expanded upon these aspects below.

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.




Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Arosio 2001	?	?	+	?	?	+	?
Castro-Sanchez 2013	+	+	+	+	+	-	+
Ciuffetti 1994	+	+	+	?	+	+	?
Collins 2005	+	+	+	?	+	+	+
Crowther 2012	?	+	+	?	-	+	-
Cucato 2013	?	?	+	?	?	+	?
Dahllof 1974	?	?	+	-	?	+	?
Gardner 2002	?	?	+	?	-	+	+
Gelin 2001	+	+	+	?	+	+	?
GOALS 2013	+	+	+	+	+	+	?
Guidon 2010	?	?	+	?	-	-	?
Hiatt 1990	+	-	+	?	-	-	?
Hiatt 1994	?	?	+	?	+	+	?
Hobbs 2005	+	+	+	?	+	+	?
Jansen 1991	?	?	+	?	?	+	+
Kakkos 2005	+	+	+	?	-	+	-
Larsen 1966	+	-	+	?	+	+	?
Leicht 2011	+	+	+	?	+	+	?
Mannarino 1991	?	?	+	?	?	+	?
McDermott 2008	+	+	+	+	+	+	?

Figure 3. (Continued)

McDermott 2008	+	+	+	+	+	+	?
McGuigan 2001	?	?	+	?	?	+	?
Mika 2005	?	?	+	+	+	+	+
Mika 2006	?	?	+	?	+	+	?
Mika 2011	?	?	+	+	+	+	+
Sanderson 2006	+	+	+	?	+	+	+
Schlager 2011	+	+	+	+	+	+	+
Tew 2009	+	+	+	?	+	+	?
Tew 2015	+	+	+	?	+	+	?
Tisi 1997	+	?	+	?	-	+	+
Tsai 2002	?	?	+	+	+	+	+
Wood 2006	?	?	+	?	+	+	?
Zwierska 2005	+	+	+	?	+	+	+

Allocation

All studies were RCTs, but many reports provided no details on random sequence generation other than a statement of 'randomised' (Arosio 2001; Crowther 2012; Cucato 2013; Dahllof 1974; Gardner 2002; Guidon 2010; Hiatt 1994; Jansen 1991; Mannarino 1991; McGuigan 2001; Mika 2005; Mika 2006; Mika 2011; Tsai 2002; Wood 2006). See Figure 3. To ensure that all trials were dealt with fairly, we deemed that any described as 'randomised' with no explanation as to how this was done had unclear risk of bias. We deemed that the remaining studies were at low risk of bias for random sequence generation.

A total of 15 studies did not report details of allocation concealment (Arosio 2001; Cucato 2013; Dahllof 1974; Gardner 2002; Guidon 2010; Hiatt 1994; Jansen 1991; Mannarino 1991; McGuigan 2001; Mika 2005; Mika 2006; Mika 2011; Tisi 1997; Tsai 2002; Wood 2006); we deemed these studies to be at unclear risk of selection bias. We deemed two studies to be at high risk of bias because the randomisation procedure was not blinded (Hiatt 1990; Larsen 1966). The remaining studies provided details of allocation concealment, and we judged them to be at low risk of bias.

Blinding

As the nature of exercise-based studies involved an activity versus standard care, medication, or an intervention, review authors deemed that blinding of participants was not possible. To ensure that all trials used a standardised approach, we scored all as having low risk of bias secondary to participant blinding. Inevitably in trials of exercise, blinding was not possible; therefore significant placebo responses may have occurred in trials comparing exercise versus usual care.

Blinding of participants was not possible, and included trials have additional risk of bias as outcome assessors may not be blinded to

the group to which a participant was randomised. Seven studies did specify that outcome assessors were blinded; we judged these to be at low risk of bias (Castro-Sanchez 2013; GOALS 2013; McDermott 2008; Mika 2005; Mika 2011; Schlager 2011; Tsai 2002). For some, however, blinding involved a separate aspect of the trial that was not focused on exercise, that is, vitamin E in Collins 2005, or carnitine analysis in Hiatt 1990; we judged these studies as having unclear risk. We deemed Dahllof 1974 to be at high risk of bias because the outcome assessor was not blinded to treatment groups. We judged the remainder of the included studies to be at unclear risk of bias owing to lack of reporting on blinding of outcome assessors.

Incomplete outcome data

Most trials reported no or minimal losses to follow-up (Arosio 2001; Castro-Sanchez 2013; Ciuffetti 1994; Collins 2005; Cucato 2013; Dahllof 1974; Gelin 2001; GOALS 2013; Hiatt 1994; Hobbs 2005; Jansen 1991; Larsen 1966; Leicht 2011; Mannarino 1991; McDermott 2008; Mika 2005; Mika 2006; Mika 2011; Sanderson 2006; Schlager 2011; Tew 2009; Tew 2015; Tsai 2002; Wood 2006; Zwierska 2005). We judged studies that had an attrition rate of 20% or more to be associated with higher risk of bias (Crowther 2012; Gardner 2002; Guidon 2010; Hiatt 1990; Kakkos 2005; McGuigan 2001; Tisi 1997).

We judged five studies to be at unclear risk of bias (Arosio 2001; Cucato 2013; Dahllof 1974; Jansen 1991; Mannarino 1991); Arosio 2001, Dahllof 1974, Jansen 1991, and Mannarino 1991 because investigators did not mention whether all enrolled participants completed the studies, and Cucato 2013 because only participants from the exercise group were lost to follow-up.

Selective reporting

All studies reported their prespecified outcome measures and were at low risk of reporting bias. Guidon 2010 discussed

results of the Walking Impairment Questionnaire (WIQ) and the Intermittent Claudication Questionnaire (ICQ) for 30 of 44 randomised participants; therefore, we judged this study to be at high risk of selective reporting bias. [Castro-Sanchez 2013](#) did not report on intermittent claudication distance; we therefore classified it as having high risk. [Hiatt 1990](#) did not specifically report on maximum walking distance or intermittent claudication in the treatment group but correlated it with treadmill performance; therefore, we classified this study as having high risk of reporting bias.

Other potential sources of bias

We initially labelled studies as having high risk of bias when they failed to meet sample size calculations ([Kakkos 2005](#); [McDermott 2008](#); [McGuigan 2001](#)). [Kakkos 2005](#) reported an attrition rate of 26% (8 of 34 discontinued); we therefore kept it at high risk. However, we reclassified [McDermott 2008](#) to unclear risk of bias, as the attrition rate was low; according to power calculations, 50 were needed in each group, and 50 completed the supervised exercise therapy (SET) whilst 48 in the control group completed the study. We also reclassified [McGuigan 2001](#) to unclear risk of bias as the reported attrition rate was low.

Reporting of treatment group numbers varied in the results section of [Crowther 2012](#); effects on outcomes were unclear. We therefore judged the study to be at high risk of bias.

Eleven studies were at low risk of bias ([Castro-Sanchez 2013](#); [Collins 2005](#); [Gardner 2002](#); [Jansen 1991](#); [Mika 2005](#); [Mika 2011](#); [Sanderson 2006](#); [Schlager 2011](#); [Tisi 1997](#); [Tsai 2002](#); [Zwierska 2005](#)).

Eleven studies included small sample sizes; we therefore deemed these studies to be at unclear risk of bias ([Arosio 2001](#); [Ciuffetti 1994](#); [Cucato 2013](#); [Dahllof 1974](#); [Hiatt 1990](#); [Hiatt 1994](#); [Hobbs 2005](#); [Larsen 1966](#); [Leicht 2011](#); [Mannarino 1991](#); [Wood 2006](#)). We deemed the remaining six studies to be at unclear risk of bias for a variety of other reasons ([Gelin 2001](#); [GOALS 2013](#); [Guidon 2010](#); [Mika 2006](#); [Tew 2009](#); [Tew 2015](#)). Please see the risk of bias tables for additional details.

Effects of interventions

See: [Summary of findings for the main comparison Is exercise an effective intervention in intermittent claudication?](#)

Exercise regimen compared with placebo or usual care

Overall outcomes

The wide range of reported time points meant that the overall analysis includes the last data time point at which data were presented in the study publications. This section shows a high degree of heterogeneity, with low heterogeneity noted in [Analysis 1.2](#) and [Analysis 1.8](#).

Maximum walking distance

(Analysis 1.1)

Ten trials with 500 participants reported this outcome and showed significant statistical heterogeneity ($I^2 = 89\%$, $P < 0.00001$) therefore we used a random-effects model ([Cucato 2013](#); [Gardner 2002](#); [Gelin 2001](#); [Hobbs 2005](#); [Jansen 1991](#); [Leicht 2011](#); [Schlager 2011](#); [Tew 2009](#); [Tew 2015](#); [Zwierska 2005](#)). The exercise group showed overall improvement in maximum walking distance (MD 120.36 metres,

95% CI 50.79 to 189.92, $P < 0.00007$, high quality of evidence as assessed via GRADE).

Pain-free walking distance

(Analysis 1.2)

Nine trials with a total of 391 participants reported on this outcome ([Cucato 2013](#); [Gardner 2002](#); [Hobbs 2005](#); [Jansen 1991](#); [Leicht 2011](#); [Mika 2005](#); [Tew 2009](#); [Tew 2015](#); [Zwierska 2005](#)) and noted improvement in pain-free walking distance in the exercise group (MD 82.11 metres, 95% CI 71.73 to 92.48, $P < 0.00001$, low quality of evidence as assessed via GRADE). Trials showed no significant heterogeneity ($I^2 = 41\%$, $P = 0.1$) therefore a fixed-effect model was used.

Maximum walking time

(Analysis 1.3)

Twelve studies with a total of 577 participants reported on maximum walking time ([Collins 2005](#); [Crowther 2012](#); [GOALS 2013](#); [Hiatt 1990](#); [Hiatt 1994](#); [Larsen 1966](#); [McDermott 2008](#); [Mika 2006](#); [Mika 2011](#); [Sanderson 2006](#); [Tsai 2002](#); [Wood 2006](#)). Data show overall improvement in walking time for those who underwent exercise (MD 4.51 minutes, 95% CI 3.11 to 5.92, $P < 0.00001$, high-quality evidence). Heterogeneity between studies was found to be significant ($I^2 = 82\%$, $P < 0.00001$) therefore a random-effects model was used. Walking time improved in eight trials ([Crowther 2012](#); [Hiatt 1990](#); [Hiatt 1994](#); [Larsen 1966](#); [McDermott 2008](#); [Mika 2006](#); [Mika 2011](#); [Tsai 2002](#)), which demonstrated exercise to be effective.

Pain-free walking time

(Analysis 1.4)

Eleven studies with a total of 534 participants reported on pain-free walking time ([Collins 2005](#); [Crowther 2012](#); [GOALS 2013](#); [Hiatt 1994](#); [Larsen 1966](#); [McDermott 2008](#); [Mika 2006](#); [Mika 2011](#); [Sanderson 2006](#); [Tsai 2002](#); [Wood 2006](#)). Meta-analysis demonstrated improvement in pain-free walking time for the exercise groups (MD 2.93 minutes, 95% CI 1.77 to 4.09, $P < 0.0001$, high-quality evidence). Heterogeneity between studies was found to be significant ($I^2 = 89\%$, $P < 0.00001$) therefore a random-effects model was used. Heterogeneity was most likely secondary to differences in reporting times and variable length of exercise programmes. Data show 100% improvement in pain-free walking time, which was likely to be of clinical significance. Only two of the studies found no real improvement with exercise ([Sanderson 2006](#); [Wood 2006](#)).

Percentage change in maximum walking distance or time

(Analysis 1.5)

We calculated the percentage change in maximum walking distance or time for 15 studies with a total of 656 participants ([Arosio 2001](#); [Crowther 2012](#); [Cucato 2013](#); [Gardner 2002](#); [GOALS 2013](#); [Hiatt 1990](#); [Hiatt 1994](#); [Kakkos 2005](#); [Leicht 2011](#); [Mannarino 1991](#); [McDermott 2008](#); [Mika 2006](#); [Tisi 1997](#); [Tsai 2002](#); [Wood 2006](#)). Meta-analysis reported overall improvement in maximum walking distance or time for those who underwent exercise (MD 40.25%, 95% CI 28.64 to 51.86, $P < 0.00001$). Heterogeneity between these studies was found to be significant ($P < 0.00001$, $I^2 = 93\%$) therefore a random-effects model was used.

Percentage change in pain-free walking distance (intermittent claudication distance (ICD)) or time (ICT)

(Analysis 1.6)

We calculated the percentage change in ICD or ICT for 15 studies with 703 participants (Arosio 2001; Collins 2005; Crowther 2012; Cucato 2013; Gardner 2002; GOALS 2013; Kakkos 2005; Mannarino 1991; McDermott 2008; Mika 2005; Mika 2006; Sanderson 2006; Tew 2009; Tsai 2002; Wood 2006). Results showed overall improvement in percentage change for ICD or ICT in favour of exercise (MD 58.42%, 95% CI 44.20 to 72.64, $P < 0.00001$). Heterogeneity between studies was found to be significant ($P < 0.001$, $I^2 = 70\%$) when a random-effects model was used.

Ankle brachial index (ABI)

(Analysis 1.7)

Thirteen trials with a total of 570 participants reported this outcome (Castro-Sanchez 2013; Collins 2005; Crowther 2012; Gardner 2002; Gelin 2001; Hiatt 1990; Hiatt 1994; Hobbs 2005; Leicht 2011; McGuigan 2001; Schlager 2011; Tew 2009; Tisi 1997). Meta-analysis showed a small change in ABI (MD 0.04, 95% CI 0.00 to 0.08, $P = 0.06$, moderate-quality evidence), which was supported by three studies (Castro-Sanchez 2013; Crowther 2012; Schlager 2011). Studies showed significant statistical heterogeneity ($P = 0.002$, $I^2 = 63\%$) therefore a random-effects model was used. Heterogeneity may be attributed to variations in exercise programme type or duration. Of note, as the number of trials included in updates has increased, the significance of change in ABI with exercise has diminished to show no improvement.

Mortality

(Analysis 1.8)

Five studies with 540 participants reported on mortality (Gelin 2001; GOALS 2013; McDermott 2008; Schlager 2011; Tew 2009). Results show no differences in effect between groups (risk ratio (RR) 0.92, 95% CI 0.39 to 2.17, $P = 0.85$, moderate-quality evidence). We updated these figures for this update, as recent published data included mortality figures for GOALS 2013. We noted no significant heterogeneity between trials ($I^2 = 0\%$, $P = 0.76$).

Amputation

(Analysis 1.9)

Only Gelin 2001 reported on amputation. Two amputations occurred in the usual care group, and none in the exercise group (RR 0.20, 95% CI 0.01 to 4.15, $P = 0.3$, low-quality evidence).

Quality of life

Overall, researchers reported quality of life (QoL) in numerous different ways (see Table 1). SF-36 provided the only consistently reported outcome for QoL, but this was reported at three and six months, rather than at one year. We have reported below results at these time points.

Other QoL questionnaires included the ICD Questionnaire (Tew 2015; Guidon 2010; Kakkos 2005), the Nottingham Health Profile (NHP) (Tisi 1997), and the Intermittent Claudication-Specific Sickness Impact Profile (SIP) scale (SIPIC) (Gelin 2001).

Tew 2015 used the ICD Questionnaire to assess changes between zero and six weeks. Data show no differences between intervention and control groups in EQ-5D score at six weeks, but improvement in ICD score in the intervention group (ICD score -10.6, 95% CI -18.9 to -2.3, $P < 0.05$) (see Table 1).

Cardiovascular events

None of the included studies reported on non-fatal cardiovascular events. We have examined all-cause mortality in Analysis 1.8.

Peak exercise calf blood flow

(Analysis 1.10)

We included in the meta-analysis for this outcome four studies with a total of 103 participants (Dahllof 1974; Gardner 2002; Hiatt 1990; Larsen 1966). Results show no significant overall improvement in blood flow between groups (MD 0.94 mL/100 mL/min, 95% CI -0.81 to 2.69, $P = 0.29$, low-quality evidence) ($I^2 = 59\%$, $P = 0.06$, moderate heterogeneity between trials). We downgraded the evidence owing to small sample size and wide confidence intervals.

Three-monthly outcomes

Seven trials provided one or more outcomes at the three-month time point for exercise compared with placebo or usual care (Collins 2005; Hiatt 1990; Hiatt 1994; Mika 2011; Schlager 2011; Tew 2009; Tsai 2002).

Maximum walking distance

(Analysis 2.1)

Three studies with 116 participants reported outcomes for maximum walking distance at the three-month time point (Cucato 2013; Schlager 2011; Tew 2009). Results showed no clear differences in distance between groups (MD 104.46 metres, 95% CI -64.33 to 273.24, $P = 0.23$) along with significant heterogeneity ($P = 0.008$, $I^2 = 79\%$) between studies when a random-effects model was used.

Pain-free walking distance

(Analysis 2.2)

Three trials with 156 participants provided pain-free walking distance outcomes at three months (Cucato 2013; Mika 2005; Tew 2009). Data showed significant improvement in the exercise group (MD 88.70 metres, 95% CI 58.25 to 119.15, $P < 0.00001$) ($I^2 = 16\%$, $P = 0.3$, no significant heterogeneity).

Maximum walking time

(Analysis 2.3)

Meta-analysis of five trials with 172 participants reporting maximum walking time at three months (Collins 2005; Hiatt 1990; Hiatt 1994; Mika 2011; Tsai 2002) showed improvement with exercise (MD 6.05 minutes, 95% CI 5.47 to 6.62, $P < 0.00001$) and no significant heterogeneity ($P = 0.62$, $I^2 = 0\%$).

Pain-free walking time

(Analysis 2.4)

Three trials with 132 participants provided data on pain-free walking time at three months (Hiatt 1994; Mika 2011; Tsai 2002). Meta-analysis demonstrated improvement in the exercise group

(MD 4.95 minutes, 95% CI 4.38 to 5.53, $P < 0.00001$) and significant heterogeneity between trials ($P = 0.02$, $I^2 = 73\%$).

Ankle brachial index (ABI)

(Analysis 2.5)

Four studies with a total of 130 participants reported on ABI outcomes at the three-month time point (Hiatt 1990; McGuigan 2001; Schlager 2011; Tew 2009). In contrast to the overall analysis, data show small differences in ABI between groups (MD 0.06, 95% CI 0.01 to 0.11, $P = 0.02$) ($P = 0.06$, $I^2 = 61\%$, moderate heterogeneity).

Mortality

(Analysis 2.6)

Tew 2009 and GOALS 2013 reported on this outcome for a total of 229 participants. No deaths occurred in either group in GOALS 2013, and Tew 2009 reported one death in the control group compared with none in the exercise group (RR 0.30, 95% CI 0.01 to 6.98).

Quality of life

(Analysis 2.7)

Tsai 2002 and Guidon 2010 reported a quality of life analysis for the SF-36 at three months. Domains found to improve secondary to exercise included 'physical function' (MD 6.60, 95% CI 2.37 to 10.83), 'vitality' (MD 5.55, 95% CI 1.54 to 9.56), and 'role physical' (MD 10.31, 95% CI 3.64 to 16.98). A random-effects model applied to assess for any variation revealed that the domain 'vitality' continued to show improvement secondary to exercise, but significance was lost for 'role physical' (MD 16.06, 95% CI -8.41 to 40.53) and 'physical function' (MD 5.95, 95% CI -2.45 to 14.34).

Peak exercise calf blood flow

(Analysis 2.8)

Only Hiatt 1990 reported on this outcome at three months and included 19 participants. Results show no clear differences between exercise and control groups at the end of the study (MD 2.50 mL/100 mL/min, 95% CI -1.49 to 6.49).

Six-monthly outcomes

Maximum walking distance

(Analysis 3.1)

Three studies with a total of 156 participants reported maximum walking distance at six months (Gardner 2002; Schlager 2011; Zwierska 2005). Results showed an increase in the exercise group (MD 138.36 metres, 95% CI 22.39 to 254.34, $P = 0.02$) and significant heterogeneity between studies ($P = 0.002$, $I^2 = 84\%$) when a random-effects model was used.

Pain-free walking distance

(Analysis 3.2)

Two studies with 116 participants reported on pain-free walking distance at six months (Gardner 2002; Zwierska 2005), noting improvement in the exercise group (MD 52.14 metres, 95% CI 6.83 to 97.45, $P = 0.02$) ($I^2 = 79\%$, $P = 0.03$, high heterogeneity).

Maximum walking time

(Analysis 3.3)

Four studies with a total of 295 participants reported outcomes for maximum walking time at six months (Crowther 2012; GOALS 2013; Larsen 1966; McDermott 2008). Meta-analysis showed improvement in favour of exercise (MD 3.20 minutes, 95% CI 2.04 to 4.36, $P < 0.00001$) and high heterogeneity ($P = 0.01$, $I^2 = 72\%$).

Pain-free walking time

(Analysis 3.4)

Five trials with 292 participants provided pain-free walking time outcomes at six months (Collins 2005; Crowther 2012; GOALS 2013; Larsen 1966; McDermott 2008), showing improvement with exercise (MD 2.32 minutes, 95% CI 0.91 to 3.74, $P < 0.001$) and significant heterogeneity between studies ($P = 0.005$, $I^2 = 73\%$) when a random-effects model was used.

Ankle brachial index (ABI)

(Analysis 3.5)

Six studies with 240 participants reported on ABI at six months (Castro-Sanchez 2013; Collins 2005; Crowther 2012; Gardner 2002; McGuigan 2001; Schlager 2011). Data showed no clear differences in ABI between the two groups (MD 0.05, 95% CI -0.01 to 0.11, $P = 0.07$) and significant heterogeneity ($P = 0.007$, $I^2 = 69\%$) between studies when a random-effects model was used.

Mortality

(Analysis 3.6)

Only GOALS 2013 provided mortality data for the six-month time point. Overall data show no clear difference in mortality between exercise and non-exercise groups at the six-month time period (RR 0.51, 95% CI 0.05 to 5.54, $P = 0.58$).

Quality of life

(Analysis 3.7)

Five studies with a total of 429 participants provided outcomes for the SF-36 generic quality of life measure (Collins 2005; Gardner 2002; GOALS 2013; McDermott 2008; Zwierska 2005), which showed improvement in quality of life secondary to exercise in certain domains, namely, 'physical summary score' (MD 2.15, 95% CI 1.26 to 3.04, $P < 0.00001$, moderate-quality evidence; supported with a random-effects model) and 'mental summary score' (MD 3.76, 95% CI 2.70 to 4.82, $P < 0.00001$, 4 studies, 343 participants, moderate-quality evidence). The 'mental summary score' no longer showed improvement when a random-effects model was applied (MD 2.85, 95% CI -1.01 to -6.71). Heterogeneity was significant (87%) for 'mental summary score'. Only two studies (43 participants) gave details on all domains; 'physical function' (MD 9.78, 95% CI 0.82 to 18.74) and 'general health' (MD 10.19, 95% CI 1.83 to 18.55) improved with exercise (Collins 2005; Zwierska 2005); this was supported again when a random-effects model was used. The other domains - 'role physical', 'bodily pain', 'vitality', 'social function', 'role emotional', and 'mental health' - did not show improvement.

Peak exercise calf blood flow

(Analysis 3.8)

Two studies with 66 participants reported on blood flow measurements at six months ([Gardner 2002](#); [Larsen 1966](#)). As with overall and three-month analyses, six-month blood flow measurements showed no differences between groups (MD 3.79 mL/100 mL/min, 95% CI 0.51 to 7.07, $P = 0.02$). Results showed no

significant heterogeneity ($P = 0.33$, $I^2 = 0\%$) between studies when a fixed-effect model was used.

Publication bias

We assessed overall publication bias using a funnel plot in three meta-analyses ([Analysis 1.3](#); [Analysis 1.4](#); [Analysis 1.7](#)) (see [Figure 4](#); [Figure 5](#); and [Figure 6](#)).

Figure 4. Funnel plot of comparison: 1 Overall outcomes: exercise regimen compared with placebo or usual care, outcome: 1.1 Maximum walking time (min).




Figure 5. Funnel plot of comparison: 1 Overall outcomes: exercise regimen compared with placebo or usual care, outcome: 1.4 Pain-free walking time (min).





Figure 6. Funnel plot of comparison: 1 Overall outcomes: Exercise regimen compared with placebo or usual care, outcome: 1.5 Ankle brachial index.



The funnel plot in [Figure 4](#) shows that 10 studies were within the 95% CI, and two were outside this range. Overall plot data were symmetrical, suggesting no evidence of publication bias or asymmetry secondary to the presence of smaller studies.

The funnel plot in [Figure 5](#) indicates that seven studies were within the 95% CI, and three were outside this range. Overall plot data were asymmetrical, suggesting bias, possibly attributed to the presence of smaller studies.

The funnel plot in [Figure 6](#) revealed that nine studies were within the 95% CI, and three were outside this range. Overall plot data were asymmetrical, suggesting bias, possibly attributed to the presence of smaller studies.

Exercise regimen compared with antiplatelet therapy

One trial involving 10 participants compared exercise with antiplatelet therapy ([Mannarino 1991](#)) and reported the following results.

Maximum walking time

([Analysis 4.1](#))

After six months of treatment, maximum walking time was improved in the exercise group compared with the group treated with antiplatelet therapy (MD 1.06 minutes, 95% CI 0.15 to 1.97). Maximum walking time was increased by 86% in the exercise group and by 38% in antiplatelet therapy group.

Ankle brachial index (ABI)

([Analysis 4.2](#))

Results show no differences between exercise and antiplatelet therapy groups at the end of the trial (MD 0.00, 95% CI -0.22 to 0.22).

Peak exercise calf blood flow

([Analysis 4.3](#))

Results show no clear differences in calf blood flow between antiplatelet therapy and exercise groups after six months, although flow tended to be higher in the exercise group (MD 2.18 mL/100 mL/min, 95% CI -0.28 to 4.64).

[Mannarino 1991](#) did not report the remaining outcomes of this review.

Exercise regimen compared with pentoxifylline

One trial involving 30 participants compared exercise versus pentoxifylline ([Ciuffetti 1994](#)) and reported the following results.

Maximum walking time

([Analysis 5.1](#))

After 13 weeks of therapy, maximum walking time was greater in the pentoxifylline group than in the exercise group (MD -0.45 minutes,

95% CI -0.66 to -0.24). Walking distance increased by 62% in the exercise group and by 88% in pentoxifylline group.

Adverse events

Two participants experienced gastroenteritis during treatment with pentoxifylline, but investigators did not consider this to be a side effect of study treatments.

Ciuffetti 1994 did not report the remaining outcomes of this review.

Exercise regimen compared with iloprost therapy

One study with 24 participants compared iloprost versus exercise (Arosio 2001) and reported the following results.

Pain-free walking distance

(Analysis 6.2)

Data showed improvement in pain-free walking distance in the exercise group at two weeks (MD 188.7 metres, 95% CI 15.38 to 362.02).

Maximum walking distance

(Analysis 6.1)

Data showed no clear effect on maximum walking distance in the exercise group at two weeks (MD 196.80 metres, 95% CI -83.8 to 477.40).

Arosio 2001 did not report the remaining outcomes of this review.

Exercise regimen compared with pneumatic foot and calf compression

One study with 25 participants compared exercise versus pneumatic foot and calf compression (Kakkos 2005).

Pain-free walking distance

(Analysis 7.2)

Data showed no clear effect on pain-free walking distance in the pneumatic compression group compared with the exercise group (MD -160.30 metres, 95% CI -438.88 to 118.28).

Maximum walking distance

(Analysis 7.1)

Data showed no clear effect on maximum walking distance in the pneumatic compression group compared with the exercise group (MD -61.90 metres, 95% CI -391.59 to 267.79).

Mortality

(Analysis 7.3)

One death occurred within the pneumatic calf compression group, and no deaths occurred in the exercise group (odds ratio (OR) 3.52, 95% CI 0.13 to 95.09).

Kakkos 2005 did not report the remaining outcomes of this review.

Exercise regimen compared with vitamin E

One study involving 24 participants compared effects of vitamin E and exercise (Collins 2005).

Maximum walking time

(Analysis 8.1; Analysis 8.2)

Collins 2005 demonstrated improvement secondary to exercise, which was greater than with vitamin E alone, at three and six months, respectively (MD 15.22 minutes, 95% CI 2.38 to 28.05; MD 22.60 minutes, 95% CI 8.05 to 37.15).

Ankle brachial index (ABI)

(Analysis 8.3; Analysis 8.4)

ABI was unchanged at three and six months, respectively (MD -0.03, 95% CI -0.21 to 0.15; MD 0.10, 95% CI -0.09 to 0.29).

Collins 2005 did not report the remaining outcomes of this review.

DISCUSSION

Summary of main results

The data presented in this update generally confirm the findings of previous versions of this review - that exercise has a significant positive effect on walking times and walking distances in people considered to be fit for exercise intervention, compared with placebo or usual care. Although most studies examined outcomes at three or six months, it is important to note that this benefit would appear to be sustained for up to two years (Jansen 1991). Of note, however, data show improvement in maximum walking distance at six months but not at three months.

Mean improvements in walking distance and walking time with exercise were clinically and statistically significant; however in most cases, the data were not normally distributed. Some individuals responded with improvement of considerably larger magnitude than the mean, whereas others responded less well, which may reflect varying compliance with exercise programmes. Successful programmes generally comprised physiotherapy with supervised exercise two or three times per week for 30 to 60 minutes, often with walking, leg exercises, or treadmill training. Some programmes encouraged additional home exercise.

We used percentage change from baseline to allow a more holistic understanding of change in walking ability. These additional parameters have demonstrated an increase in both initial claudication and maximum walking percentage changes, in keeping with changes documented for initial claudication walking distance and time, as well as for maximum walking distance and time. Of note, use of combined data allowed inclusion of 15 studies and a larger cohort of participants in the analysis.

Data related to ankle brachial index (ABI) and to other important outcomes are sparse. Investigators provided no data on non-fatal cardiovascular events and inconclusive data regarding mortality and amputation.

Antiplatelet therapy was less effective than exercise in improving walking distance and other measures of lower limb function, but researchers presented no data pertaining to fatal and non-fatal cardiovascular events. A previous meta-analysis has shown that antiplatelet agents reduce the incidence of cardiovascular events in people with claudication (Trialists 1994). Therefore, aspirin should be of benefit despite lack of effect on the lower limb.

The single small trial comparing pentoxifylline with exercise showed that participants on drug therapy had significantly longer walking distances after three months than those on exercise therapy (Ciuffetti 1994). Investigators reported no cardiovascular or adverse events among those on pentoxifylline, but disadvantages of drug treatment might include cost and lack of general cardiovascular improvement. Arosio 2001 showed that exercise improved pain-free walking time significantly more than iloprost, but this was a small study (24 participants).

Pneumatic foot and calf compression showed no clear effect on walking distances compared with exercise, again in a small trial (Kakkos 2005). However, alternatives for improving blood flow among those with poor mobility represent an important area for future research in those who are unsuitable for exercise programmes. Therefore further research into such avenues may yield more certain results and may therefore be of interest.

Researchers reported quality of life data on a range of scales, which made it difficult to incorporate this information into a meta-analysis. Results were variable depending on the comparisons made; this is a topic for future research.

Overall completeness and applicability of evidence

Overall, it is not contested that exercise may provide benefit for individuals with intermittent claudication. However, the overall benefit and duration of exercise programmes differ. A standardised approach with a minimum duration of six months appears to confer benefit and would improve both pain-free and maximum walking distance. Review authors found scant data on the benefits of a six-month programme compared with a three-month programme when measured outcomes focused on quality of life, mortality, or amputation risk. The general assumption is that exercise training is safe in peripheral arterial disease, but the data presented are often focussed on mortality and limb loss. Other cardiovascular morbidity appears to be under-reported, and additional details on safety data within exercise studies would be of value.

Criteria for participant selection resulted in exclusion of many individuals with stable claudication for whom exercise was not practical or safe owing to pre-existing medical conditions. As most people with intermittent claudication are elderly, comorbidities are common. For up to a third of people, exercise may not be a suitable option. Other research trials and reviews have examined optimum modes of exercise for people with intermittent claudication, for example, a Cochrane review explored supervised versus unsupervised exercise for intermittent claudication (Fokkenrood 2013). The present review cannot resolve uncertainties about different exercise regimens.

This review did not investigate the issue of non-compliance, which has a large impact on the use of supervised exercise classes and how funding should be allocated for this treatment as advocated by the National Institute for Health and Care Excellence (NICE) (NCGC 2012). Further information on the cost-effectiveness of exercise programmes is needed.

Quality of the evidence

We the review authors continue to believe that the quality of evidence is high in supporting exercise as first-line treatment for symptomatic claudicants. See [Summary of findings for the main](#)

[comparison](#). In this review update, we have added only two studies that support the body of literature already presented.

Compared with the previous version of this review, the evidence presented has improved in quality, as reporting criteria have become more stringent. In general, the bias present within papers was often difficult to assess owing to absence of relevant information. In these cases, we marked risk of bias as unclear, as evidence on which to base a judgement was insufficient. Overall, included studies were of moderate methodological quality.

Pooled outcomes for walking distance and maximum walking distance at all time points demonstrated significant heterogeneity. This most likely was secondary to pooling of walking times from some studies at two years and reporting of outcomes after three months in other studies. When possible we addressed this heterogeneity by analysing data at specific time points, such as three and six months post intervention. Regardless of this, data showed improvement in initial walking distance and time at three months, and in maximum walking distance and time by six months.

Publication bias remains an additional confounding factor for the quality of evidence pooled within this review. Older studies have been reported more poorly, as journal reporting criteria were previously less rigorous, resulting in greater inherent bias.

Potential biases in the review process

We searched the Specialised Register and the Cochrane Central Register of Controlled Trials (CENTRAL) for new publications about exercise in patients with intermittent claudication. Two review authors independently assessed all new studies in keeping with recommendations provided in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011).

The main cause of bias within this review is that walking advice provided by consultants in clinics is now seen as the best medical advice but may also be deemed as unsupervised exercise. Therefore this review excluded all studies that reported best medical treatment, which includes walking advice, as we believe this would overlap with the Cochrane review on supervised versus unsupervised exercise (Fokkenrood 2013). It remains possible that usual care included walking advice, but this was not explicitly stated by trialists in their study papers, suggesting one possible aspect of reporting bias within studies.

It is the review authors' feeling that as claudication research develops, a usual care non-exercise control group will cease to exist, as its inclusion will be deemed unethical.

Finally, earlier studies used treadmill walking distance as the primary outcome measure, and this has continued to serve as a method for assessment of improvement in claudication. Whilst treadmill distances remain a reliable means of assessment, this method is being superseded by other functional and subjective assessments. Newer assessment tools to assess changes in walking distance include GPS tracking, six-minute walks, subjective reporting, and assessments based on the Walking Impairment Questionnaire (WIQ). It is important that future reviews consider inclusion of these as outcome measures (Cucato 2013b; Tew 2013).

Agreements and disagreements with other studies or reviews

This review is consistent with previous versions ([Lane 2014](#); [Leng 2000](#); [Watson 2008](#)) and adds to the body of available evidence supporting exercise for people with intermittent claudication. This evidence is also in line with NICE guidelines, which recommend that exercise should be provided as first-line treatment for people with intermittent claudication ([NCGC 2012](#)). Other meta-analyses such as [Parmenter 2011](#) and other reviews such as [Lauret 2012](#) and [Lauret 2012a](#) also support this finding.

AUTHORS' CONCLUSIONS

Implications for practice

This review provides high-quality evidence showing that exercise therapy should play an important part in the care of selected patients with intermittent claudication, to improve walking times and distances. Effects were demonstrated following three months of supervised exercise, although some programmes lasted longer than one year. Limited data suggest that an effect is sustained for up to two years. Exercise did not improve ankle brachial index (ABI), and investigators detected no differences in the effect of exercise between groups in terms of amputation or mortality. Exercise may improve quality of life when compared with placebo or usual care.

Antiplatelet agents were less effective than exercise in improving walking distance but should continue to be used because of benefits in reducing cardiovascular events and death. In contrast, pentoxifylline was more effective than exercise but may have fewer beneficial effects on the cardiovascular system in general. Iloprost led to less improvement in walking time than exercise. Data show no clear effect on walking distances when pneumatic foot and calf compression was compared with exercise. However the number of participants in these studies is small and data are limited.

Implications for research

Important questions involve the degree of supervision required in any exercise regimen and how long any change in exercise habits can be expected to last. Therefore, a trial with long follow-

up - of five years - is needed to compare the effectiveness of different supervised and unsupervised regimens in terms of changing long-term exercise patterns. Behavioural changes and attitudes towards exercise have been items of key interest in areas such as cardiovascular rehabilitation ([Jolly 2009](#)). More recent studies have demonstrated that brief psychological interventions may play a key role in improving walking distances among patients with intermittent claudication up to one year ([Cunningham 2012](#)). A telling aspect of research into claudicants' beliefs is their poor overall understanding of the disease and of why walking is recommended ([Cunningham 2014](#)). Future holistic exercise programmes, which may include cognitive-behavioural therapy and lifestyle and risk factor modification, could provide great benefit in encouraging people to start and maintain a better overall lifestyle.

Future research should also focus on compliance with exercise and how this could be improved. Outcome measures should include fatal and non-fatal cardiovascular events. In addition, expansion of this review to assess the benefit of exercise for asymptomatic patients and its impact on cardiovascular morbidity is an important goal.

Further cost-effectiveness analysis is required to determine whether the cost of supervised sessions might offset the cost of deterioration in terms of surgery or occupation of in-patient beds for complications such as myocardial infarction.

A trial is needed to compare exercise treatment with pentoxifylline to determine whether the benefit of drug treatment is sustained over a longer period, and whether data show any differences in cardiovascular events. Further investigation of pneumatic foot and calf compression for treatment of individuals with intermittent claudication is also needed. Consistency in use of quality of life measures among different trials would be helpful in linking outcomes to patient-assessed improvements.

ACKNOWLEDGEMENTS

We are very grateful to Cochrane Vascular for assistance provided in the search for new trials and to Dr Brian Ellis for his input in previous versions of this review.

REFERENCES

References to studies included in this review

Arosio 2001 {published data only}

Arosio E, Cuzzolin L, De Marchi S, Minuz P, Degan M, Crivellente F, et al. Increased endogenous nitric oxide production induced by physical exercise in peripheral arterial occlusive disease patients. *Life Sciences* 1999;**65**(26):2815-22.

* Arosio E, Minuz P, Prior M, Zuliani V, Gaino S, De Marchi S, et al. Vascular adhesion molecule-1 and markers of platelet function before and after a treatment with iloprost or a supervised physical exercise program in patients with peripheral arterial disease. *Life Sciences* 2001;**69**(4):421-33.

Castro-Sanchez 2013 {published data only}

* Castro-Sanchez AM, Mataran-Penarrocha GA, Feriche-Fernandez-Castanys B, Fernandez-Sola C, Sanchez-Labraca N, Moreno-Lorenzo C. A program of 3 physical therapy modalities improves peripheral arterial disease in diabetes type 2 patients: a randomized controlled trial. *Journal of Cardiovascular Nursing* 2013;**28**(1):74-82.

Castro-Sanchez AM, Moreno-Lorenzo C, Mataran-Penarrocha GA, Feriche-Fernandez-Castanys B, Sanchez-Labraca N, Sanchez Joya Mdel M. Efficacy of a massage and exercise programme on the ankle-brachial index and blood pressure in patients with diabetes mellitus type 2 and peripheral arterial disease: a randomized clinical trial. [Spanish]. *Medicina Clinica* 2010;**134**(3):107-10.

Ciuffetti 1994 {published data only}

Ciuffetti G, Paltriccia R, Lombardini R, Lupattelli G, Pasqualini L, Mannarino E. Treating peripheral arterial occlusive disease: pentoxifylline vs exercise. *International Angiology* 1994;**13**(1):33-9.

Collins 2005 {published data only}

* Collins EG, Langbein WE, Orebaugh C, Bammert C, Hanson K, Reda D, et al. Cardiovascular training effect associated with polestriding exercise in patients with peripheral arterial disease. *Journal of Cardiovascular Nursing* 2005;**20**(3):177-85.

Collins EG, Langbein WE, Orebaugh C, Bammert C, Hanson K, Reda D, et al. Polestriding exercise and vitamin E for management of peripheral vascular disease. *Medicine and Science in Sports and Exercise* 2003;**35**(3):384-93.

Langbein WE, Collins EG, Orebaugh C, Littooy FN, Edwards L, Reda D. Polestriding exercise and vitamin E for management of claudication pain. *Journal of Rehabilitation Research and Development* 1998;**35**(Suppl 1):181-2.

Langbein WE, Collins EG, Orebaugh C, Maloney C, Williams KJ, Littooy FN, et al. Increasing exercise tolerance of persons limited by claudication pain using polestriding. *Journal of Vascular Surgery* 2002;**35**(5):887-93.

Crowther 2012 {published data only}

Crowther RG, Leicht AS, Spinks WL, Sangla K, Quigley F, Golledge J. Effects of a 6-month exercise program pilot study on walking economy, peak physiological characteristics,

and walking performance in patients with peripheral arterial disease. *Vascular Health and Risk Management* 2012;**8**:225-32.

Cucato 2013 {published data only}

Cucato GG, Chehuen Mda R, Costa LA, Ritti-Dias RM, Wolosker N, Saxton JM, et al. Exercise prescription using the heart of claudication pain onset in patients with intermittent claudication. *Clinics (Sao Paulo, Brazil)* 2013;**68**(7):974-8.

Dahllof 1974 {published data only}

Dahllof AG, Bjorntorp P, Holm J, Schersten T. Metabolic activity of skeletal muscle in patients with peripheral arterial insufficiency. *European Journal of Clinical Investigation* 1974;**4**(1):9-15.

Gardner 2002 {published data only}

Gardner AW, Katzel LI, Sorkin JD, Bradham DD, Hochberg MC, Flinn WR, et al. Exercise rehabilitation Improves functional outcomes and peripheral circulation in patients with intermittent claudication: a randomised controlled trial. *Journal of the American Geriatric Society* 2001;**49**(6):755-62.

* Gardner AW, Katzel LI, Sorkin JD, Goldberg AP. Effects of long-term exercise rehabilitation on claudication distances in patients with peripheral arterial disease: a randomized controlled trial. *Journal of Cardiopulmonary Rehabilitation* 2002;**22**(3):192-8.

Gelin 2001 {published data only}

* Gelin J, Jivegard L, Taft C, Karlsson J, Sullivan M, Dahllöf AG, et al. Treatment efficacy of intermittent claudication by surgical intervention, supervised physical exercise training compared to no treatment in unselected randomised patients I: one year results of functional and physiological improvements. *European Journal of Vascular and Endovascular Surgery* 2001;**22**(2):107-13.

Taft C, Karlsson J, Gelin J, Jivegard L, Sandström R, Arfvidsson B, et al. Treatment efficacy of intermittent claudication by invasive therapy, supervised physical exercise training compared to no treatment in unselected randomised patients II: one year results of health-related quality of life. *European Journal of Vascular and Endovascular Surgery* 2001;**22**(2):114-23.

GOALS 2013 {published data only}

McDermott M. A home-based exercise intervention significantly improves walking performance in peripheral arterial disease: one-year follow-up from a randomized controlled trial. *Circulation* 2013;**128**:A14789.

McDermott M, Criqui M, Domanchuk K, Guralnik J, Kibbe M, Liu K, et al. Home-based exercise improves walking speed and prevents mobility loss in peripheral artery disease: a randomized controlled trial. *Circulation* 2014; Vol. 130:A11744.

McDermott MM, Guralnik JM, Criqui MH, Ferrucci L, Liu K, Spring B, et al. Unsupervised exercise and mobility loss in peripheral artery disease: a randomized controlled trial. *Journal of the American Heart Association* 2015;**4**(5):e001659.

McDermott MM, Guralnik JM, Criqui MH, Ferrucci L, Zhao L, Liu K, et al. Home-based walking exercise in peripheral artery disease: 12-month follow-up of the GOALS randomized trial. *Journal of the American Heart Association* 2014;**3**(3):e000711.

McDermott MM, Liu K, Guralnik JM, Criqui MH, Spring B, Tian L, et al. Home-based walking exercise intervention in peripheral artery disease: a randomized clinical trial. *JAMA* 2013;**310**(1):57-65.

Rejeski WJS, Spring B, Domanchuk K, Tao H, Tian L, Zhao L, et al. A group-mediated, home-based physical activity intervention for patients with peripheral artery disease: effects on social and psychological function. *Journal of Translational Medicine* 2014;**12**(29):1-8.

Guidon 2010 {published data only}

Guidon M, McGee H. One-year effect of a supervised exercise programme on functional capacity and quality of life in peripheral arterial disease. *Disability and Rehabilitation* 2013;**35**(5):397-404.

Guidon MM, McGee H, Kelly C. Quality of life and functional capacity following peripheral arterial disease exercise programme. *European Journal of Cardiovascular Prevention and Rehabilitation* 2010;**17**:S102.

Hiatt 1990 {published data only}

Hiatt WR, Regensteiner JG, Hargarten ME, Wolfel EE, Brass EP. Benefit of exercise conditioning for patients with peripheral arterial disease. *Circulation* 1990;**81**(2):602-9.

Regensteiner JG, Steiner JF, Panzer RJ, Hiatt WR. Evaluation of walking impairment by questionnaire in patients with peripheral arterial disease. *Journal of Vascular Medicine and Biology* 1990;**2**:142-52.

Hiatt 1994 {published data only}

Hiatt WR, Regensteiner JG, Wolfel EE, Carry MR, Brass EP. Effect of exercise training on skeletal muscle histology and metabolism in peripheral arterial disease. *Journal of Applied Physiology* 1996;**81**(2):780-8.

* Hiatt WR, Wolfel EE, Meier RH, Regensteiner JG. Superiority of treadmill walking exercise versus strength training for patients with peripheral arterial disease. Implications for the mechanism of the training response. *Circulation* 1994;**90**(4):1866-74.

Regensteiner JG, Steiner JF, Hiatt WR. Exercise training improves functional status in patients with peripheral arterial disease. *Journal of Vascular Surgery* 1996;**23**(1):104-15.

Hobbs 2005 {published data only}

* Hobbs SD, Marshall T, Fegan C, Adam DJ, Bradbury AW. The effect of supervised exercise and cilostazol on coagulation and fibrinolysis in intermittent claudication: a randomised controlled trial. *Journal of Vascular Surgery* 2007;**45**(1):65-70, discussion 70.

Hobbs SD, Marshall T, Fegan C, Adam DJ, Bradbury AW. The effect of supervised exercise and cilostazol on coagulation and fibrinolysis in patients with intermittent claudication. Yearbook

2005, The Vascular Society of Great Britain & Ireland. 2005. Abstract 68.

Jansen 1991 {published data only}

Jansen T, Weiss T, Amendt K, Hsu E, Hubsch-Muller C, Diehm C. Effect of a 2-year ambulatory vascular sports program on walking distance in claudication patients - a controlled study [German]. *Vasa Supplementum* 1991;**33**:175.

Kakkos 2005 {published data only}

Kakkos SK, Geroulakos G, Nicolaides AN. Improvement of the walking ability in intermittent claudication due to superficial femoral artery occlusion with supervised exercise and pneumatic foot and calf compression: a randomised controlled trial. *European Journal of Vascular and Endovascular Surgery* 2005;**30**(2):164-75.

Larsen 1966 {published data only}

* Larsen OA, Lassen NA. Effect of daily muscular exercise in patients with intermittent claudication. *Lancet* 1966;**2**(7473):1093-6.

Larsen OA, Lassen NA. Effect of daily muscular exercise in patients with intermittent claudication. *Scandinavian Journal of Clinical and Laboratory Investigation* 1967;**Suppl 99**:168-71.

Leicht 2011 {published data only}

Leicht A, Crowther R, Golledge J. Influence of regular exercise on body fat and eating patterns of patients with intermittent claudication. *International Journal of Molecular Sciences* 2015;**16**(5):11339-54.

Leicht AS, Crowther RG, Golledge J. Influence of peripheral arterial disease and supervised walking on heart rate variability. *Journal of Vascular Surgery* 2011;**54**(5):1352-9.

Mannarino 1991 {published data only}

Mannarino E, Pasqualini L, Innocente S, Scricciolo V, Rignanese A, Ciuffetti G. Physical training and antiplatelet treatment in stage II peripheral arterial occlusive disease: alone or combined?. *Angiology* 1991;**42**(7):513-21.

McDermott 2008 {published data only}

McDermott MM, Ades P, Guralnik JM, Dyer A, Ferrucci L, Liu K, et al. Treadmill exercise and resistance training in patients with peripheral arterial disease with and without intermittent claudication: a randomized controlled trial. *JAMA* 2009;**301**(2):165-74, Erratum in *JAMA* 2012;**307**(16):1694.

* McDermott MM, Ades PA, Dyer A, Guralnik JM, Kibbe M, Criqui MH. Corridor-based functional performance measures correlate better with physical activity during daily life than treadmill measures in persons with peripheral arterial disease. *Journal of Vascular Surgery* 2008;**48**(5):1231-7.

NCT00106327. Improving functioning in peripheral arterial disease. clinicaltrials.gov/ct2/show/NCT00106327. Date first received: 22 March 2005.

Roitman JL. Treadmill exercise and resistance training in patients with peripheral arterial disease with and without intermittent claudication: a randomized controlled trial.

Journal of Cardiopulmonary Rehabilitation and Prevention 2010;**30**(1):62.

McGuigan 2001 {published data only}

McGuigan MR, Bronks R, Newton RU, Sharman MJ, Graham JC, Cody DV, et al. Resistance training in patients with peripheral arterial disease: effects on myosin isoforms, fiber type distribution, and capillary supply to skeletal muscle. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* 2001;**56**:B302-10.

Mika 2005 {published data only}

Mika P, Spodaryk K, Cencora A. Effects of treadmill training on walking distance and lower limb blood flow in patients with intermittent claudication [Zmiany dystansu marszu i przepływu tętniczego w kończynach dolnych podczas treningu marszowego u pacjentów z chropaniem przestankowym]. *Medical Rehabilitation* 2005;**9**(1):3-9.

* Mika P, Spodaryk K, Cencora A, Unnithan VB, Mika A. Experimental model of pain-free treadmill training in patients with claudication. *American Journal of Physical Medicine and Rehabilitation* 2005;**84**(10):756-62.

Mika 2006 {published data only}

* Mika P, Spodaryk K, Cencora A, Mika A. Red blood cell deformability in patients with claudication after pain-free treadmill training. *Clinical Journal of Sport Medicine* 2006;**16**(4):335-40.

Mika 2011 {published data only}

Mika P, Wilk B, Marchewka A, Nizankowski R. The effect of pain-free treadmill training on fibrinogen, haematocrit, and lipid profile in patients with claudication. *European Journal of Cardiovascular Prevention and Rehabilitation* 2011;**18**(5):754-60.

Sanderson 2006 {published data only}

Sanderson B, Askew C, Stewart I, Walker P, Gibbs H, Green S. Short-term effects of cycle and treadmill training on exercise tolerance in peripheral arterial disease. *Journal of Vascular Surgery* 2006;**44**(1):119-27.

Schlager 2011 {published data only}

* Schlager O, Giurgea A, Schuhfried O, Seidinger D, Hammer A, Gröger M, et al. Exercise training increases endothelial progenitor cells and decreases asymmetric dimethylarginine in peripheral arterial disease: a randomized controlled trial. *Atherosclerosis* 2011;**217**(1):240-8.

Schlager O, Hammer A, Giurgea A, Schuhfried O, Fialka-Moser V, Gschwandtner M, et al. Impact of exercise training on inflammation and platelet activation in patients with intermittent claudication. *Swiss Medical Weekly* 2012;**142**:w13623.

Steiner-Boeker S, Schlager O, Giurgea A, Schuhfried O, Seidinger D, Koppensteiner R. Exercise training on top of standard medication increases endothelial progenitor cells and decreases ADMA levels in patients with PAD. *European Heart Journal* 2009;**30**(984):985.

Tew 2009 {published data only}

Tew G, Nawaz S, Zwierska I, Saxton JM. Limb-specific and cross-transfer effects of arm-crank exercise training in patients with symptomatic peripheral arterial disease. *Clinical Science* 2009;**117**(12):405-13.

Tew 2015 {published data only}

Tew GA, Humphreys L, Crank H, Hewitt C, Nawaz S, Al-Jundi W, et al. The development and pilot randomised controlled trial of a group education programme for promoting walking in people with intermittent claudication. *Vascular Medicine (United Kingdom)*. SAGE Publications Ltd, 2015; Vol. 20, issue 4:348-57.

Tisi 1997 {published data only}

Tisi P, Shearman C. The impact of treatment of intermittent claudication on subjective health of the patient. *Health Trends* 1998/99;**30**:109-14.

* Tisi PV, Hulse M, Chulakadabba A, Gosling P, Shearman CP. Exercise training for intermittent claudication: does it adversely affect biochemical markers of the exercise-induced inflammatory response?. *European Journal of Vascular and Endovascular Surgery* 1997;**14**(5):344-50.

Tsai 2002 {published data only}

Tsai JC, Chan P, Wang CH, Jeng C, Hsieh MH, Kao PF, et al. The effects of exercise training on walking function and perception of health status in elderly patients with peripheral arterial occlusive disease. *Journal of Internal Medicine* 2002;**252**(5):448-55.

Wood 2006 {published data only}

Wood RE, Sanderson BE, Askew CD, Walker PJ, Green S, Stewart IB. Effect of training on the response of plasma vascular endothelial growth factor to exercise in patients with peripheral arterial disease. *Clinical Science* 2006;**111**(6):401-9.

Zwierska 2005 {published data only}

Saxton JM, Zwierska I, Blagojevic M, Choksy SA, Nawaz S, Pockley AG. Upper- versus lower-limb aerobic exercise training on health-related quality of life in patients with symptomatic peripheral arterial disease. *Journal of Vascular Surgery* 2011;**53**(5):1265-73.

Saxton JM, Zwierska I, Hopkinson K, Espigares E, Choksy S, Nawaz S, et al. Effect of upper- and lower-limb exercise training on circulating soluble adhesion molecules, hs-CRP and stress proteins in patients with intermittent claudication. *European Journal of Vascular and Endovascular Surgery* 2008;**35**(5):607-13.

* Zwierska I, Walker RD, Choksy SA, Male JS, Pockley AG, Saxton JM. Upper- vs lower-limb aerobic exercise rehabilitation in patients with symptomatic peripheral arterial disease: a randomized controlled trial. *Journal of Vascular Surgery* 2005;**42**(6):1122-30.

References to studies excluded from this review

Allen 2010 {published data only}

Allen JD, Stabler T, Kenjale A, Ham KL, Robbins JL, Duscha BD, et al. Plasma nitrite flux predicts exercise performance in

peripheral arterial disease after 3 months of exercise training. *Free Radical Biology and Medicine* 2010;**49**(6):1138-44.

Andreozzi 2008 {published data only}

Andreozzi GM, Leone A, Laudani R, Martin R, Deinit G, Cataldi V. Levo-propionyl-carnitine improves the effectiveness of supervised physical training on the absolute claudication distance in patients with intermittent claudication. *Angiology* 2008;**59**(1):84-9.

Aruna 2015 {published data only}

Aruna S, Thenmozhi P. Effectiveness of Allen Buerger exercise in preventing peripheral arterial disease among people with type II diabetes mellitus. *International Journal of Pharma and Bio Sciences* 2015;**6**(2):(B) 966-70.

Beutel 1985 {published data only}

Beutel G, Sobanski R. Results of the complex spa-therapy in peripheral obstructive disease Stage II. *Zeitschrift fur Physiotherapie* 1985;**37**:309-11.

Bronas 2011 {published data only}

* Bronas UG, Treat-Jacobson D, Leon AS. Comparison of the effect of upper body-ergometry aerobic training vs treadmill training on central cardiorespiratory improvement and walking distance in patients with claudication. *Journal of Vascular Surgery* 2011;**53**(6):1557-64.

Treat-Jacobson D, Bronas UG, Leon AS. Efficacy of arm-ergometry versus treadmill exercise training to improve walking distance in patients with claudication. *Vascular Medicine* 2009;**14**(3):203-13.

Brotons 2011 {published data only}

Brotons C, Soriano N, Moral I, Rodrigo MP, Kloppe P, Rodríguez AI, et al. PREseAP Study Research Team. Randomized clinical trial to assess the efficacy of a comprehensive programme of secondary prevention of cardiovascular disease in general practice: the PREseAP study. *Revista Española de Cardiología* 2011;**64**(1):13-20.

Buchwalsky 1974 {published data only}

Buchwalsky R, Blümchen, G, Schlosser V, Pohle W, Isbary J, Barmeyer J, et al. The influence of long-term training on the degree of compensation of peripheral occlusive arterial disease. *Klinische Wochenschrift* 1971; Vol. 49:1045.

* Buchwalsky R, Schnellbacher K, Roskamm H, Barmeyer J. Long-term training effects in chronic ischemia. [German]. *Medizinische Welt* 1974;**25**(3):83-6.

Bulling 1991 {published data only}

Bulling B, Von Bary S. Treatment of chronic peripheral occlusive disease with physical training and ginkgo biloba extract 761. *Medizinische Welt* 1991;**42**(8):702-8.

Cachovan 1999 {published data only}

Cachovan M, Rogatti W, Creutzig A, Diehm C, Heidrich H, Scheffler P, et al. Treadmill testing for evaluation of claudication: comparison of constant-load and graded-exercise tests. *European Journal of Vascular and Endovascular Surgery* 1997;**14**(4):238-43.

* Cachovan M, Rogatti W, Woltering F, Creutzig A, Diehm C, Heidrich H, et al. Randomized reliability study evaluating constant-load and graded-exercise treadmill test for intermittent claudication. *Angiology* 1999;**50**(3):193-200.

Carmeli 2004 {published data only}

Carmeli E, Barchad S, Masharawi Y, Coleman R. Impact of a walking program in people with down syndrome. *Journal of Strength and Conditioning Research* 2004;**18**(1):180-4.

Cheetham 2004 {published data only}

Cheetham DR, Burgess L, Ellis M, Williams A, Greenhalgh RM, Davies AH. Does supervised exercise offer adjuvant benefit over exercise advice alone for the treatment of intermittent claudication? A randomised trial. *European Journal of Vascular and Endovascular Surgery* 2004;**27**(1):17-23.

Choi 2012 {published data only}

Choi KKS, Etnyre G, Fighi SF, Kunkel CF, Ornelas CC, Scremin E. Comparison of calf exercise and treadmill training in peripheral arterial disease. American Academy of Physical Medicine and Rehabilitation Conference 2012; Vol. 4, issue 10 Suppl:S346 (Poster 457).

Cina 1996 {published data only}

Cina G, Vernich M, Campisi C, Cascone C, Ofria F, Leopardi N, et al. Physical training and low-dose heparin-calcium in patients suffering from chronic obliterating arteriopathy of the lower limbs with intermittent claudication [Training fisico ed eparina calcica a basse dosi in pazienti affetti da arteriopatia obliterante cronica degli arti inferiori con claudicatio intermittens]. *Minerva Cardioangiologica* 1996;**44**(4):179-85.

CLEVER 2009 {published data only}

Bronas UG, Hirsch AT, Murphy T, Badenhop D, Collins TC, Ehrman JK, et al. CLEVER Research Group. Design of the multicenter standardized supervised exercise training intervention for the claudication: exercise vs endoluminal revascularization (CLEVER) study. *Vascular Medicine* 2009;**14**(4):313-21.

Melton C. Supervised exercise program helps patients with peripheral artery disease walk longer. *Annals of Long-Term Care* 2011;**19**(12):15-6.

Melton C. Supervised exercise program helps patients with peripheral artery disease walk longer. *Clinical Geriatrics* 2011;**19**(12):17.

Murphy TP. Supervised exercise vs. primary stenting for claudication due to aorto-iliac peripheral artery disease: 6-month outcomes from the CLEVER Study. *European Heart Journal* 2012;**33**(1):144-5.

Murphy TP, Cutlip DE, Regensteiner JG, Mohler ER, Cohen DJ, Reynolds MR, et al. CLEVER Study Investigators. Supervised exercise versus primary stenting for claudication resulting from aortoiliac peripheral artery disease: six-month outcomes from the claudication: exercise versus endoluminal revascularization (CLEVER) study. *Circulation* 2012;**125**(1):130-9.

Murphy TP, Hirsch AT, Cutlip DE, Regensteiner JG, Comerota AJ, Mohler E, et al. CLEVER Investigators. Claudication: exercise vs endoluminal revascularization (CLEVER) study update. *Journal of Vascular Surgery* 2009;**50**(4):942-5, e2.

Murphy TP, Hirsch AT, Ricotta JJ, Cutlip DE, Mohler E, Regensteiner JG, et al. CLEVER Steering Committee. The Claudication: Exercise Vs. Endoluminal Revascularization (CLEVER) study: rationale and methods. *Journal of Vascular Surgery* 2008;**47**(6):1356-63.

Collins 2010 {published data only}

Collins T. Walking therapy for patients with diabetes mellitus and peripheral arterial disease who are limited by leg pain and/or fatigue. *Atherosclerosis Supplements* 2011;**12**(1):118-9.

Collins T, Ghidei W, Ahluwalia J. Benefits of walking therapy in patients with diabetes mellitus and peripheral arterial disease who are limited by leg pain or fatigue. *Journal of General Internal Medicine* 2011;**26**:S176-7.

Collins T, Lunos S. Home-based walking therapy improves walking ability and quality of life in patients with diabetes mellitus and peripheral arterial disease. *Journal of General Internal Medicine* 2010;**25**(3 Suppl):S296-7.

Collins T, Lunos S. Home-based walking therapy improves walking ability and quality of life in persons with diabetes mellitus and peripheral arterial disease. *Vascular Medicine* 2010;**15**:155.

Collins TC, Lunos S, Carlson T, Henderson K, Lightbourne M, Nelson B, et al. Effects of a home-based walking intervention on mobility and quality of life in people with diabetes and peripheral arterial disease: a randomized controlled trial. *Diabetes Care* 2011;**34**(10):2174-9.

Collins TC, Lunos S, Hodges J. Pilot study of whether a walking intervention reduces inflammation in patients with diabetes and peripheral arterial disease (PAD). *Vascular Medicine* 2011;**16**(3):230.

NCT00611988. Self-managed walking improves function. clinicaltrials.gov/ct2/show/NCT00611988. Date first received: 25 January 2008.

Collins 2012 {published data only}

Collins EG, McBurney C, Butler J, Jelinek C, O'Connell S, Fritsch C, et al. The effects of walking or walking-with-poles training on tissue oxygenation in patients with peripheral arterial disease. *International Journal of Vascular Medicine* 2012;**2012**:Article 985025.

Collins EG, O'Connell S, McBurney C, Jelinek C, Butler J, Reda D, et al. Comparison of walking with poles and traditional walking for peripheral arterial disease rehabilitation. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2012;**32**(4):210-8.

Creasy 1990 {published data only}

Creasy TS, Fletcher EW, Walton J, Collin J, Morris P. Prospective randomized trial of PTA versus supervised exercise therapy

for intermittent claudication. *British Journal of Radiology* 1992;**65**(Suppl):108.

* Creasy TS, McMillan PJ, Fletcher EWL, Collin J, Morris PJ. Is percutaneous transluminal angioplasty better than exercise for claudication? Preliminary results from a prospective randomised trial. *European Journal of Vascular Surgery* 1990;**4**(2):135-40.

Perkins JM, Collin J, Creasy TS, Fletcher EW, Morris PJ. Exercise training versus angioplasty for stable claudication. Long and medium term results of a prospective, randomised trial. *European Journal of Endovascular Surgery* 1996;**11**(4):409-13.

Crowther 2008 {published data only}

* Crowther RG, Spinks WL, Leicht AS, Sangla K, Quigley F, Golledge J. Effects of a long-term exercise program on lower limb mobility, physiological responses, walking performance, and physical activity levels in patients with peripheral arterial disease. *Journal of Vascular Surgery* 2008;**47**(2):303-9.

Crowther RG, Spinks WL, Leicht AS, Sangla K, Quigley F, Golledge J. The influence of a long term exercise program on lower limb movement variability and walking performance in patients with peripheral arterial disease. *Human Movement Science* 2009;**28**(4):494-503.

Cucato 2011 {published data only}

Cucato GG, Ritti-Dias RM, Wolosker N, Santarem JM, Jacob Filho W, Forjaz CL, et al. Post-resistance exercise hypotension in patients with intermittent claudication. *Clinics (Sao Paulo, Brazil)* 2011;**66**(2):221-6.

Cucato 2011a {published data only}

Cucato GG, Forjaz CL, Kanegusuku H, da Rocha Chehuen M, Riani Costa LA, Wolosker N, et al. Effects of walking and strength training on resting and exercise cardiovascular responses in patients with intermittent claudication. *Vasa* 2011;**40**(5):390-7.

Cucato 2015 {published data only}

Cucato GG, Saxton JM. Post-walking exercise hypotension in patients with intermittent claudication. *Medicine and Science in Sports and Exercise* 2015;**47**(3):460-7.

Cunningham 2012 {published data only}

Cunningham MA, Swanson V, Holdsworth RJ, O'Carroll RE. Late effects of a brief psychological intervention in patients with intermittent claudication in a randomized clinical trial. *British Journal of Surgery* 2013;**100**(6):756-60.

* Cunningham MA, Swanson V, O'Carroll RE, Holdsworth RJ. Randomized clinical trial of a brief psychological intervention to increase walking in patients with intermittent claudication. *British Journal of Surgery* 2012;**99**(1):49-56.

Dahllof 1976 {published data only}

Dahllof AG, Holm J, Schersten T, Sivertsson R. Peripheral arterial insufficiency, effect of physical training on walking tolerance, calf blood flow and blood flow resistance. *Scandinavian Journal of Rehabilitation Medicine* 1976;**8**(1):19-26.

Dantas 2016 {published data only}

Dantas FF, da Silva Santana F, da Silva TS, Cucato GG, Farah BQ, Ritti-Dias RM. Acute effects of T'ai Chi Chuan exercise on blood pressure and heart rate in peripheral artery disease patients. *Journal of Alternative and Complementary Medicine (New York, N.Y.)* 2016;**22**:375-9.

Dedes 2010 {published data only}

Dedes H, Figoni SF, Kalioundji G, Kunkel C, Peter A, Phillips A, et al. Prospective trial of calf ergometry training on walking ability in peripheral arterial disease. *Physical Medicine and Rehabilitation* 2010;**2**(9 Suppl 1):S26.

Degischer 2002 {published data only}

Degischer S, Labs KH, Hochstrasser J, Aschwanden M, Tschoepf M, Jaeger KA. Physical training for intermittent claudication: a comparison of structured rehabilitation versus home-based training. *Vascular Medicine* 2002;**7**(2):109-15.

Dittmar 1977 {published data only}

Dittmar K, Krause D. Success of physiotherapeutic interval training combined with a standardized hemoderivative in intermittent claudication. *Munchener Medizinische Wochenschrift* 1977;**119**(11):369-72.

Ericsson 1970 {published data only}

Ericsson B, Haeger K, Lindell SE. Effect of physical training of intermittent claudication. *Angiology* 1970;**21**(3):188-92.

Ernst 1987 {published data only}

Ernst EE, Matrai A. Intermittent claudication, exercise, and blood rheology. *Circulation* 1987;**76**(5):1110-4.

Ernst 1990 {published data only}

Ernst E, Kollar L, Resch KL, Bergmann H. Arterial occlusive disease. Comparison between pentoxifylline and exercise vs. exercise alone in patients with stage II of disease. *Munchener Medizinische Wochenschrift* 1990;**132**(28-29):456-8.

EXERT 2009 {published data only}

NCT00895635. Evaluating two exercise training programs to reduce leg pain in people with peripheral arterial disease (The EXERT study). clinicaltrials.gov/ct2/show/NCT00895635 Date first received: 6 May 2009.

EXITPAD 2010 {published data only}

Gommans LNM, Scheltinga MRM, Van Sambeek MRHM, Maas AHEM, Bendermacher BLW, Teijink JAW. Gender differences following supervised exercise therapy in patients with intermittent claudication. *Journal of Vascular Surgery* 2015;**62**:681-8.

Nicolai SP, Hendriks EJ, Prins MH, Teijink JA, EXITPAD Study Group. Optimizing supervised exercise therapy for patients with intermittent claudication. *Journal of Vascular Surgery* 2010;**52**(5):1226-33.

* Nicolai SP, Teijink JA, Prins MH, Exercise Therapy in Peripheral Arterial Disease Study Group. Multicenter randomized clinical trial of supervised exercise therapy with or without feedback

versus walking advice for intermittent claudication. *Journal of Vascular Surgery* 2010;**52**(2):348-55.

Nicolai SPA, Prins MH, Teijink JAW. Supervised exercise therapy is more effective than walking advice in patients with intermittent claudication; a randomized multicenter study. [Dutch]. *Nederlands Tijdschrift voor Geneeskunde* 2011;**155**(2):55-62.

van Asselt AD, Nicolai SP, Joore MA, Prins MH, Teijink JA, Exercise Therapy in Peripheral Arterial Disease Study Group. Cost-effectiveness of exercise therapy in patients with intermittent claudication: supervised exercise therapy versus a 'go home and walk' advice. *European Journal of Vascular and Endovascular Surgery* 2011;**41**(1):97-103.

Fakhry 2011 {published data only}

Fakhry F, Spronk S, de Ridder M, den-Hoed PT, Hunink MG. Long-term effects of structured home-based exercise program on functional capacity and quality of life in patients with intermittent claudication. *Archives of Physical Medicine and Rehabilitation* 2011;**92**(7):1066-73.

Fitzgerald 1971 {published data only}

Fitzgerald DE, Keates JS, MacMillan D. Angiographic and plethysmographic assessment of graduated physical exercise in the treatment of chronic occlusive arterial disease of the leg. *Angiology* 1971;**22**(2):99-106.

Fowler 2002 {published data only}

Fowler B, Jamrozik K, Norman P, Allen Y, Wilkinson E. Improving maximum walking distance in early peripheral arterial disease: randomised controlled trial. *Australian Journal of Physiotherapy* 2002;**48**(4):269-75.

Gardner 2005 {published data only}

Gardner AW, Montgomery PS, Flinn WR, Katzel LI. The effect of exercise intensity on the response to exercise rehabilitation in patients with intermittent claudication. *Journal of Vascular Surgery* 2005;**42**(4):702-9.

Gardner 2011 {published data only}

Gardner AW, Parker DE, Montgomery PS, Scott KJ, Blevins SM. Efficacy of quantified home-based exercise and supervised exercise in patients with intermittent claudication: a randomized controlled trial. *Circulation* 2011;**123**(5):491-8.

Gardner 2012 {published data only}

* Gardner AW, Montgomery PS, Parker DE. Optimal exercise program length for patients with claudication. *Journal of Vascular Surgery* 2012;**55**(5):1346-54.

Gardner AW, Montgomery PS, Parker DE. Optimal exercise program length for patients with claudication: a randomized controlled trial. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2011; Vol. 31:E4-5.

Gardner 2014 {published data only}

Gardner AW, Parker DE, Montgomery PS, Blevins SM. Diabetic women are poor responders to exercise rehabilitation in the treatment of claudication. *Journal of Vascular Surgery* 2014;**59**(4):1036-43.

Gardner 2014a {published data only}

Gardner AW, Parker DE, Montgomery PS, Blevins SM. Step-monitored home exercise improves ambulation, vascular function, and inflammation in symptomatic patients with peripheral artery disease: a randomized controlled trial. *Journal of the American Heart Association* 2014;**3**(5):e001107.

Gibbellini 2000 {published data only}

Gibellini R, Fanello M, Bardile AF, Salerno M, Aloï T. Exercise training in intermittent claudication. *International Angiology* 2000;**19**(1):8-13.

Gibbs 2013 {published data only}

Gibbs BB, Dobrosielski DA, Althouse AD, Stewart KJ. The effect of exercise training on ankle-brachial index in type 2 diabetes. *Atherosclerosis* 2013;**230**(1):125-30.

Gottstein 1987 {published data only}

Gottstein P, Schollmayer E. Additive effect of Buflomedil under simultaneous training. Buflomedil-therapy in combination with intensive training in patients with intermittent claudication. *Zeitschrift für Allgemeinmedizin* 1987;**63**(28):836-9.

Greenhalgh 2008 {published data only}

Greenhalgh RM. MIMIC Trials: Angioplasty effective in randomised controlled trials for peripheral arterial disease. <http://www.cxvasc.com/interventionalnews/latestnews.cfm?ccs=485&cs=4222>. accessed 2 February 2009.

* Greenhalgh RM, Belch JJ, Brown LC, Gaines PA, Gao L, Reise JA, et al. The adjuvant benefit of angioplasty in patients with mild to moderate intermittent claudication (MIMIC) managed by supervised exercise, smoking cessation advice and best medical therapy: results from two randomised trials for stenotic femoropopliteal and aortoiliac arterial disease. *European Journal of Vascular and Endovascular Surgery* 2008;**36**(6):680-8.

Guidon 2013 {published data only}

Guidon M, McGee H. Recruitment to clinical trials of exercise: challenges in the peripheral arterial disease population. *Physiotherapy* 2013;**99**(4):305-10.

Guirro 2015 {published data only}

Guirro EC, Guirro RR, Dibai-Filho AV, Pascote SCS, Rodrigues-Bigaton D. Immediate effects of electrical stimulation, diathermy, and physical exercise on lower limb arterial blood flow in diabetic women with peripheral arterial disease: a randomized crossover trial. *Journal of Manipulative and Physiological Therapeutics* 2015;**38**(3):195-202.

Hobbs 2006 {published data only}

Hobbs SD, Bradbury AW. The EXercise versus Angioplasty in Claudication Trial (EXACT): reasons for recruitment failure and the implications for research into and treatment of intermittent claudication. *Journal of Vascular Surgery* 2006;**44**(2):432-3.

* Hobbs SD, Marshall T, Fegan C, Adam DJ, Bradbury AW. The constitutive procoagulant and hypofibrinolytic state in patients with intermittent claudication due to infrainguinal disease

significantly improves with percutaneous transluminal balloon angioplasty. *Journal of Vascular Surgery* 2006;**43**(1):40-6.

Hodges 2008 {published data only}

Hodges LD, Sandercock GR, Das SK, Brodie DA. Randomized controlled trial of supervised exercise to evaluate changes in cardiac function in patients with peripheral atherosclerotic disease. *Clinical Physiology and Functional Imaging* 2008;**28**(1):32-7.

Holm 1973 {published data only}

Holm J, Dahllöf AG, Björntorp P, Schersten T. Enzyme studies in muscles of patients with intermittent claudication. Effect of training. *Scandinavian Journal of Clinical Laboratory Investigation* 1973;**31**(Suppl 128):201-5.

Jones 1996 {published data only}

Jones PP, Skinner JS, Smith LK, John FM, Bryant CX. Functional improvements following StairMaster vs. treadmill exercise training for patients with intermittent claudication. *Journal of Cardiopulmonary Rehabilitation* 1996;**16**(1):47-55.

Kiesewetter 1987 {published data only}

Kiesewetter H, Blume J, Jung F, Gerhards M, Leipnitz G. Training by walking and drug therapy of peripheral arterial occlusive disease. [German]. *Vasa Supplementum* 1987;**20**:384-7.

Kono 2013 {published data only}

Kono Y, Yamada S, Yamaguchi J, Hagiwara Y, Iritani N, Ishida S, et al. Secondary prevention of new vascular events with lifestyle intervention in patients with noncardioembolic mild ischemic stroke: a single-center randomized controlled trial. *Cerebrovascular Diseases* 2013;**36**(2):88-97.

Krause 1976 {published data only}

Krause D, Dittmar K. Combination of physiotherapeutic exercise therapy with bencyclane in intermittent claudication. *Munchener Medizinische Wochenschrift* 1976;**118**(40):1281-4.

Kruidenier 2011 {published data only}

Kruidenier LM, Nicola SP, Rouwet EV, Peters RJ, Prins MH, Teijink JAW. Additional supervised exercise therapy after a percutaneous vascular intervention for peripheral arterial disease: a randomized clinical trial. *Journal of Vascular and Interventional Radiology* 2011;**22**(7):961-8.

Labs 1999 {published data only}

Labs KH, Nehler MR, Roessner M, Jaeger KA, Hiatt WR. Reliability of treadmill testing in peripheral arterial disease: a comparison of a constant load with a graded load treadmill protocol. *Vascular Medicine* 1999;**4**(4):239-46.

Lee 2007 {published data only}

Lee HL, Mehta T, Ray B, Heng MS, McCollum PT, Chetter IC. A non-randomised controlled trial of the clinical and cost effectiveness of a Supervised Exercise Programme for claudication. *European Journal of Endovascular and Vascular Surgery* 2007;**33**(2):202-7.

Leon 2005 {published data only}

Leon AS. Does long-term aerobic exercise slow progression of atherosclerosis?. *Clinical Journal of Sport Medicine* 2005;**15**(4):285-6.

Lepantalo 1984 {published data only}

Lepantalo M, Sundberg S, Gordin A. The effects of physical training and flunarizine on walking capacity in intermittent claudication. *Scandinavian Journal of Rehabilitation Medicine* 1984;**16**(4):159-62.

LIFE Study {published data only}

McDermott M, Bonds D, Buford T, Church T, Dodson J, Fielding J, et al. A physical activity intervention improves 400-meter walking velocity as compared to a successful aging intervention in patients with peripheral artery disease: the LIFE study randomized trial. *Circulation* 2014; Vol. 130:A19294.

Newman ABD, Kritchevsky S, Beavers MM. Cardiovascular events in a physical activity intervention as compared to a successful aging intervention: the LIFE study randomized trial. *Circulation* 2014; Vol. 130:A19294.

Lundgren 1989 {published data only}

* Lundgren F, Dahllof AG, Lundholm K, Schersten T, Volkmann R. Intermittent claudication - surgical reconstruction or physical training? A prospective randomised trial of treatment efficiency. *Annals of Surgery* 1989;**209**(3):346-55.

Lundgren F, Dahllof AG, Schersten T, Bylund-Fellenius AC. Muscle enzyme adaptation in patients with peripheral arterial insufficiency: spontaneous adaptation, effect of different treatments and consequences on walking performance. *Clinical Science* 1989;**77**(5):485-93.

Maejima 2005 {published data only}

Maejima Y, Yasu T, Ueba H, Kobayashi N, Hashimoto S, Kubo N, et al. Exercise after heparin administration: new therapeutic program for patients with-option arteriosclerosis obliterans. *Circulation Journal* 2005;**69**(9):1099-104.

Mannarino 1988 {published data only}

Mannarino E, Menna M, Pasqualini L, Innocente S, Regni O. Effect of physical exercise in patients with peripheral occlusive arteriopathy. Controlled study. *Riabilitazione* 1988;**21**(2):123-31.

Mannarino 1989 {published data only}

Mannarino E, Pasqualini L, Menna M, Maragoni G, Orlandi U. Effects of physical training on peripheral vascular disease: a controlled study. *Angiology* 1989;**40**(1):5-10.

Martinez 2009 {published data only}

Martinez CA, Carmeli E, Barak S, Stopka CB. Changes in pain-free walking based on time in accommodating pain-free exercise therapy for peripheral arterial disease. *Journal of Vascular Nursing* 2009;**27**(1):2-7.

Mays 2015 {published data only}

Mays RJ. Community-based walking exercise for patients with peripheral artery disease: a pilot study moderated. *Journal of*

the American College of Cardiology 2014; Vol. 63, issue 12(Suppl 1):A2035.

Mays RJ, Hiatt WR, Casserly IP, Rogers RK, Main DS, Kohrt WM, et al. Community-based walking exercise for peripheral artery disease: an exploratory pilot study. *Vascular Medicine* 2015;**20**(4):339-47.

Mazari 2010 {published data only}

Lee HLD, Gulati S, Mehta T, Mekako AI, Rahman MNA, McCollum P, et al. Early result of a randomised controlled trial of treatment for intermittent claudication. *The Vascular Society of Great Britain & Ireland Yearbook*. 2007:78.

Mazari F, Khan J, Abdul Rahman MNA, Mehta T, Gulati S, McCollum P, et al. Cost utility analysis of a randomised control trial of percutaneous transluminal angioplasty (PTA), supervised exercise programme (SEP) and combined treatment (PTA+SEP) for patients with intermittent claudication (IC) due to femoropopliteal disease. *The Vascular Society of Great Britain and Ireland Yearbook*. 2009:44.

* Mazari FA, Gulati S, Rahman MN, Lee HL, Mehta TA, McCollum PT, et al. Early outcomes from a randomized, controlled trial of supervised exercise, angioplasty, and combined therapy in intermittent claudication. *Annals of Vascular Surgery* 2010;**24**(1):69-79.

Mazari FA, Khan JA, Carradice D, Samuel N, Abdul Rahman MN, Gulati S, et al. Randomized clinical trial of percutaneous transluminal angioplasty, supervised exercise and combined treatment for intermittent claudication due to femoropopliteal arterial disease. *British Journal of Surgery* 2012;**99**(1):39-48.

Mazari FAK, Mehta T, Rahman MNA, McCollum P, Chetter IC. A RCT of non-surgical treatment for intermittent claudication in femoro-popliteal disease: 12-month results. *The Vascular Society of Great Britain & Ireland Yearbook*. 2008:75.

McDermott 2004 {published data only}

McDermott MM, Tiukinhoy S, Greenland P, Liu K, Pearce WH, Guralnik JM, et al. A pilot exercise intervention to improve lower extremity functioning in peripheral arterial disease unaccompanied by intermittent claudication. *Journal of Cardiopulmonary Rehabilitation* 2004;**24**(3):187-96.

Nawaz 1999 {published data only}

Nawaz S, Walker RD, Pockley AG, Wood RF. A safer way to exercise claudicants?. *British Journal of Surgery* 1999;**86**(Suppl 1):10-1.

Nawaz 2001 {published data only}

Nawaz S, Walker RD, Wilkinson CH, Saxton JM, Pockley AG, Wood RF. The inflammatory response to upper and lower limb exercise and the effects of exercise training in patients with claudication. *Journal of Vascular Surgery* 2001;**33**(2):392-9.

NCT01065740 {published data only}

NCT01065740. Optimized supervised education program for peripheral arterial disease (POPART). clinicaltrials.gov/ct2/show/NCT01065740. Date first received: 8 February 2010.

NCT01241747 {published data only}

NCT01241747. Exercise for women with peripheral arterial disease. clinicaltrials.gov/show/NCT01241747. Date first received: 15 November 2010.

NCT02075502 {published data only}

NCT02075502. Community walking exercise for patients with peripheral artery disease (GAIT). clinicaltrials.gov/ct2/show/NCT02075502. Date first received: 13 February 2014.

NCT02879019 {published data only}

NCT02879019. Walking training in peripheral artery disease (GrEnADa Sub-study) (GrEnADa). clinicaltrials.gov/ct2/show/NCT02879019. Date first received: 11 August 2016.

Necker 2003 {published data only}

Necker A, Schumm T, Liu Y, Baur C, Steinacker JM, Lehmann M. Effects of training after angioplasty on performance and circulation of the lower extremities in peripheral occlusive arterial disease. *Deutsche Zeitschrift für Sportmedizin* 2001;**52**:s67.

* Necker A, Stilgenbauer F, Schumm T, Liu Y, Steinacker JM. Effects of training after angioplasty on walking distance, blood flow and muscular HSP70 expression in peripheral occlusive arterial disease. *Deutsche Zeitschrift für Sportmedizin* 2003;**54**:S46.

Nicolai 2010 {published data only}

Nicolai SP, Leffers P, Kruidenier LM, De-Bie RA, Prins MH, Teijink JA. Extending the range of treadmill testing for patients with intermittent claudication. *Medicine and Science in Sports and Exercise* 2010;**42**(4):640-5.

Nielsen 1977 {published data only}

Nielsen SL, Gyntelberg F, Larsen B, Lassen NA. Hospital versus home training, a clinical trial. *Aktuelle Probleme in der Angiologie* 1975;**30**:121-6.

* Nielsen SL, Larsen B, Prah M, Jensen CT, Jensen BE, Wenkens V. Hospital training compared with home training in patients with intermittent claudication. [Danish]. *Ugeskrift for Læger* 1977;**139**(46):2733-6.

Nordanstig 2011 {published data only}

NCT01219842. Invasive revascularization or not in intermittent claudication - a randomised controlled trial IRONIC. clinicaltrials.gov/ct2/show/NCT01219842. Date first received: 12 October 2010.

* Nordanstig J, Gelin J, Hensater M, Taft C, Osterberg K, Jivegard L. Walking performance and health-related quality of life after surgical or endovascular invasive versus non-invasive treatment for intermittent claudication - a prospective randomised trial. *European Journal of Vascular and Endovascular Surgery* 2011;**42**(2):220-7.

Parr 2009 {published data only}

Parr BM, Noakes TD, Derman EW. Peripheral arterial disease and intermittent claudication: efficacy of short-term upper body strength training, dynamic exercise training, and

advice to exercise at home. *South African Medical Journal* 2009;**99**(11):800-4.

Patterson 1997 {published data only}

Patterson RB, Pinto B, Marcus B, Colucci A, Braun T, Roberts M. Value of a supervised exercise program for the therapy of arterial claudication. *Journal of Vascular Surgery* 1997;**25**(2):312-9.

Pinto 1997 {published data only}

Pinto BM, Marcus BH, Patterson RB, Roberts M, Colucci A, Braun C. On-site versus home exercise programs: psychological benefits for individuals with arterial claudication. *Journal of Aging and Physical Activity* 1997;**5**(4):311.

Presern-Strukelj 200 {published data only}

Presern-Strukelj M, Poredos P. The influence of electrostimulation on the circulation of the remaining leg in patients with one-sided amputation. *Angiology* 2002;**53**(3):329-35.

PROPEL study {published data only}

Domanchuk K, Ferrucci L, Guralnik JM, Criqui MH, Tian L, Liu K, et al. Progenitor cell release plus exercise to improve functional performance in peripheral artery disease: the PROPEL study. *Contemporary Clinical Trials* 2013;**36**(2):502-9.

Saber R, Liu K, Ferrucci L, Criqui MH, Zhao L, Tian L, et al. Ischemia-related changes in circulating stem and progenitor cells and associated clinical characteristics in peripheral artery disease. *Vascular Medicine* 2015;**20**(6):534-43. [DOI: [10.1177/1358863X15600255](https://doi.org/10.1177/1358863X15600255)]

Riccioni 2010 {published data only}

Riccioni C, Sarcinella R, Izzo A, Palermo G, Liguori M, Caliumi C, et al. Rehabilitative treatment in peripheral artery disease: protocol application and follow-up. *Minerva Cardioangiologica* 2010;**58**(5):551-65.

Richardson 1991 {published data only}

Richardson JK. Rocker-soled shoes and walking distance in patients with calf claudication. *Archives of Physical Medicine and Rehabilitation* 1991;**72**(8):554-8.

Riebe 2001 {published data only}

Riebe D, Patterson RB, Braun CM. Comparison of two progressive treadmill tests in patients with peripheral arterial disease. *Vascular Medicine* 2001;**6**(4):215-21.

Ritti-Dias 2010 {published data only}

Meneses AL, de Lima GH, Forjaz CL, Lima AH, Silva GQ, Cucato GG, et al. Impact of a supervised strength training or walking training over a subsequent unsupervised therapy period on walking capacity in patients with claudication. *Journal of Vascular Nursing* 2011;**29**(2):81-6.

Ritti-Dias RM, Wolosker N, de Moraes Forjaz CL, Carvalho CR, Cucato GG, Leão PP, et al. Strength training increases walking tolerance in intermittent claudication patients: randomized trial. *Journal of Vascular Surgery* 2010;**51**(1):89-95.

Rodrigues 2014 {published data only}

Rodrigues L, Forjaz C. A single bout of resistance exercise does not modify cardiovascular responses during daily activities in patients with peripheral artery disease. *Blood Pressure Monitoring* 2014;**19**(2):64-71.

Saleem 2011 {published data only}

Saleem H, Delaney C, Puckridge P, Foreman RT, Spark JI. What is the most appropriate exercise programme for patients with intermittent claudication? A study based on clinical response and inflammatory markers [Abstract]. *ANZ Journal of Surgery* 2011;**81**:A95.

Savage 2001 {published data only}

Savage P, Ricci MA, Lynn M, Gardner A, Knight S, Brochu M, et al. Effects of home versus supervised exercise for patients with intermittent claudication. *Journal of Cardiopulmonary Rehabilitation* 2001;**21**(3):152-7.

Scheffler 1991 {published data only}

Scheffler P, de la Hamette D, Muller H. Controlled vascular training in IIb peripheral arterial occlusive disease: additive effect of intravenous PGE1 versus intravenous pentoxifylline during training [German]. *Vasa Supplementum* 1991;**33**:350-2.

Schlager 2011a {published data only}

Schlager O, Giurgea A, Schuhfried O, Seidinger D, Gschwandner ME, Fialka-Moser V, et al. Supervised exercise training has no effect on inflammation and platelet activation in peripheral arterial disease. *Circulation* 2011;**124**:A11379.

Schoneberger 1994 {published data only}

Schoneberger A, Horeis M, Raschka C. The effects of kinetotherapy and electrotherapy on the maximum walking distance of patients with peripheral arterial occlusion at stage II according to Fontaine. *International Journal of Sports Medicine*. 15 1994; Vol. 15:356.

Silvestro 2002 {published data only}

Silvestro A, Scopacasa F, Oliva G, de Cristofaro T, Iuliano L, Brevetti G. Vitamin C prevents endothelial dysfunction induced by acute exercise in patients with intermittent claudication. *Atherosclerosis* 2002;**165**(2):277-83.

Slordahl 2005 {published data only}

Slordahl SA, Wang E, Hoff J, Kemi OJ, Amundsen BH, Helgerud J. Effective training for patients with intermittent claudication. *Scandinavian Cardiovascular Journal* 2005;**39**(4):244-9.

Snabl 1958 {published data only}

Snabl P, Fiala Z, Polak F. The treatment of lower extremity arterial disease. *Vnitřní Lekarství* 1958;**4**(3):253-8.

Sonaglia 2013 {published data only}

Sonaglia F, Milia P, Caserio M, Bigazzi B, Bigazzi B, Ricotta S, et al. Efficacy and safety of oral porcine relaxin (pRLX) in adjunct to physical exercise in the treatment of peripheral arterial disease (PAD). *Italian Journal of Anatomy and Embryology* 2013;**118**(1 Suppl):84-91.

Spronk 2009 {published data only}

ISRCTN64443682. Comparing exercise training and angioplasty for claudication: a randomised controlled trial. <http://www.controlled-trials.com/ISRCTN64443682> (accessed December 2013).

Spronk S, Bosch JL, den Hoed PT, Veen HF, Pattynama PM, Hunink MG. Cost-effectiveness of endovascular revascularization compared to supervised hospital-based exercise training in patients with intermittent claudication: a randomized controlled trial. *Journal of Vascular Surgery* 2008;**48**(6):1472-80.

* Spronk S, Bosch JL, den Hoed PT, Veen HF, Pattynama PM, Hunink MG. Intermittent claudication: clinical effectiveness of endovascular revascularization versus supervised hospital-based exercise training - randomized controlled trial. *Radiology* 2009;**250**(2):586-95.

Stewart 2008 {published data only}

Stewart AH, Smith FC, Baird RN, Lamont PM. Local versus systemic mechanisms underlying supervised exercise training for intermittent claudication. *Vascular and Endovascular Surgery* 2008;**42**(4):314-20.

Streminski 1992 {published data only}

Streminski JA, de la Haye R, Rettig K, Kuntz G. Comparison of the effectiveness of physical training with parental drug therapy in Fontaine stage IIb peripheral arterial occlusive disease. *VASA* 1992;**21**(4):392-402.

SUPER study {published data only}

* Frans FA, Bipat S, Reekers JA, Legemate DA, Koelemay MJ, SUPER Study Collaborators. SUPERvised exercise therapy or immediate PTA for intermittent claudication in patients with an iliac artery obstruction - a multicentre randomised controlled trial; SUPER study design and rationale. *European Journal of Vascular and Endovascular Surgery* 2012;**43**(4):466-71.

NTR2776. SUPERvised exercise therapy or immediate PTA for intermittent claudication in patients with an iliac artery obstruction: a randomized controlled trial. SUPER study. <http://apps.who.int/trialsearch/Trial.aspx?TrialID=NTR2776> (accessed December 2013).

Taft 2004 {published data only}

Taft C, Sullivan M, Lundholm K, Karlsson J, Gelin J, Jivegard L. Predictors of treatment outcome in intermittent claudication. *European Journal of Vascular and Endovascular Surgery* 2004;**27**(1):24-32.

Tebbutt 2011 {published data only}

Tebbutt N, Robinson L, Todhunter J, Jonker L. A plantar flexion device exercise programme for patients with peripheral arterial disease: a randomised prospective feasibility study. *Physiotherapy* 2011; Vol. 97, issue 3:244-9.

Thomson 1999 {published data only}

Thomson IA, van Rij AM, Morrison ND, Packer SGK, Christie R. A ten year randomised controlled trial of percutaneous femoropopliteal angioplasty for claudication. *Australia and New Zealand Journal of Surgery* 1999;**69**(Suppl):98.

Treat-Jacobson 2012 {published data only}

Treat-Jacobson DJ, Bronas UG, Krause BJ, Robinson CA, Santili SM, Leon AS. Aerobic arm exercise training to improve outcomes for patients with severe claudication and ischemic rest pain. *Vascular Medicine* 2012;**17**(3):204.

Ventura 1984 {published data only}

Ventura MR, Young DE, Feldman MJ, Pastore P, Pikula S, Yates MA. Effectiveness of health promotion interventions. *Nursing Research* 1984;**33**(3):162-7.

Walker 2000 {published data only}

Walker RD, Nawaz S, Wilkinson CH, Saxton JM, Pockley AG, Wood RF. Influence of upper- and lower-limb exercise training on cardiovascular function and walking distances in patients with intermittent claudication. *Journal of Vascular Surgery* 2000;**31**(4):662-9.

Waller 1988 {published data only}

Waller PC, Solomon SA, Ramsay LE. Cigarette smoking and treadmill exercise distances in patients with intermittent claudication: the effects of suggestion. Proceedings of The British Pharmacological Society. 1988:632P-3P.

Wang 2008 {published data only}

Wang E, Hoff J, Loe H, Kaehler N, Helgerud J. Plantar flexion: an effective training for peripheral arterial disease. *European Journal of Applied Physiology* 2008;**104**(4):749-56.

Wang 2010 {published data only}

Wang E, Helgerud J, Loe H, Indseth K, Kaehler N, Hoff J. Maximal strength training improves walking performance in peripheral arterial disease patients. *Scandinavian Journal of Medicine and Science in Sports* 2010;**20**(5):764-70.

Winterfeld 1983 {published data only}

Winterfeld HJ, Siewert H, Strangfeld D, Schmidt HH. Experiences with the application of physiotherapy in peripheral arterial circulatory disorders of the lower extremities (stage I and IIa) under ambulatory conditions. *Zeitschrift für die gesamte innere Medizin und ihre Grenzgebiete* 1983;**38**(7):221-5.

Zwierska 2004 {published data only}

Zwierska I, Nawaz S, Walker RD, Wood RF, Pockley AG, Saxton JM. Treadmill versus shuttle walk tests of walking ability in intermittent claudication. *Medicine and Science in Sports and Exercise* 2004;**36**(11):1835-40.

References to ongoing studies
NCT01231360 {published data only}

NCT01231360. The effect of exercise training on skeletal muscle metabolism in peripheral artery disease (PAD). clinicaltrials.gov/ct2/show/NCT01231360. Date first received: 28 October 2010.

NCT01822457 {published data only}

NCT01822457. Effect of Nike Fuel Band on exercise and function in claudicants: a randomised controlled trial. clinicaltrials.gov/ct2/show/NCT01822457. Date first received: 19 February 2013.

Additional references
Antoniou 2017

Antoniou GA, Georgiadis GS, Antoniou SA, Makar RR, Smout JD, Torella F. Bypass surgery for chronic lower limb ischaemia. *Cochrane Database of Systematic Reviews* 2017, Issue 4. [DOI: [10.1002/14651858.CD002000.pub3](https://doi.org/10.1002/14651858.CD002000.pub3)]

Baumgartner 1998

Baumgartner TA, Strong CH. Conducting and Reading Research. 2nd Edition. New York: WCB/Mcgraw-Hill, 1998.

Boutroux 1980

Boutroux YC, Baccard D, Bouchet JY. Programmed physical training in physiotherapy for obliterative arteriopathies of the lower limbs at the stage of claudication (authors' transl). *Journal des Maladies Vasculaires* 1980;**5**(3):173-6.

Clyne 1985

Clyne CAC, Mears H, Weller RO, O'Donnell TF. Calf muscle adaptation to peripheral vascular disease. *Cardiovascular Research* 1985;**19**(8):507-12.

Collins 2003

Collins EG, Langbein WE, Orebaugh C, Bammert C, Hanson K, Reda D, et al. Polestriding exercise and vitamin E for management of peripheral vascular disease. *Medicine and Science in Sports and Exercise* 2003;**35**(3):384-93.

Cucato 2013b

Cucato GG, Zerati AE, da Rocha Chehuen M, Ritti-Dias RM, Saez G, Ragazzo L, et al. Comparison between subjective and objective methods to assess functional capacity during clinical treatment in patients with intermittent claudication [Comparação entre os métodos subjetivo e objetivo para avaliação da capacidade funcional durante tratamento clínico em pacientes com claudicação intermitente]. *Einstein (Sao Paulo)* 2013;**11**(4):495-9.

Cunningham 2014

Cunningham MA, Swanson V, Pappas E, O'Carroll RE, Holdsworth RJ. Illness beliefs and walking behavior after revascularization for intermittent claudication: a qualitative study. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2014;**34**(3):195-201.

Câmara 2012

Câmara LC, Ritti-Dias RM, Meneses AL, D'Andréa Greve JM, Filho WJ, Santarém JM, et al. Isokinetic strength and endurance in proximal and distal muscles in patients with peripheral artery disease. *Annals of Vascular Surgery* 2012;**26**(8):1114-9.

Delaney 2014

Delaney CL, Miller MD, Chataway TK, Spark JI. A randomised controlled trial of supervised exercise regimens and their impact on walking performance, skeletal muscle mass and calpain activity in patients with intermittent claudication. *European Journal of Vascular and Endovascular Surgery* 2014;**47**(3):304-10.

Ernst 1987a

Ernst E. Physical exercise for peripheral vascular disease: a review. *VASA* 1987;**16**(3):227-31.

Ernst 1992

Ernst E. Exercise: the best therapy for intermittent claudication?. *British Journal of Hospital Medicine* 1992;**48**(6):303-4, 307.

Fakhry 2013

Fakhry F, Fokkenrood HJP, Rouwet EV, Teijink JAW, Spronk S, Hunink MGM. Endovascular revascularisation versus conservative management for intermittent claudication. *Cochrane Database of Systematic Reviews* 2013, Issue 5. [DOI: [10.1002/14651858.CD010512](https://doi.org/10.1002/14651858.CD010512)]

Fokkenrood 2013

Fokkenrood HJP, Bendermacher BLW, Lauret GJ, Willigendael EM, Prins MH, Teijink JAW. Supervised exercise therapy versus non-supervised exercise therapy for intermittent claudication. *Cochrane Database of Systematic Reviews* 2013, Issue 8. [DOI: [10.1002/14651858.CD005263.pub3](https://doi.org/10.1002/14651858.CD005263.pub3)]

Fowkes 1998

Fowkes G, Gillespie IN. Angioplasty (versus non surgical management) for intermittent claudication. *Cochrane Database of Systematic Reviews* 1998, Issue 2. [DOI: [10.1002/14651858.CD000017](https://doi.org/10.1002/14651858.CD000017)]

Gardner 1995

Gardner AW, Poehlman ET. Exercise rehabilitation programs for the treatment of claudication pain. A meta analysis. *JAMA* 1995;**274**(12):975-80.

Gohil 2013

Gohil RA, Mockford KA, Mazari FA, Khan JA, Van Vanicek N, Chetter IC, et al. Percutaneous transluminal angioplasty results in improved physical function but not balance in patients with intermittent claudication. *Journal of Vascular Surgery* 2013;**58**(6):1533-9.

Gohil 2013a

Gohil RA, Lane TRA, Coughlin PA. Review of the adaptation of skeletal muscle in intermittent claudication. *World Journal of Cardiovascular Diseases* 2013;**3**:347-60.

Gommans 2016

Gommans LN, Smid AT, Scheltinga MR, Brooijmans FA, van Disseldorp EM, van der Linden FT, et al. Altered joint kinematics and increased electromyographic muscle activity during walking in patients with intermittent claudication. *Journal of Vascular Surgery* 2016;**63**(3):664-72.

GRADE 2004

GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ* 2004;**328**:1490-4.

GRADEpro GDT 2015 [Computer program]

McMaster University (developed by Evidence Prime). GRADEpro GDT. Version accessed 4 July 2016. Hamilton (ON): McMaster University (developed by Evidence Prime), 2015.

Haas 2012

Haas TL, Lloyd PG, Yang HT, Terjung RL. Exercise training and peripheral arterial disease. *Comprehensive Physiology* 2012;**2**(4):2933-3017.

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Housley 1988

Housley E. Treating claudication in five words. *BMJ (Clinical Research Edition)* 1988;**296**(6635):1483-4.

Jolly 2009

Jolly K, Lip GYH, Taylor RS, Raftery J, Mant J, Lane D, et al. The Birmingham rehabilitation uptake maximisation study (BRUM): a randomised controlled trial comparing home-based with centre-based cardiac rehabilitation. *Heart* 2009;**95**(1):36-42.

King 2012

King S, Vanicek N, Mockford KA, Coughlin PA. The effect of a 3-month supervised exercise programme on gait parameters of patients with peripheral arterial disease and intermittent claudication. *Clinical Biomechanics (Bristol, Avon)* 2012;**27**(8):845-51.

King 2015

King S, Vanicek N, O'Brien TD. Dynamic muscle quality of the plantar flexors is impaired in claudicant patients with peripheral arterial disease and associated with poorer walking endurance. *Journal of Vascular Surgery* 2015;**62**(3):689-97. [DOI: [10.1016/j.jvs.2015.03.039](https://doi.org/10.1016/j.jvs.2015.03.039)]

Koutakis 2010

Koutakis P, Pipinos II, Myers SA, Stergiou N, Lynch TG, Johanning JM. Joint torques and powers are reduced during ambulation for both limbs in patients with unilateral claudication. *Journal of Vascular Surgery* 2010;**51**(1):80-8.

Koutakis 2015

Koutakis P, Myers SA, Cluff K, Ha DM, Haynatzki G, McComb RD, et al. Abnormal myofiber morphology and limb dysfunction in claudication. *Journal of Surgical Research* 2015;**196**(1):172-9. [DOI: [0.1016/j.jss.2015.02.011](https://doi.org/10.1016/j.jss.2015.02.011)]

Lauret 2012

Lauret GJ, van Dalen DC, Willigendael EM, Hendriks EJ, De Bie RA, Spronk S, et al. Supervised exercise therapy for intermittent claudication: current status and future perspectives. *Vascular* 2012;**20**(1):12-9.

Lauret 2012a

Lauret GJ, van Dalen HC, Hendriks HJ, van Sterkenburg SM, Koelemay MJ, Zeebregts CJ, et al. When is supervised exercise therapy considered useful in peripheral arterial occlusive disease? A nationwide survey among vascular surgeons. *European Journal of Vascular and Endovascular Surgery* 2012;**43**(3):308-12.

Leng 1993

Leng GC, Fowkes FGR. The epidemiology of peripheral arterial disease. *Vascular Medicine Review* 1993;**4**(1):5-18.

Lundgren 1989a

Lundgren F, Dahllöf AG, Scherstén T, Bylund-Fellenius AC. Muscle enzyme adaptation in patients with peripheral arterial insufficiency: spontaneous adaptation, effect of different treatments and consequences on walking performance. *Clinical Science* 1989;**77**(5):485-93.

McDermott 2006

McDermott MM. The magnitude of the problem of peripheral arterial disease: epidemiology and clinical significance. *Cleveland Clinic Journal of Medicine* 2006;**73**(Suppl 4):S2-7.

McDermott 2011

McDermott MM, Liu K, Ferrucci L, Tian L, Guralnik JM, Liao Y, et al. Greater sedentary hours and slower walking speed outside the home predict faster declines in functioning and adverse calf muscle changes in peripheral arterial disease. *Journal of the American College of Cardiology* 2011;**57**(23):2356-64.

McDermott 2012

McDermott MM, Liu K, Tian L, Guralnik JM, Criqui MH, Liao Y, et al. Calf muscle characteristics, strength measures, and mortality in peripheral arterial disease: a longitudinal study. *Journal of the American College of Cardiology* 2012;**59**(13):1159-67.

McDermott 2013

McDermott M. A home-based exercise intervention significantly improves walking performance in peripheral arterial disease: one-year follow-up from a randomized controlled trial. *Circulation* 2013;**128**:A14789.

McDermott 2013a

McDermott MM, Liu K, Guralnik JM, Criqui MH, Spring B, Tian L, et al. Home-based walking exercise intervention in peripheral artery disease: a randomized clinical trial. *JAMA* 2013;**310**(1):57-65.

McDermott 2014

McDermott M, Criqui M, Domanchuk K, Guralnik J, Kibbe M, Liu K, et al. Home-based exercise improves walking speed and prevents mobility loss in peripheral artery disease: a randomized controlled trial. *Circulation* 2014; Vol. 130:A11744.

McDermott 2014a

McDermott MM, Guralnik JM, Criqui MH, Ferrucci L, Zhao L, Liu K, et al. Home-based walking exercise in peripheral artery disease: 12-month follow-up of the GOALS randomized trial. *Journal of the American Heart Association* 2014;**3**(3):e000711.

Mockford 2014

Mockford KA, Gohil RA, Mazari F, Khan JA, Vanicek N, Coughlin PA, et al. Effect of supervised exercise on physical function and balance in patients with intermittent claudication. *British Journal of Surgery* 2014;**101**(4):356-62.

NCGC 2012

National Clinical Guideline Centre. Lower limb peripheral arterial disease. Diagnosis and management. NICE Clinical Guideline 147. <http://www.nice.org.uk/nicemedia/live/13856/60426/60426.pdf> (accessed September 2013).

Parmenter 2011

Parmenter BJ, Raymond J, Dinnen P, Singh MA. A systematic review of randomized controlled trials: walking versus alternative exercise prescription as treatment for intermittent claudication. *Atherosclerosis* 2011;**218**(1):1-12.

Parmenter 2013

Parmenter BJ, Raymond J, Fiatarone Singh MA. The effect of exercise on fitness and performance-based tests of function in intermittent claudication: a systematic review. *Sports Medicine (Auckland, N.Z.)* 2013;**43**(6):513-24.

Parmenter 2013a

Parmenter BJ, Raymond J, Dinnen PJ, Lusby RJ, Fiatarone Singh MA. Preliminary evidence that low ankle-brachial index is associated with reduced bilateral hip extensor strength and functional mobility in peripheral arterial disease. *Journal of Vascular Surgery* 2013;**57**(4):963-73.

Regensteiner 1993

Regensteiner JG, Wolfel EE, Brass EP, Carry MR, Ringel SP, Hargarten ME, et al. Chronic changes in skeletal muscle histology and function in peripheral arterial disease. *Circulation* 1993;**87**(2):413-21.

Rejeski 2014

Rejeski WJS, Spring B, Domanchuk K, Tao H, Tian L, Zhao L, et al. A group-mediated, home-based physical activity intervention for patients with peripheral artery disease: effects on social and psychological function. *Journal of Translational Medicine* 2014;**12**(29):1-8.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Ruell 1984

Ruell PA, Imperial ES, Bonar FJ, Thursby PF, Gass GC. Intermittent claudication. The effect of physical training on walking tolerance and venous lactate concentration. *European Journal of Applied Physiology and Occupational Physiology* 1984;**52**(4):420-5.

Sjöström 1982

Sjöström M, Neglén P, Fridén J, Eklöf B. Human skeletal muscle metabolism and morphology after temporary incomplete ischaemia. *European Journal of Clinical Investigation* 1982;**12**(1):69-79.

Terjung 1988

Terjung RL, Mathien GM, Erney TP, Ogilvie RW. Peripheral adaptations to low blood flow in muscle during exercise. *American Journal of Cardiology* 1988;**62**(8):15E-9E.

Tew 2013

Tew G, Copeland R, Le Faucheur A, Gernigon M, Nawaz S, Abraham P. Feasibility and validity of self-reported walking capacity in patients with intermittent claudication. *Journal of Vascular Surgery* 2013;**57**(5):1227-34.

Trialists 1994

Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy - I: Prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. *BMJ* 1994;**308**(6921):81-106.

Wang 2009

Wang J, Zhou S, Bronks R, Graham J, Myers S. Effects of supervised treadmill walking training on calf muscle capillarization in patients with intermittent claudication. *Angiology* 2009;**60**(1):36-41.

Willigendael 2005

Willigendael EM, Bendermacher BL, van der Berg C, Welten RJ, Prins MH, de Bie RA, et al. The development and implementation of a regional network of physiotherapists for exercise therapy in patients with peripheral arterial disease, a preliminary report. *BMC Health Services Research* 2005;**5**:49.

Yang 1991

Yang HT, Ogilvie RW, Terjung RL. Low-intensity training produces muscle adaptations in rats with femoral artery stenosis. *Journal of Applied Physiology* 1991;**71**(5):1822-9.

References to other published versions of this review

Lane 2014

Lane R, Ellis B, Watson L, Leng GC. Exercise for intermittent claudication. *Cochrane Database of Systematic Reviews* 2014, Issue 7. [DOI: [10.1002/14651858.CD000990.pub3](https://doi.org/10.1002/14651858.CD000990.pub3)]

Leng 2000

Leng GC, Fowler B, Ernst E. Exercise for intermittent claudication. *Cochrane Database of Systematic Reviews* 2000, Issue 2. [DOI: [10.1002/14651858.CD000990](https://doi.org/10.1002/14651858.CD000990)]

Watson 2008

Watson L, Ellis B, Leng GC. Exercise for intermittent claudication. *Cochrane Database of Systematic Reviews* 2008, Issue 4. [DOI: [10.1002/14651858.CD000990.pub2](https://doi.org/10.1002/14651858.CD000990.pub2)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Arosio 2001

Methods	Study design: RCT Method of randomisation: 'randomly divided' Exclusions post randomisation: not reported Losses to follow-up: not reported
Participants	Country: Italy Setting: hospital No. of participants: 24 Age: mean 65 years; 64.5 (55 to 69) years in exercise group; 66.4 (57 to 72) years in iloprost group Sex: all male Inclusion criteria: Fontaine stage II PAD with IC during past 4 to 6 years, confirmed by Doppler ultrasound, angiography, and ABI Exclusion criteria: not clearly stated but inferred to be smoking, severe hypertension, stroke, ischaemic attack, cerebrovascular disease, any drugs for PAD or other disease except transdermal clonidine for mild to moderate hypertension
Interventions	Intervention 1: physical exercise (n = 10) - walking, running, squat thrusts, 2 × day for 30 min, cycle ergometer for 30 min, plus 30 min constant load treadmill (2 mph, slope 0%) Intervention 2: iloprost (0.5 to 2 ng/kg/min) (n = 10) Duration: 14 days

Arosio 2001 (Continued)

Outcomes	Endogenous NO products Neutrophil adhesion ICD (m) and ACD (m) from constant load treadmill test (2 mph, slope 0%)
Notes	Exercise was interrupted at onset of pain, participant rested for 3 min, or until pain had gone, then re-summed activity until each 30-min block.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No description of randomisation method provided. Simply stated, 'the groups were randomly divided in two treatment regimens'
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No mention of whether all enrolled completed the study
Selective reporting (reporting bias)	Low risk	Prespecified outcome measures reported
Other bias	Unclear risk	Small numbers (n = 10 per treatment arm) and duration for only 14 days

Castro-Sanchez 2013

Methods	Study design: RCT Method of randomisation: randomised balanced (stratified) selection process Groups were balanced for type of medication received and Leriche-Fontaine stage (I or IIa), via a stratification system that generates a sequence of letters (from a table of correlatively ordered permutations) for each category and combination of categories. Exclusions post randomisation: 5; 3 in placebo (reasons: working hours, care for grandparents, and sprained ankle) and 2 in the physical therapy modality (reasons: terminally ill husband, and bed rest) Losses to follow-up: not reported
Participants	Country: Spain Setting: healthcare district No. of participants: 68

Castro-Sanchez 2013 (Continued)

Age: mean (SD) age, 53 (12) years; range, 41 to 65 years

Sex: 30 women and 38 men

Inclusion criteria: diagnosis of type 2 diabetes, postexercise ABI of 0.6 to 0.9, HbA1c of 7% to 13%, BMI of 25 to 40, and sedentary lifestyle

Exclusion criteria: other stages of PAD; peripheral venous insufficiency; cardiac, renal, or hepatic insufficiency; a cardiovascular event in the previous year; arterial pressure > 160/90 mmHg; LDL cholesterol > 160 mg/dL; an active smoking habit (in 3 previous years); and the presence of walking disorder, impaired skin integrity, or psychological or neurological disorder

Interventions	Intervention 1: Exercise group (n = 34) performed a session of 3 physical therapy modalities 2 times/d at home. Intervention 2: Placebo group (n = 34) received a treatment protocol 1 day/week with disconnected ultrasound electrotherapy equipment in dorsal and lumbar regions (15 minutes per region); these patients were instructed on the use of ultrasound equipment and were unaware that it was switched off. Duration: 20 weeks
Outcomes	Initial claudication walking distance (not reported in results); constant load treadmill test (3 km/h, 10% grade) Blood parameters: fibrinogen (mg/dL), haemoglobin (g/dL), glucose (mg/dL), cholesterol (mg/dL), high-density lipoprotein (HDL) cholesterol (mg/dL), LDL cholesterol (mg/dL), triglycerides (mg/dL), and HbA1c (%) Doppler flow velocity ABPI Cardiovascular risk score Heart rate during exercise test Assessments at 3 time points: baseline (before treatments), immediately after the final treatment session, and at 6 months after the conclusion of treatment
Notes	Home exercise protocol described by Boutroux 1980

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised balanced (stratified) selection process
Allocation concealment (selection bias)	Low risk	Groups were balanced for type of medication received and Leriche-Fontaine stage (I or IIa) via a stratification system that generates a sequence of letters (from a table of correlatively ordered permutations) for each category and combination of categories.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessors were blinded to the treatment group.

Exercise for intermittent claudication (Review)

Castro-Sanchez 2013 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Five dropouts were reported and excluded.
Selective reporting (reporting bias)	High risk	All outcomes stated - with the exception of initial claudication walking distance
Other bias	Low risk	No evidence of other bias

Ciuffetti 1994

Methods	Study design: RCT Method of randomisation: allocated by predetermined computer code. Not blinded Exclusions post randomisation: not reported Losses to follow-up: no dropouts reported
Participants	Country: Italy Setting: hospital No. of participants: 30 Age: 48 to 64 years Sex: male and female Inclusion criteria: stable maximal walking time (90 to 130 s) at 2 previous 6-monthly checks, plus stage II PAD confirmed by velocimetry and angiography Exclusion criteria: no h/o vascular surgery, coronary or cerebrovascular disease, DM; no factors affecting oxygen demand (e.g. anaemias); recent infection; treatment with vasodilators, antiplatelets, anti-coagulants, or drugs affecting haemorrhological parameters for previous 1 month
Interventions	Treatment: 1 h exercise at home daily plus twice-weekly supervision as out-patients. Home regimen: week 1, 500 m in 20 min; week 2, 1000 m in 40 min; week 3, 2000 m in 60 min on the flat Control: pentoxifylline 800 mg tds Duration: 3 months
Outcomes	Primary: treadmill test maximal walking distance (2 km/h on 12° slope)
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised by means of a predetermined computer code
Allocation concealment (selection bias)	Low risk	Predetermined computer code

Ciuffetti 1994 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated in paper
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts reported
Selective reporting (reporting bias)	Low risk	All outcomes stated
Other bias	Unclear risk	n = 15 per group; small numbers; groups matched for baseline gender and age

Collins 2005

Methods	Study design: RCT Method of randomisation: participants randomised after baseline testing completed; 4 groups, randomised 2 × 2 factorial analysis, permuted blocks: computer generated for vit E, sealed envelopes for exercise Exclusions post randomisation: none Losses to follow-up: 6 (4 polestriding group, 2 control group)
Participants	Country: USA Setting: community No. of participants: 52 (49 analysed) Age: 65.8 ± 7.1 years polestriding group; 68.0 ± 8.6 control group Sex: 51 M, 1 F Inclusion criteria: history of IC, ABI < 0.95 at rest or < 0.85 after exercise, IC pain a factor for arrested walking Exclusion criteria: vascular surgery, angioplasty in previous 6 months; other comorbid conditions that would interfere with participation in an exercise programme; currently taking vit E, warfarin, or pentoxifylline; unable to give informed consent
Interventions	Treatment: polestriding (n = 27), supervised training 3 times per week for 4 weeks, twice weekly for 8 weeks, once weekly for 4 weeks, biweekly for 4 weeks, unsupervised for 4 weeks Control: no exercise (n = 25), usual care. Seen biweekly for 3 months and monthly thereafter Duration: 24 weeks
Outcomes	Primary: ABI, maximum walking distance, oxygen uptake Secondary: health-related quality of life
Notes	All participants given \$6 travel for each visit and \$5 for each test battery completed, starting at \$25

Exercise for intermittent claudication (Review)

Collins 2005 (Continued)

Secondary analysis of the 2 × 2 factorial design of [Collins 2003](#); because results showed no influence of vit E on exercise, researchers combined exercise groups and compared findings with those of the non-exercise groups

The treadmill protocol began at a speed of 1.8 meter per hour (mph) and 0% grade. After the first 6 minutes, speed increased by 0.2 mph every 3 minutes. Per cent grade increased by 0.5% every 30 seconds

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants randomised after baseline testing completed; 4 groups, randomised 2 × 2 factorial analysis, permuted blocks, computer generated
Allocation concealment (selection bias)	Low risk	Randomised 2 × 2 factorial analysis, permuted blocks, computer generated for vit E. For exercise: sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinded to tablets, but not to exercise
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Assessors blinded to serum vit E levels
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported
Selective reporting (reporting bias)	Low risk	All outcomes stated
Other bias	Low risk	Controlled for additional visits by making it fair across all 4 groups

Crowther 2012

Methods	Study design: RCT Method of randomisation: "randomly allocated using a blinded protocol" Exclusions post randomisation: nil stated Losses to follow-up: For personal reasons, 6 participants (n = 5 CPAD-IC, and n = 1 TPAD-IC) withdrew over the course of the research study.
Participants	Country: Australia Setting: hospital No. of participants: N = 22; control (CPAD-IC, n = 11) or treatment (TPAD-IC, n = 11) group Age: CPAD-IC group: 67.1 (± 6.8), TPAD-IC group: 71.3 (± 8.5) Sex: 50% male Inclusion criteria: symptoms of IC, appropriate history of IC, imaging confirmation of PAD on lower limb duplex or CTA, and ability and willingness to attend for regular supervised exercise

Crowther 2012 (Continued)

Exclusion criteria: selection for surgical or endovascular intervention (n = 30), patient preference (n = 20) and requirement for mobility aids, obvious gait abnormalities (e.g. steppage, vaulting, circumduction, hip hiking), and medical conditions that influenced gait (e.g. orthopaedic conditions, neurological impairment)

Interventions	Treatment: 6-month supervised exercise programme Control: standard medical treatment as outlined in the Trans-Atlantic Inter-Society Consensus (TASC II) guidelines Duration: 6 months
Outcomes	Primary: submaximal walking economy and walking performance during a graded treadmill exercise test Secondary: body composition; resting ABI
Notes	Exercise programme initially consisted of intermittent supervised treadmill walking 3 days per week for a total time of 25 minutes at 3.2 km/h (0.88 m/s). Participants were required to walk until the pain level was perceived as 3 or 4 on the CPS. Exercise intensity (via treadmill grade and walking speed) and duration (25 minutes up to a maximum of 40 minutes) were progressively increased once the participant could walk continuously for 25 minutes at a level below 3 on the CPS. This exercise progression strategy was continued over the 6-month period of the study. The graded treadmill walking protocol consisted of a constant speed of 3.2 km/h (0.88 m/s) at an incline of 0% for the first 2 min, which was then increased by 2% every 2 minutes.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Description of randomisation limited to: "Participants were randomly allocated using a blinded protocol"
Allocation concealment (selection bias)	Low risk	Blinded protocol; no further details provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	High risk	6 participants lost to follow-up. Groups left uneven. However baseline data for the 2 groups were similar after participants who dropped out were excluded.
Selective reporting (reporting bias)	Low risk	All outcomes reported
Other bias	High risk	In Results section, numbers in CPAD and TPAD groups alternate. In Table 1, CPAD (n = 6), TPAD (n = 10); Table 2, CPAD (n = 10), TPAD (n = 6)

Cucato 2013

Methods	Study design: RCT Method of randomisation: randomised; nil else stated Exclusions post randomisation: nil stated Losses to follow-up: 4 from the walking group (reason: personal reasons)
Participants	Country: Brazil Setting: hospital No. of participants: 29 Age: control 61 ± 8 years; walking training 64 ± 6 years Sex: male Inclusion criteria: Stable symptoms of IC were recruited from a tertiary centre specialising in vascular disease. Male patients with Fontaine stage II symptoms of IC for longer than 6 months, an ABI < 0.90 at rest in 1 or 2 legs, and who were able to walk for at least 2 min at 3.2 km/h on a treadmill were invited to participate. Exclusion criteria: obesity (BMI < 30 kg/m²); use of beta-blockers, non-dihydropyridine calcium channel blockers, or peripheral vasodilators; inability to obtain the ABI; exercise tolerance limited by factors other than claudication (i.e. arrhythmias, cardiac symptoms, or exaggerated blood pressure rise); electrocardiogram response suggestive of myocardial ischaemia, and h/o revascularisation in the previous year
Interventions	Treatment: 15 × 2-min walking bouts, with 2-min intermittent rest periods. Exercise intensity was adjusted to maintain heart rate within 4 bpm above or below the heart rate of claudication pain onset (e.g. if the patient reported claudication pain onset in the maximal treadmill test at 100 bpm, the heart rate exercise zone was set at 96 to 104 bpm). Treadmill speed was set at 3.2 km/h, while the grade was adjusted to achieve the target heart rate. Twice-weekly 12-week walking exercise training programme Control: stretching exercise classes Duration: 12 weeks
Outcomes	Primary: to analyse the pain, cardiovascular and metabolic responses experienced by patients during walking exercise performed at the heart rate corresponding to claudication pain onset, and to investigate the effects of a 12-week walking training programme at this intensity on walking capacity in patients with IC Secondary: to determine whether low-volume walking exercise training programme at this intensity would induce walking performance improvements in this patient group Graded treadmill test: PFWD, MWD

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomised into 2 groups
Allocation concealment (selection bias)	Unclear risk	Not stated

Cucato 2013 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants given the nature of the intervention
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Participants from exercise group lost to follow-up only. Baseline demographics between 2 groups matched in spite of attrition
Selective reporting (reporting bias)	Low risk	All stated
Other bias	Unclear risk	Small numbers

Dahllof 1974

Methods	Study design: RCT Method of randomisation: states "randomised". Not blinded Exclusions post randomisation: not stated Losses to follow-up: not stated
Participants	Country: Sweden Setting: not stated No. of participants: 18 Age: 54 to 71 years Sex: male and female Inclusion criteria: IC for > 1 year Exclusion criteria: DM or IHD
Interventions	Treatment: 30-min training sessions 3 times weekly, supervised by a physiotherapist, including dynamic leg exercises beyond the appearance of pain Control: placebo tablets Duration: 6 months
Outcomes	Primary: treadmill test pain-free and maximal walking distance (max 1000 m), at 4 km/h Secondary: calf blood flow - venous occlusion plethysmography after ischaemic foot exercises
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Exercise for intermittent claudication (Review)

Dahllöf 1974 (Continued)

Random sequence generation (selection bias)	Unclear risk	Randomised, nil else reported
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinded to tablets, but not to exercise
Blinding of outcome assessment (detection bias) All outcomes	High risk	Physiotherapist ran exercise aspect.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No mention of whether all enrolled completed the study
Selective reporting (reporting bias)	Low risk	All outcomes stated
Other bias	Unclear risk	Small numbers (n = 18)

Gardner 2002

Methods	Study design: RCT Method of randomisation: not described Exclusions post randomisation: nil Losses to follow-up: 3 in treatment group, 6 in control group dropped out. More at 18 months. 31 remained.
Participants	Country: USA Setting: vascular clinics, newspaper and radio adverts No. of participants: 61 originally, 31 remained at 18 months Age: over 60 years Sex: 90% male Inclusion criteria: Fontaine II PAD Rose Questionnaire. ABI < 0.97 at rest. IC limiting factor on treadmill Exclusion criteria: other significant medical conditions limiting exercise tolerance, poorly controlled DM
Interventions	Treatment: 30-min training sessions 3 times weekly for 6 months, then twice weekly for 12 months Control: usual care Duration: 6 months and 18 months
Outcomes	Treadmill distance to claudication Maximum claudication distance ABI

Exercise for intermittent claudication (Review)

Gardner 2002 (Continued)

Peak oxygen uptake
Walking economy
Six-minute walking test distance
Accelerometer-derived physical activity
Walking Impairment Questionnaire
Calf blood flow
Health-related quality of life on SF-36
Self-perceived ambulatory measures

Notes Participants performed a progressive, graded treadmill protocol (2 mph, 0% grade with 2% increase every 2 minutes) until maximal claudication pain. Distance walked to onset of claudication pain, distance walked to maximal claudication pain, time to relief of claudication pain after the test, and peak oxygen uptake were measured.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomisation not specified
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants given the nature of the intervention
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated in paper
Incomplete outcome data (attrition bias) All outcomes	High risk	Numbers for attrition provided for all outcomes except the HRQoL, but reasons not stated 61 originally, 31 remained at 18 months: high loss to follow-up
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported on
Other bias	Low risk	No evidence of other bias

Gelin 2001

Methods Study design: RCT
Method of randomisation: block, computer
Exclusions post randomisation: nil stated
Losses to follow-up: 39 (13 control, 15 training, 11 invasive therapy) did not complete evaluation at 1 year. A total of 90 patients did not complete treatment as allocated, although analysis was based on intention to treat. 15 were unsuitable for surgery and 17 received angioplasty in surgery arm. 25 withdrew from quality of life section.

Participants Country: Sweden
Setting: vascular out-patients
No. of participants: 253
Age: mean 67 (range 45 to 81) years

Exercise for intermittent claudication (Review)

Gelin 2001 (Continued)

Sex: 67% male
Inclusion criteria: stable IC for > 6 months, ABI < 0.6 Maximum postischaemic blood flow < 25 mL/min/100 g, willing to undergo operations
Exclusion criteria: contraindication to surgery; other disorder limiting treadmill walking

Interventions	Treatment: • Supervised exercise training 30 min 3 times per week for 6 months, then 6 to 12 months 2 sessions per week • Invasive surgery/endovascular procedure, based on angiographic findings Control: observation only Duration: 1 year
Outcomes	Primary: mortality, ABI, amputation, treadmill distance; max postischaemic calf blood flow, big toe systolic pressure Secondary: blood pressure, haemoglobin, cholesterol, triglycerides, creatine, quality of life
Notes	Quality of life (data for 171 only, 18 changed group but were analysed in original) Randomised before pretreatment investigations. Low compliance in exercise group but classes offered for longer than most studies Results for ABI omitted standard deviations. Walking test was performed on a treadmill with a progressively inclining slope from 0° to 12°, simulating a gradually increasing workload.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised into 3 groups 'utilising a computer based algorithm, taking 21 assumed long-term prognostic variables into account'
Allocation concealment (selection bias)	Low risk	Randomisation performed by an independent nurse, who communicated the allocation group to the responsible physician
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated in paper
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	No competing interests declared. Funding transparent. High number of cross-overs described and discussed. However, greater degree of treatment dropouts in the exercise group; total numbers for this section (58%); has the potential to bias reported outcomes (PTA/surgery 80%)

GOALS 2013

Methods	Study design: RCT Method of randomisation: parallel-design randomised controlled clinical trial Exclusions post randomisation: 5 lost interest (control), 8 lost interest in exercise group Losses to follow-up: in CONSORT diagram; 3 deaths overall
Participants	Country: USA Setting: community No. of participants: 194 Age: 69.3 (9.5) exercise group; 71.0 (9.6) control group Sex: 49 (50.5%) men in exercise group; 48 (49.5%) in control group Inclusion criteria: ABI ≤ 0.90 in either leg. Individuals with a resting ABI ≥ 0.91 to ≤ 1.00 at baseline were eligible if their ABI dropped by at least 20% after a heel-rise test. Individuals with a resting ABI > 0.90 were eligible if they provided medical record documentation of lower extremity revascularisation or evidence of PAD from an accredited vascular laboratory. Exclusion criteria: potential participants with a below- or above-knee amputation; wheelchair confinement; inability to walk at least 50 feet without stopping; use of a walking aid other than a cane; inability to attend weekly sessions; walking impairment for a reason other than PAD, foot ulcer, or critical limb ischaemia; significant visual or hearing impairment; non-completion of the study run-in (attendance at 2 weekly health education sessions over a 3-week period); major surgery or lower extremity revascularisation during the previous 3 months or planned during the next 12 months; major medical illness including cancer treatment during the prior 12 months; current participation in another clinical trial or in another exercise trial within the past 3 months; completion of cardiac rehabilitation during the past 3 months; Parkinson disease; requirement of oxygen with activity or exercise; determination that exercise may be unsafe including having more than a class II New York Heart Association level of heart failure or angina; an increase in angina pectoris during the prior 6 months; an abnormal baseline exercise stress test; an exercise level similar to that targeted in the intervention at the time of recruitment; and a Mini-Mental State Examination score of < 23 at baseline
Interventions	Treatment: Our intervention applied principles from social cognitive theory, the group dynamics literature, and research on self-regulation to motivate participants to adhere to home-based walking exercise. Participants met once weekly for 90 minutes in a group with other PAD participants for the entire 6 months of the intervention. Forty-five minutes was devoted to facilitator-led discussions and 45 minutes to walking around an indoor track. Control: The health education control group attended weekly 60-minute group sessions with other PAD participants. Physicians and other healthcare professionals provided lectures on topics including managing hypertension, cancer screening, and vaccinations. Control group attended weekly lectures about topics not related to exercise. Duration: 6 months
Outcomes	Primary: 6-minute walk test Secondary: maximal treadmill walking time, pain-free treadmill walking time, physical activity, Walking Impairment Questionnaire (WIQ) scores, and Physical Health Composite Score (PCS) and Mental Health Composite Score (MCS) from the 12-item Medical Outcomes Study Short-Form Health Survey (SF-12)
Notes	
Risk of bias	

GOALS 2013 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	RCT: parallel-design randomised controlled clinical trial randomised by computer via a randomly permuted block method
Allocation concealment (selection bias)	Low risk	Eligible participants were randomised by computer via a randomly permuted block method, stratifying by baseline 6-minute walk performance.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All outcomes were measured before randomisation and at 6-month follow-up by assessors unaware of participants' group assignment.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Follow-up rates were 90.7% and 92.8% in the intervention and control groups. After exclusion of 3 participants who died before follow-up testing, follow-up rates were 91.7% and 94.7%.
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Multiple imputation was used by study authors to account for missing data at 6-month follow-up, including additional analyses in which a value of zero was assigned for missing 6-month follow-up data for decedents.

Guidon 2010

Methods	Study design: RCT Method of randomisation: "randomly allocated" Exclusions post randomisation: nil stated Losses to follow-up: 13 withdrew before follow-up
Participants	Country: Ireland Setting: not stated No. of participants: 44 initially randomised, 30 completed study Age: 67 ± 8.12 years Sex: 70.5% male Inclusion criteria: IC with ABI < 0.9 Exclusion criteria: nil stated
Interventions	Treatment: twice-weekly supervised exercise programme for 12 weeks Control: usual care Duration: 12 weeks
Outcomes	Walking Impairment Questionnaire

Exercise for intermittent claudication (Review)

Guidon 2010 (Continued)

Intermittent Claudication Questionnaire

SF-36

Notes RCT performed by School of Physiotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"randomly allocated"; no further information available
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	High risk	25% lost to follow-up
Selective reporting (reporting bias)	High risk	Only WIQ and ICQ discussed for 30/44 participants
Other bias	Unclear risk	RCT performed by School of Physiotherapy: who will require funding and may therefore be undertaking the study to provide weight for sustaining or undertaking additional classes

Hiatt 1990

Methods	Study design: RCT Method of randomisation: randomised in pairs by coin toss Exclusions post randomisation: 6 were discontinued from the study. 4 treated participants were discontinued: 3 for non-compliance with exercise training sessions and 1 for development of medical problems unrelated to PAD. Two control participants were discontinued: 1 for medical problems not due to PAD and the second for progression of arterial disease, as previously defined. Losses to follow-up: nil
Participants	Country: USA Setting: not stated No. of participants: 19 (25 enrolled, 6 discontinued); 10 treatment and 9 control Age: mean 61 years (treatment), 59 years (control) Sex: all male

Exercise for intermittent claudication (Review)

Hiatt 1990 (Continued)

Inclusion criteria: IC due to PAD (ABI < 0.95 at rest or < 0.85 after exercise)

Exclusion criteria: critical limb ischaemia; resting ankle blood pressure < 50 mmHg; unable to walk on the treadmill at a speed of 2 mph or an exercise capacity limited by angina, congestive heart failure, COPD, or arthritis; DM; vascular intervention in previous year; treatment with beta-blockers or pentoxifylline

Interventions	Treatment: programme of exercise 3 times each week (5 min warm up, 50 min intermittent isotonic resistive exercise, 5 min cool down) Control: maintain usual level of exercise Duration: 12 weeks
Outcomes	Primary: treadmill test maximal walking time (2 mph at 0% slope, with a subsequent 3.5% increase in slope every 3 min until forced to stop) Secondary: calf blood flow venous occlusion plethysmography, concentration of plasma carnitine at rest Subjective indicators: perceived pain during exercise; walking-limited distance
Notes	Hiatt 1990 included distance walked and claudication pain. Unfortunately these results were not reported by treatment group but simply correlated with treadmill performance. The correlation was good; therefore it may be assumed that significant improvement was experienced by those receiving exercise therapy. Treadmill test: graded treadmill test (2 mph 0% grade, with a subsequent 3.5% increase in grade every 3 minutes)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Coin toss
Allocation concealment (selection bias)	High risk	Coin toss
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Only for the carnitine analysis
Incomplete outcome data (attrition bias) All outcomes	High risk	All outcome data clearly reported. 25 enrolled, 6 discontinued (24% lost)
Selective reporting (reporting bias)	High risk	Maximum walking distance and claudication pain not reported for individual treatment group but correlated with treadmill performance.
Other bias	Unclear risk	Small numbers (n = 19)

Hiatt 1994

Methods	Study design: RCT Method of randomisation: not stated Exclusions post randomisation: 3 did not consent for muscle biopsy Losses to follow-up: 2 in control group
Participants	Country: USA Setting: not stated No. of participants: 29 Age: mean 67 years $\pm$ 6 years Sex: male only Inclusion criteria: 3-month h/o stable IC (Rose Questionnaire) limiting exercise sufficiently to affect ability to perform routine activity; ABI < 0.94 at rest, < 0.73 after exercise Exclusion criteria: rest pain, ulcer, gangrene; inability to walk on treadmill at > 2 mph; exercise limited by angina, CHF, COAD, or arthritis; no DM, vascular surgery, or PTA in previous year
Interventions	Treatment: supervised treadmill walking exercise (n = 10) or strength training (n = 9) and encouraged to walk alone for 2 days each week (treadmill 1 h 3 times a week, walking until moderate pain, then rest) Control: maintain usual level of activity (n = 10) Duration: 12 weeks
Outcomes	Primary: treadmill test pain-free and maximal walking distance (2 mph, 0% slope, with a subsequent 3.5% increase in slope every 3 min until forced to stop) ABI (resting and post exercise)
Notes	Trial continued after 12 weeks without a control group; therefore later results not included Strength training was less effective than treadmill exercise; only the latter was included in the meta-analysis for statistical reasons.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomised
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not specified
Incomplete outcome data (attrition bias)	Low risk	All outcome data reported

Hiatt 1994 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Small numbers

Hobbs 2005

Methods	Study design: RCT Method of randomisation: 2 × 2 factorial design, random number table Exclusions post randomisation: not reported Losses to follow-up: 4 of original 38 withdrew
Participants	Country: UK Setting: university department of vascular surgery, patients referred to IC clinic from primary or secondary care No. of participants: 34 Age: median 67 (63 to 72) years Sex: 27 men, 7 women Inclusion criteria: IC diagnosed by Edinburgh Claudication Questionnaire and reduced ABI < 0.9, reviewed after 3 to 6 months; max walking distance 20 to 500 m Exclusion criteria: significant aorto-iliac disease, inability to complete treadmill distance to absolute claudication distance, MI, transient ischaemic attack, stroke or PTA in past 3 months, CHF, bleeding diathesis, glomerular filtration rate < 20 mL/min, CYP3A4 or CYP2C19 inhibitor use
Interventions	Treatment: - Supervised exercise: 3-month, twice-weekly 1-hour physiotherapist-led exercise programme. Given videotape of programme and encouraged to take log of exercise at home - Cilostazol 100 mg twice daily, if side effects dosing halved for 1 week - Exercise as above plus cilostazol Control: best medical therapy Duration: 6 months
Outcomes	MWD Pain-free MWD ABI Thrombin antithrombin complex Prothrombin fragments 1 and 2 Plasminogen activator inhibitor Tissue plasminogen activator
Notes	Study authors contacted for means and SDs - received for BMT and supervised exercise groups Treadmill test: 3 km/h, 10% incline to MWD or 1000 m

Risk of bias

Hobbs 2005 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	2 × 2 factorial design randomisation, random number table
Allocation concealment (selection bias)	Low risk	Eligible participants were randomised in a 2 × 2 factorial design to continue BMT only or to receive BMT + supervised exercise, BMT+ cilostazol, or BMT + supervised exercise + cilostazol. The 2 × 2 factorial design is well recognised as one of the most robust study designs and allows for greater interrogation of the data, as well as allowing interaction of different treatments to be assessed.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated in paper
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Small numbers

Jansen 1991

Methods	Study design: RCT Method of randomisation: not described Exclusions post randomisation: not described Losses to follow-up: not described
Participants	Country: Germany Setting: community No. of participants: 48 Age: not described Sex: not described Inclusion criteria: PAD stage II Exclusion criteria: not stated
Interventions	Treatment: training on treadmill 3.5 km/h 10% slope for 2 hours twice per week under medical/physiotherapy supervision Control: no training

Jansen 1991 (Continued)

Duration: 2 years

Outcomes	Primary: treadmill walking distance, pain-free and maximum Secondary: ultrasound Doppler of arm and leg arteries
Notes	Treadmill test: 3.5 km/h, 10% incline Translated from German

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomised; no further information available
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Unable to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No mention of whether all enrolled completed the study
Selective reporting (reporting bias)	Low risk	All prespecified outcomes stated
Other bias	Low risk	No evidence of other bias

Kakkos 2005

Methods	Study design: RCT Method of randomisation: "blind block telephone procedure" by means of computer Exclusions post randomisation: 2 participants in the IPC group withdrew their consent before their 6 weekly appointments Losses to follow-up: 4 in supervised exercise group; 4 in IPC group
Participants	Country: England Setting: vascular out-patient clinic No. of participants: 34 Age: median and IQR: 66 (10.5) unsupervised exercise (USE) group; 69 (11.8) supervised exercise (SE) group; 66 (7) intermittent pneumatic compression (IPC) group Sex: 27 M in total; 8 M USE group, 11 M SE group, 8 M IPC group

Exercise for intermittent claudication (Review)

Kakkos 2005 (Continued)

Inclusion criteria: stable IC for > 6 months due to superficial femoral artery occlusion of ≥ 6 cm in length on ultrasound and/or angiogram

Exclusion criteria: duration of symptoms < 6 months, previous angioplasty or arterial surgery to symptomatic leg, MI within previous 6 months, inability to manage treadmill or training, any psychiatric illness or other reason making follow-up difficult, ischaemic rest pain, gangrene, ischaemic ulceration, inability to attend supervised programme, severe peripheral neuropathy, ABI > 0.9 at enrolment, non-compressible calf arteries, iliac occlusions or stenoses amenable to surgery or angioplasty, femoral artery occlusion < 6 cm; exercise capacity limited by angina, congestive heart failure, COPD, disease of spinal column, venous disease, neurological disease, mental illness, or arthritis

Interventions	Treatment: • Supervised (n = 12): 5-minute warm-up, 50-minute intermittent exercise, 5-minute cool-down. Attendance was 3 times a week for 6 months. • Unsupervised exercise (n = 9); advised to walk for approximately 45 minutes each day • Pneumatic foot compression (n = 13); to be used daily for 3-hour periods Duration: 6-month treatment period plus further follow-up at 12 months after treatment began (exercise advice given for second 6-month period)
Outcomes	Primary: ICD, ACD, ABI Secondary: SF-36, WIQ, IC Questionnaire
Notes	Study authors were successfully contacted for means and SDs. Treadmill test: 3.5 km/h, 10% incline

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised: blind, block telephone computer generated randomisation, undertaken by Clinical Trials and Evaluation Unit
Allocation concealment (selection bias)	Low risk	Independent allocation A blind, block 'telephone' randomisation procedure was performed by means of a computer. Randomisation was performed by the Clinical Trials and Evaluation Unit at the Royal Brompton Hospital in London.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	High risk	8 of 34 discontinued at 6 months (attrition rate of 26%)
Selective reporting (reporting bias)	Low risk	All data reported

Exercise for intermittent claudication (Review)

Kakkos 2005 (Continued)

Other bias	High risk	n = 8 per group. Power calculation required 15 participants per group.
------------	-----------	--

Larsen 1966

Methods	Study design: RCT Method of randomisation: randomised in pairs matched by age and disease, not blinded Exclusions post randomisation: 0 Losses to follow-up: 0
Participants	Country: Denmark Setting: not stated No. of participants: 14 Age: 44 to 65 years Sex: male and female Inclusion criteria: typical IC, stable for > 6 months Exclusion criteria: none stated
Interventions	Treatment: instructed to walk daily, in addition to normal activities Control: 1 placebo (lactose) tablet bd Duration: 6 months
Outcomes	Primary: treadmill test MWD (4.6 km/h, elevation of 0, 8, or 16 cm/m) Secondary: calf blood flow xenon clearance method
Notes	Treadmill test differed for each participant.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised in pairs matched by age and disease. Not blinded
Allocation concealment (selection bias)	High risk	Not blinded
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Unable to blind participants to exercise
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias)	Low risk	All outcomes stated

Exercise for intermittent claudication (Review)

Larsen 1966 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	All outcomes reported
Other bias	Unclear risk	Treadmill test different for each participant, small numbers (n = 14)

Leicht 2011

Methods	Study design: RCT Method of randomisation: (info from study author) Consecutive IC patients were randomly allocated to each group via sealed envelopes that were developed by a researcher not associated with the study. Exclusions post randomisation: nil stated Losses to follow-up: described in detail in CONSORT chart (Leicht 2011)
Participants	Country: Australia Setting: local hospital and university centre No. of participants: 25 PAD, 24 healthy age-matched controls Age: 66.9 years ($\pm$ 8 years) Sex: 14 male (56%) Inclusion criteria: PAD was confirmed based on absence of lower limb peripheral pulses, lower limb artery stenosis, or occlusion on duplex or computed tomographic angiography, and ankle brachial index (ABI) < 0.94. Exclusion criteria: inability to attend potential regular supervised exercise (n = 48), selected for surgical or endovascular intervention (n = 30), declined (n = 20), other medical condition influencing gait (n = 20), exhibited significant ectopy at rest (n = 3)
Interventions	- Conservative medical treatment (n = 13) - Supervised exercise (n = 12) consisted of treadmill walking 3 days per week for 25 to 40 minutes per day at a workload that induced intense to maximal claudication pain. Control: healthy age-matched controls (n = 24) Duration: 12 months
Outcomes	Primary: to compare heart rate variability (HRV) in patients with IC and in age-matched healthy adults Secondary: to examine the influence of an intense long-term (12-month) exercise programme on HRV in patients with IC PFWD, MWD, ABI
Notes	Treadmill test: 3.2 km/h and incline of 0%. The incline increased by 2% every 2 minutes until voluntary exhaustion or a maximum time of 25 minutes.

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Leicht 2011 (Continued)

Random sequence generation (selection bias)	Low risk	Randomised, consecutive IC patients were randomly allocated to each group via sealed envelopes that were developed by a researcher not associated with the study.
Allocation concealment (selection bias)	Low risk	Consecutive IC patients were randomly allocated to each group via sealed envelopes that were developed by a researcher not associated with the study.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data reported
Selective reporting (reporting bias)	Low risk	All outcome data reported
Other bias	Unclear risk	Small numbers

Mannarino 1991

Methods	Study design: RCT Method of randomisation: states random. Not blinded Exclusions post randomisation: not stated Losses to follow-up: none
Participants	Country: Italy Setting: not stated No. of participants: 20 Age: 48 to 75 years Sex: male and female Inclusion criteria: IC for > 2 years, stable for past 3 months, pain-free walking distance < 300 m Exclusion criteria: h/o angina, recent MI, or stroke; vascular surgery or PTA in previous 6 months; impaired cardiac or lung function; major liver, kidney, or metabolic disorders; infection; cancer; peptic ulcer
Interventions	Treatment: 1 h home exercises daily supervised via out-patients. Week 1: 500 m in 20 min; week 2: 1000 m in 40 min; week 3: 2000 m in 60 min Control: dipyridamole 75 mg tds plus aspirin 330 mg od Duration: 6 months
Outcomes	Primary: treadmill test pain-free and maximal walking time (2 km/h on 12° slope), ABI

Exercise for intermittent claudication (Review)

Mannarino 1991 (Continued)

Secondary: calf blood flow strain-gauge plethysmography

Notes A third group of 10 participants received antiplatelet treatment and exercise. These results are discussed but are not formally included in a meta-analysis.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomised; no further information available
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated in paper
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No mention of whether all enrolled completed the study
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Small numbers

McDermott 2008

Methods	Study design: RCT Method of randomisation: parallel assignment by computer via a randomly permuted block method; outcome assessor blinded Exclusions post randomisation: not described Losses to follow-up: described in flow chart (McDermott 2008)
Participants	Country: Chicago Setting: urban academic medical centre No. of participants: 156 Age: 70.6 ± 10.3 years Sex: male and female Inclusion criteria: 150 peripheral arterial disease patients with and without IC Exclusion criteria: above or below knee amputation, wheelchair confinement, inability to walk on a treadmill or perform progressive resistance training, inability to attend 3 times a week for 6 months, Class II NYHA heart failure or angina at rest or on minimal exertion, silent coronary ischaemia (ST de-

Exercise for intermittent claudication (Review)

McDermott 2008 (Continued)

pression of 1 mm) during baseline exercise test, ST-T wave changes or LBBB on baseline ECG, walking impairment not attributed to PAD, planned lower extremity revascularisation or major surgery within 12 months, MI or CABG during previous 3 months, current foot ulcer, ABI > 0.95, life expectancy < 12 months, does not speak English, dementia, poorly controlled BP, treated for cancer in the past 12 months, current significant exercise

Interventions	Treatment group 1: supervised treadmill exercise (6 months, 3 times a week; followed by a 6-month home-based programme). Initially 15 minutes of exercise, increased to 40 minutes by week 8. Initial treadmill walking speed 2.0 mph. Between weeks 8 and 24, intensity was increased weekly; either by grade or by speed, n = 51 Treatment group 2: lower extremity resistance training (6 months, 3 times a week; followed by a 6-month home-based programme). Participants performed 3 sets of 8 repetitions of knee extensions, leg press, leg curl exercises. Weights adjusted monthly until lifting 80% of 1 maximum repetition. 3 sets of 8 repetitions of squat and toe rises were also performed, n = 52 (1 dropped out) Control: usual care with diet and nutrition advice (11 sessions lasting 1 hour each over 6 months), n = 53 (1 dropped out) Duration: 6 months
Outcomes	6-Minute walking test distance Short physical performance battery Brachial artery flow-mediated dilatation Treadmill walking performance distance WIQ SF-36 physical functioning score
Notes	Treadmill test: Gardner Skinner Protocol

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised parallel assignment by computer via a randomly permuted block method
Allocation concealment (selection bias)	Low risk	Parallel assignment by computer via a randomly permuted block method
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Examiners were blinded to participant group assignment.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported

Exercise for intermittent claudication (Review)

McDermott 2008 (Continued)

Other bias	Unclear risk	Underpowered. Power calculations assumed that 50 people in each group would complete 6-month follow-up, and that 2 separate 2-sample t-tests using a 2-sided alpha of 0.05 would be conducted. The study was designed to have 80% power to detect a difference of 30 meters change in 6-minute walk distance and a difference of 0.97 change in the SPPB between baseline and 6-month follow-up between each exercise and control group. Only 50 completed the SET; 48 were in the control group, and 46 were in the lower extremity resistance group.
------------	--------------	---

McGuigan 2001

Methods	Study design: RCT Method of randomisation: randomly assigned to either of the 2 experimental groups Exclusions post randomisation: 2 participants from the training group and 2 from the control group withdrew from the study because of circumstances unrelated to the investigation. Losses to follow-up: not stated
Participants	Country: Australia Setting: not stated No. of participants: 20 (sample size of 18 needed) Age: not stated Sex: male and female Inclusion criteria: ABI < 0.94 at rest that decreased to < 0.73 after exercise Exclusion criteria: • Leg pain at rest • Ischaemic ulceration or gangrene • Inability to walk at least 2 km/h on a treadmill • Limited exercise capacity by factors other than IC (e.g. symptoms of angina, CCF, COPD, arthritis) • Vascular surgery or angioplasty undergone within the previous year • Smoking
Interventions	Treatment: n = 11. Resistance training program 3 days per week throughout the 24-week period. Minimum of 48 hours between sessions Control: n = 9 Duration: 24 weeks
Outcomes	Aims: to investigate effects of resistance training on walking performance, strength, and skeletal muscle adaptation Primary: fibre area and shifts in MHC isoforms Secondary: • Graded treadmill protocol - initially 3 km/h at 0% for 2 min, then increased 2% every 2 min with speed constant • Rate of perceived pain (scale 0 to 10)

McGuigan 2001 (Continued)

- Time in seconds to onset of claudication pain or maximal claudication pain
- 6-minute walk
- ABI
- Fibre type and distribution
- Muscle capillarisation

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not stated. Randomly assigned to either of the 2 experimental groups
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Two participants from the training group and 2 from the control group withdrew from the study because of circumstances unrelated to the investigation. Unclear what total numbers were
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Sample size was 18, which was not met, as 4 withdrew.

Mika 2005

Methods	Study design: RCT Method of randomisation: not described Exclusions post randomisation: not described Losses to follow-up: 18 withdrew (10 control, 8 exercise)
Participants	Country: Poland Setting: university department out-patient clinics No. of participants: 98 Age: 50 to 70 years Sex: male and female Inclusion criteria: PAD and IC (Fontaine stage II) stable for 3 months, PFWD 50 to 200 m at 3.2 km/h

Mika 2005 (Continued)

Exclusion criteria: angina, recent MI, vascular surgery in past 3 months, impaired cardiac or lung function, DM, cancer, kidney and liver disease, arthritis limiting walking, other contraindication to walking, those taking beta-blockers, pentoxifylline or other haemorheologically active drugs

Interventions	Treatment: 12-week programme of supervised pain-free treadmill exercise, 1 hour per day 3 times a week of repetitive walking exercise Control: usual care Duration: 3 months
Outcomes	PFWD Total leukocyte count Neutrophil count Microalbuminuria
Notes	Treadmill test: 3.2 km/h, 12% incline

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomised; no further details available
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinded assessor. Testing and analyses were conducted by qualified medical staff blinded to participants' group assignment.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Low risk	No evidence of other bias

Mika 2006

Methods	Study design: RCT Method of randomisation: not described Exclusions post randomisation: not described Losses to follow-up: 5
Participants	Country: Poland

Exercise for intermittent claudication (Review)

Mika 2006 (Continued)

Setting: university department clinics

No. of participants: 60

Age: 50 to 70 years

Sex: male and female

Inclusion criteria: Fontaine stage II PAD, IC limiting walking and stable for 3 months, ABI < 0.9

Exclusion criteria: angina; recent MI; vascular surgery in past 3 months; impaired cardiac or lung function; DM; cancer, kidney, and liver disease; arthritis limiting walking; inability to walk at 3.2 km/h; taking beta-blockers, pentoxifylline, or other haemorheologically active drugs

Interventions	Treatment: 12-week programme of supervised pain-free MD treadmill exercise, 1 hour per day 3 times a week of repetitive walking exercise Control: usual care, to maintain normal activity level Duration: 3 months
Outcomes	Pain-free walking time Maximum walking time Red cell deformability - erythrocyte elongation index
Notes	Treadmill test: 3.2 km/h 0% for 3 minutes followed by an increase in grade of 3.5% every 3 minutes (speed remained constant)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomised; no further details available
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Unable to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	5 lost to follow-up (3 in exercise group, 2 in control group). All data on remaining 55 reported
Selective reporting (reporting bias)	Low risk	All data on remaining 55 reported
Other bias	Unclear risk	Patients who were unable to walk on the treadmill at a speed of at least 3.2 km/h were also excluded.

Mika 2011

Methods	Study design: RCT Method of randomisation: not stated Exclusions post randomisation: not stated Losses to follow-up: 7 dropped out
Participants	Country: Poland Setting: hospital; vascular out-patient clinic No. of participants: 68 Age: 63.5 years (training group), 62.1 years (control group) 60 male, 8 female Inclusion criteria: PAD and IC – Fontaine stage II, aged 50 to 70 years, ABI < 0.9 at rest and 0.75 after exercise. All included patients had stable claudication distance and were able to walk no less than 150 m without pain. Exclusion criteria: rest pain, gangrene, ulceration, history of angina pectoris, recent MI or vascular surgery within the previous year, impaired cardiac or lung function, DM, cancer, or kidney and liver disease. Also, patients with arthritis who were unable to walk on the treadmill at a speed of at least 3.2 km/h. Additionally, women in menopausal status and those taking oestrogen were excluded because of the possible effects of these factors on HDL cholesterol level. None of the participants in the study was taking beta-blockers.
Interventions	Treatment: 12-week SEP, sessions conducted in the morning, 3 times per week Repetitive walking exercise with 3-min resting intervals. During each session, after 5 min of warm-up activities (free cycling on a stationary cycle ergometer), participants walked on the treadmill at a speed of 3.2 km/h at a grade that induced claudication pain within approximately 3 to 5 min Control: no change in physical activity All study participants were encouraged to stop smoking. Their diet was neither controlled nor modified throughout the study period. Duration: 3 months
Outcomes	Pain-free walking time (PFWT) Maximal walking time Haematocrit Plasma lipoproteins
Notes	Graded treadmill protocol (Gardner protocol): 3.2 km/h throughout the test, but the inclination from 0% (during initial stage) was raised by 2% every 2 min until maximal claudication pain occurred; without handrail

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomly assigned to either an intervention group (n = 34) or a control group (n = 34)

Mika 2011 (Continued)

Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Treadmill testing and biochemical analyses were conducted by medical staff blinded to participants' group assignment.
Incomplete outcome data (attrition bias) All outcomes	Low risk	7 dropouts discussed; rest of data complete
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Low risk	During an initial visit, participants were familiarised with treadmill walking (speed 3.2 km/h, 0% inclination) for 10 min. The treadmill test was repeated on the next day, and the mean of 2 measurements was used in data analysis. The treadmill was calibrated before each testing. Participants were instructed not to use handrail support during the test.

Sanderson 2006

Methods	Study design: RCT Method of randomisation: closed envelope system Exclusions post randomisation: nil Losses to follow-up: 1 in treadmill group
Participants	Country: Australia Setting: not stated No. of participants: 42 Age: mean 63 years Inclusion criteria: claudication lasting > 1 year, ABI < 0.9 Exclusion criteria: reduced cardiac function, rest pain, recent surgery or cardiac event, other medical conditions rendering exercise unsuitable
Interventions	Treatment: • Treadmill exercise • Cycling Both groups exercised 3 times a week for 6 weeks Control: no exercise Duration: 6 weeks
Outcomes	MWD and PFWD and cycling tests Submaximal and peak physiological response

Exercise for intermittent claudication (Review)

Sanderson 2006 (Continued)

ABI

Notes	Stratified by gender, presence of diabetes, then randomised Results given as mean differences The maximal graded walking test was performed on a motorised treadmill (TrackMaster TMX425CP, Newton, Kan) at a constant speed of 2.7 km/h. The treadmill gradient was set at 0% for the first 5 minutes of the test, then it was increased by 2% every 3 min until the participant failed to sustain the task.
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Closed envelope system
Allocation concealment (selection bias)	Low risk	Closed envelope system
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated in paper
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Low risk	No evidence of other bias

Schlager 2011

Methods	Study design: RCT single centre Method of randomisation: computer-generated random digits in sealed envelopes (block-wise randomisation by 2) Exclusions post randomisation: nil Losses to follow-up: 1 (MI 3 months after study inclusion)
Participants	Country: Austria Setting: single-centre hospital No. of participants: 40 Age: mean age of 69.1 years (standard deviation of ± 9.5 years) Inclusion criteria: symptomatic PAD (Rutherford category I to III), diagnosed by clinical evaluation, oscillometric pulse wave measurements, ABI, duplex sonography, and or CT or MR angiography

Schlager 2011 (Continued)

Exclusion criteria: asymptomatic PAD, critical limb ischaemia and reduced exercise tolerance caused by other limitations than claudication (coronary artery disease, congestive heart failure, dyspnoea, uncontrolled blood pressure, any kind of restriction of the musculoskeletal system, which might have an influence on the efficiency of exercise training)

Interventions	Treatment: SEP with BMT (twice-weekly programme for 6 months). Warm-up period of 5 to 10 min; initial duration included 35 min of intermittent walking, which was increased by 5 min each session until 50 min of intermittent walking was accomplished. The workload of exercise training was set to a walking speed that elicited claudication symptoms within 3 to 5 min. Control: BMT Duration: 12 months
Outcomes	Endothelial progenitor cells (CD34+, CD133+, KDR+) Plasma levels of vascular endothelial growth factor (VEGF), asymmetrical dimethylarginine (ADMA), stromal cell derived factor-1 (SDF-1) Maximum walking distance
Notes	Treadmill test: 3.2 km/h, 12% incline

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random digits in sealed envelopes (block-wise randomisation by 2)
Allocation concealment (selection bias)	Low risk	Computer-generated random digits in sealed envelopes (block-wise randomisation by 2)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Treadmill exercise tests were routinely performed by 2 experienced medical technical assistants, who were blinded to the assigned treatment arm and otherwise were not involved in the study. Participants were instructed to report all symptoms and discomforts (claudication and other than claudication) occurring during the test, but assistants refrained from encouraging participants during treadmill exercise tests.
Incomplete outcome data (attrition bias) All outcomes	Low risk	One out of 40 participants did not complete the study.
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Low risk	No evidence of other bias

Tew 2009

Methods	Study design: RCT
---------	-------------------

Exercise for intermittent claudication (Review)

Tew 2009 (Continued)

Method of randomisation: computer programme (nQuery Advisor 6.0 Statistical solutions)
Not blinded

Exclusions post randomisation: none

Losses to follow-up: 2 withdrew from the exercise group, and 4 withdrew from the control group; 1 participant died of a heart attack, 1 developed a lower limb ulcer that required revascularisation surgery, 1 was identified as having a popliteal artery aneurysm, and 1 returned to full-time employment. The remaining 2 participants cited lack of time as their reason for withdrawal.

Participants	Country: England Setting: single-centre hospital No. of participants: 57 Age: 69 ± 9 years Sex: male and female Inclusion criteria: Fontaine stage II PAD, 12-month history of stable IC, ambulation limited by IC, resting ABI < 0.90 or ABPI drop > 15 mmHg post maximal walking exercise Exclusion criteria: absence of PAD, ABI unobtainable owing to incompressible vessels, Fontaine I or III, exercise limited by cause other than IC, history of IC < 12 months, revascularisation/other major surgery within past 12 months, pharmacological therapy specifically for IC (e.g. cilostazol)
Interventions	Treatment: • Arm crank exercise (twice-weekly exercise programme for 12 weeks at an intensity of 60% to 70% of the peak work rate achieved of the initial arm crank assessment. Patients trained in cycles of 2-minute exercise at 50 rev/min followed by 2 min of rest for a total of 40 min/session) Control: • Non-exercise group Duration: 3 months
Outcomes	Primary: lower limb oxygen delivery Secondary: NIRS, peak VO₂ kinetics
Notes	Treadmill test: 3.2 km/h, 0% grade with 1% increase every 1 min

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised via a computer programme (nQuery Advisor 6.0; Statistical Solutions)
Allocation concealment (selection bias)	Low risk	Computer programme (nQuery Advisor 6.0; Statistical Solutions)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias)	Unclear risk	Not stated

Tew 2009 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	2 from the exercise group and 4 from the control group
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Methodological limitations need to be considered: NIRS data; exact contribution of intracellular myoglobin to the StO ₂ signal is unclear, and subcutaneous fat thickness changes might have influenced findings. No relationship exists between calf skinfold and calf muscle StO ₂ during walking in patients with IC.

Tew 2015

Methods	Study design: RCT Method of randomisation: randomised ratio 2:3. Achieved via a block randomisation sequence (block size 10) generated before recruitment Exclusions post randomisation: none Losses to follow-up: 1 lost to follow-up, all accounted for within CONSORT diagram
Participants	Country: UK Setting: not stated No. of participants: 23 Age: > 18 Sex: men and women Inclusion criteria: age > 18 years and stable IC for > 3 months Exclusion criteria: CLI, planned or previous lower-limb revascularisation, presence of contraindications to exercise (e.g. unstable angina) or co-morbidities that limited walking to a greater extent than the IC (e.g. severe arthritis); major surgery, MI, or CVA in the previous 6 months
Interventions	Treatment: usual care + SEDRIC (structured education for rehabilitation in intermittent claudication programme that promotes self-managed walking in people with IC) Control: usual care (included a brief info leaflet on PAD) Duration: 6 weeks
Outcomes	Mean daily step count Pain-free and maximum walking distances 6-min walking distance WELCH questionnaire, WIQ, EQ-5D, Intermittent Claudication Questionnaire
Notes	Gardner incremental treadmill test: 3.2 km/h, 0% grade, with 2% increase every 2 min

Tew 2015 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised ratio 2:3. Achieved via a block randomisation sequence (block size 10) generated before recruitment
Allocation concealment (selection bias)	Low risk	Implemented by an individual who was not involved in recruitment or data gathering processes
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Group allocation was concealed from participants while baseline information was gathered.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	All stated
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Baseline data showed that the intervention group had slightly higher levels of physical activity, walking capacity, and quality of life than the control group.

Tisi 1997

Methods	Study design: RCT Method of randomisation: randomised by sealed envelope, weighted 70:40:40 (angioplasty:exercise:observation) 19/25 randomised to PTA had unsuitable lesions, and the technique failed in one other. Not blinded Intention-to-treat analysis Exclusions post randomisation: none stated Losses to follow-up: none stated
Participants	Country: England Setting: general hospital No. of participants: 67 Age: mean 69.3 years Sex: male and female Inclusion criteria: stable IC > 6 months, positive Edinburgh Claudication Questionnaire, ABI < 0.8 and > 30 mmHg drop in ankle systolic pressure on exercise, walking distance 50 to 250 m on treadmill (3 km/h, 10% gradient) Exclusion criteria: intervention for IC in past 6 months, exercise limited by other factors, concurrent medical disease, treatment with steroids, inability to complete assessment visits

Tisi 1997 (Continued)

Interventions	Treatment: • Exercise - series of active and passive leg exercises performed to the limit of exercise pain, supervised by a physiotherapist once weekly for 4 weeks. Also encouraged to exercise for 45 min daily at home, plus to walk 1 mile a day; n = 22 • PTA, n = 28 • Control: observation, plus advice given to all 3 groups (leaflet advising on weight loss, smoking, and exercise, plus 75 mg aspirin daily), n = 17 • Healthy controls, n = 15 Duration: 12 months
Outcomes	Primary: treadmill test - PFWD and MWD (3 km/h on 10% slope); ABI Secondary: Nottingham Health Profile
Notes	Results for the exercise versus angioplasty comparison have not been included in this review. Claudication and walking distance results not in usable format

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised by sealed envelope, weighted 70:40:40 (angioplasty:exercise:observation)
Allocation concealment (selection bias)	Unclear risk	Randomised by sealed envelope, weighted 70:40:40 (angioplasty:exercise:observation); opacity of envelopes not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated in paper
Incomplete outcome data (attrition bias) All outcomes	High risk	Of the 22 enrolled at baseline, 10 were present at 12 months in the exercise group. In the observation group, n = 17 at baseline, and n = 9 at 12 months. No reasons given for loss to F/U
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Low risk	No evidence of other bias

Tsai 2002

Methods	Study design: RCT Method of randomisation: not described Exclusions post randomisation: not described
---------	--

Tsai 2002 (Continued)

Losses to follow-up: 11 - 5 in treatment group, 6 in control group

Participants	Country: Taiwan Setting: community No. of participants: 64 Age: 76 years $\pm$ 4 Sex: 81% male Inclusion criteria: Fontaine II PAD on Rose Questionnaire. ABI < 0.95 Exclusion criteria: intervention for IC in past 3 months, exercise limited by other factors, rest pain, MI or unstable claudication in the past 3 months, history of angina on exertion
Interventions	Treatment: 12-week progressive rehabilitation programme, 3 times a week. Up to 30 minutes on the treadmill Control: usual care Duration: 12 weeks
Outcomes	Time to onset of pain Time to maximum pain 6-Minute walking test distance WIIQ Physical function Bodily pain Role limitation physical and emotional General health Mental health Social function Vitality

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomised; no further details available
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinded outcome assessor
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported

Tsai 2002 (Continued)

Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Low risk	No evidence of other bias

Wood 2006

Methods	Study design: RCT Method of randomisation: not stated by paper Not blinded Exclusions post randomisation: none stated Losses to follow-up: nil stated
Participants	Country: Australia Setting: vascular out-patient clinic at the Royal Brisbane and Women's Hospital No. of participants: 18 Age: 56 ± 9 control, 64 ± 6 treatment Sex: male and female Inclusion criteria: history (> 6 months) of stable PAD and IC Exclusion criteria: rest pain, unstable angina and/or uncontrolled hypertension, experienced loss of consciousness as a result of dizziness, had a bone or joint problem that could be exacerbated by exercise, had undergone vascular surgery within the past 6 months, had suffered a cerebrovascular or coronary event in the past 12 months, or resided more than 1 h by car from the laboratory. Patients not limited by claudication and those who experienced angina or demonstrated ischaemic ECG abnormalities during exercise were excluded from the study.
Interventions	Treatment: • 6 weeks of treadmill walking training, three 40-min supervised walking training sessions/week (n = 7) Control: sedentary control (n = 6) Participants in both groups were asked to continue their normal daily activities. Duration: 6 weeks
Outcomes	• To determine whether plasma vascular endothelial growth factor (VEGF) increases in response to acute exercise in patients with IC, given the potentially large hypoxic stimulus for VEGF release during exercise in this population • To determine whether this response is attenuated following 6 weeks of high-intensity exercise training
Notes	Treadmill test: 2.7 km/h, 0%, with grade increased 2% every 3 min

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"In part B participants were randomised to one of two study arms".

Wood 2006 (Continued)

Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Unable to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	n = 6 for control and n = 7 for training. Nil lost to follow-up
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Unclear risk	Small numbers in each arm

Zwierska 2005

Methods	Study design: RCT Method of randomisation: Randomisation was undertaken via the fishbowl technique (Baumgartner 1998). This involved drawing patient names out of a 'hat' and allocating them to respective groups at random. Randomisation was undertaken by an independent academic based at another site in Sheffield. Intention-to-treat analysis: yes Exclusions post randomisation: nil Losses to follow-up: 10
Participants	Country: UK Setting: not stated No. of participants: 104 Age: median 69 (50 to 89) Sex: 81% male in leg group, 78% in arm group, 73% in control group Inclusion criteria: stable symptomatic peripheral arterial disease Exclusion criteria: not stated
Interventions	Treatment: • Supervised upper limb aerobic exercise • Supervised lower limb aerobic exercise Control: no exercise Duration: 24 weeks
Outcomes	ICD

Exercise for intermittent claudication (Review)

Zwierska 2005 (Continued)

MWD
Peak heart rate
Peak oxygen consumption
Perceived exertion and pain
Physical activity status

Notes Results given as medians - study authors contacted successfully for more information

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomised Randomisation was undertaken via the fishbowl technique (Baumgartner and Strong 1998). This involved drawing patient names out of a 'hat' and allocating them to respective groups at random.
Allocation concealment (selection bias)	Low risk	Randomisation was undertaken by an independent academic based at another site in Sheffield.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not possible to blind participants
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Assessment of outcome measures was conducted by the same 2 staff members. Procedures were checked for consistency at random by the lead investigator at Sheffield Hallam University (J.S.), who was blinded to group assignment. Blinding of the 2 staff members who assessed outcome measures was not stated.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcome data clearly reported
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported
Other bias	Low risk	No evidence of other bias

ABI: ankle brachial index.

ABPI: ankle brachial pressure index.

ACD: absolute claudication distance (or maximum walking distance).

bd: twice a day.

BMI: body mass index.

BMT: best medical therapy.

bpm: beats per minute.

CABG: coronary artery bypass graft.

CCF: congestive cardiac failure.

CHF: congestive heart failure.

CLI: critical limb ischaemia.

COAD: chronic obstructive airways disease.

COPD: chronic obstructive pulmonary disease.

CPAD-IC: peripheral arterial disease with intermittent claudication control group.

CPS: composite pain scale.

CT: computed tomography.

Exercise for intermittent claudication (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

CTA: computed tomographic angiography.
CVA: cerebrovascular accident.
DM: diabetes mellitus.
ECG: electrocardiography.
EQ-5D: EurQoL Group Quality of Life Questionnaire based on 5 dimensions.
HbA1c: glycated haemoglobin.
HDL: high-density lipoprotein.
h/o: history of.
HRQoL: health-related quality of life.
HRV: heart rate variability.
ICQ: Intermittent Claudication Questionnaire.
IQR: interquartile ratio.
LBBB: left bundle branch block.
LDL: low-density lipoprotein.
IC: intermittent claudication.
ICD: intermittent claudication distance.
IHD: ischaemic heart disease.
IPC: intermittent pneumatic compression.
MI: myocardial infarction.
min: minutes.
MR: magnetic resonance.
MWD: maximum or maximal walking distance.
NIRS: near-infrared spectroscopy.
NO: nitric oxide.
NYHA: New York Heart Association.
od: once a day.
PAD: peripheral arterial disease.
PFWd; pain-free walking distance.
PTA: percutaneous transluminal angioplasty.
RCT: randomised controlled trial.
SD: standard deviation.
SDF-1: stromal cell derived factor-1.
SE: supervised exercise.
SEP: supervised exercise programme.
SET: supervised exercise therapy.
SF-36: Short Form-36.
SPPB: Short Physical Performance Battery.
TASC: Trans-Atlantic Inter-Society Consensus.
tds: three times a day
TPAD-IC: peripheral arterial disease with intermittent claudication treatment group.
USE: unsupervised exercise.
VEGF: vascular endothelial growth factor.
vit E: vitamin E.
VO₂: maximum volume of oxygen.
WELCH: Walking Estimated-Limitation Calculated by History.
WIQ: Walking Impairment Questionnaire.

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Allen 2010	Compares home and supervised based programmes with a focus on nitrite flux
Andreozzi 2008	All 4 study arms utilised exercise.
Aruna 2015	Unclear whether symptomatic PAD. Methods state only that patients had type 2 diabetes mellitus for > 4 years and ABI between 0.90 and 0.40.
Beutel 1985	Both groups included exercise; the RCT compared CO ₂ mineral baths vs CO ₂ gas baths.

Exercise for intermittent claudication (Review)

Study	Reason for exclusion
Bronas 2011	The usual care group was provided with an exercise regimen that participants could decide whether to follow or not. Therefore, the study compares supervised vs non-supervised exercise. "Control participants also received written instructions on how to exercise independently if they so chose to do".
Brotons 2011	This RCT focussed on all patients with "a diagnosis of ischaemic heart disease, stroke or peripheral arterial disease", and the "primary endpoint was the combination of all-cause mortality and hospital cardiovascular readmission". Separate outcomes for each group were not provided.
Buchwalsky 1974	2 training groups
Bulling 1991	Comparison of exercise plus ginkgo biloba and exercise vs placebo. No non-exercise control group
Cachovan 1999	Comparison of 2 different sequences of exercise. No non-exercise control group
Carmeli 2004	Not randomised
Cheetham 2004	2 exercise groups and no non-exercise control group
Choi 2012	2 exercise groups and no non-exercise control group. "Controlled clinical trial with two intervention arms: calf ergometry and treadmill training".
Cina 1996	Comparison of physical training vs low-dose heparin and physical training and placebo. No non-exercise control group
CLEVER 2009	This study compares relative clinical and cost-effectiveness of invasive revascularisation with stents vs supervised exercise rehabilitation.
Collins 2010	Satisfied inclusion criteria, but data not available in usable format despite attempt to contact trial author
Collins 2012	Comparison of 2 different sequences of exercise. No non-exercise control group. "Patients were randomized to a traditional walking program or walking with poles program of exercise training".
Creasy 1990	This study compared PTA vs supervised exercise therapy.
Crowther 2008	Supervised exercise vs usual care, which included exercise advice for both groups
Cucato 2011	Methodology not of an RCT: 8 participants randomly underwent 2 experimental sessions: a session of resistance exercise (6 exercises, 3 sets of 12, 10 and 8 repetitions with perceived exertion of 11 to 13 on the 15-grade Borg scale) and a control session (resting on exercise machines)
Cucato 2011a	The aim of this study was to compare effects of walking and strength training on cardiovascular responses assessed at rest and during exercise in patients with intermittent claudication.
Cucato 2015	Participants with IC participated in 2 experimental sessions in a random order. Walking exercise (WE) (15 × 2-min bouts of WE interpolated with 2-min rest intervals) and control (standing rest on a treadmill for 60 min). BP, cardiac output (CO: CO ₂ rebreathing), and cardiovascular autonomic modulation (spectral analysis of heart rate variability) were assessed before and after both experimental sessions during supine rest, and stroke volume and systemic vascular resistance were calculated.
Cunningham 2012	This trial assessed whether a brief psychological intervention could increase daily walking at 4 months. The control group received usual care plus researcher contact, and the treatment group

Study	Reason for exclusion
	received usual care and a brief psychological intervention to modify illness and walking beliefs and to develop a personalised walking action plan.
Dahllof 1976	This trial was not fully randomised, as the first 10 of the 34 participants were allocated to exercise before randomisation was introduced. There were several differences between treatment and control groups, including significantly higher cholesterol levels and maximal calf blood flow in the control group. Treatment effects may also have been masked because many control participants spontaneously undertook increased exercise during the trial period.
Dantas 2016	Randomised cross-over intervention study with outcome measures of systolic BP, diastolic BP, and heart rate
Dedes 2010	Compares calf ergometry vs standard treadmill training (both groups involve exercise)
Degischer 2002	Not a randomised trial, and both groups had exercise
Dittmar 1977	Exercise in addition to drug treatment
Ericsson 1970	This trial compared exercise with no treatment but was excluded from the review because of insufficient evidence that participants were randomised. The paper stated that participants were "divided into two groups", and trialists were approached to confirm the method of allocation. No reply was received; therefore it was decided to exclude results from the review, unless evidence to the contrary becomes available.
Ernst 1987	This trial compared participants on an exercise programme with those not exercised, but allocation was not strictly randomised, as participants were "assigned according to space on the exercise programme".
Ernst 1990	Comparison of treadmill exercise plus pentoxifylline and treadmill exercise plus placebo. No non-exercise control group
EXERT 2009	Participants in the control group will continue to receive usual care from their regular doctor for treatment of PAD and will be provided with written exercise instructions.
EXITPAD 2010	Both groups have compare supervised exercise vs walking advice.
Fakhry 2011	Home-based exercise compared to supervised exercise
Fitzgerald 1971	This trial compared an exercise regimen with no treatment, but no evidence in the report indicates that the groups were randomly allocated. The quality of the trial appears generally poor, with variable periods of follow-up and inclusion of some patients who did not have intermittent claudication.
Fowler 2002	Some patients were asymptomatic, and it was not possible to exclude them for data extraction purposes.
Gardner 2005	2 exercise groups compared
Gardner 2011	Home-based exercise compared to supervised exercise, and third control group given walking advice
Gardner 2012	No exercise, usual care control group encouraged to walk more on their own but did not receive specific recommendations regarding an exercise programme during the study
Gardner 2014	80 participants were randomised to home-based and supervised exercise programmes. Both groups received training.

Study	Reason for exclusion
Gardner 2014a	180 participants were randomised. The NEXT Step programme and the supervised exercise programme consisted of intermittent walking to mild to moderate claudication pain for 12 weeks, whereas controls performed light resistance training.
Gibbellini 2000	Satisfied inclusion criteria but no usable data, despite attempts to contact trial authors. Results divided into asymptomatic and symptomatic groups, not aggregated
Gibbs 2013	This study is not focussed on patients with PAD. Participants (n = 140) with uncomplicated Type 2 Diabetes Mellitus, and without known cardiovascular disease or PAD, aged 40 to 65 years, were randomised to supervised aerobic and resistance training 3 times per week for 6 months, or to a usual care control group. ABI was measured before and after the intervention.
Gottstein 1987	Both groups received training.
Greenhalgh 2008	Both groups included exercise with the addition of PTA as the modality being investigated.
Guidon 2013	Focused on dropout rate of trials. Recruitment to clinical trials of exercise presents significant challenges in the PAD population owing to the presence of coexisting cardiovascular and cerebrovascular disease, reluctance to exercise due to leg pain, and acceptance of reduced mobility as part of ageing. Early identification in primary care before the onset of significant comorbidity may ameliorate some of these issues.
Guirro 2015	A cross-over study was carried out with 1 session of each therapeutic resource (high-voltage electrical stimulation (HVES), continuous short-wave diathermy, or physical exercise), with a 7-day washout period between protocols.
Hobbs 2006	This study compared exercise with angioplasty.
Hodges 2008	Groups were given supervised exercise or normal care, which included walking advice.
Holm 1973	Suitable for review but no usable numerical data, despite attempt to contact trial author. Stated significant improvement in walking time in exercise group compared to placebo group
Jones 1996	Comparison of treadmill vs StairMaster. No non-exercise control group
Kiesewetter 1987	Both groups received intensive physical therapy. No non-exercise control group
Kono 2013	This study is not focussed on patients with PAD. An observer-blind randomised controlled trial that enrolled 70 patients (48 men, mean age 63.5 years) with acute non-cardioembolic mild ischaemic stroke. Participants were allocated in equal numbers to a lifestyle intervention group or a control group.
Krause 1976	Combines exercise with drug
Kruidenier 2011	Treatment groups received angioplasty or exercise and angioplasty. A third group of exercise is required to make the trial eligible for inclusion in the review.
Labs 1999	Comparison of constant-load and graded-load treadmill testing with and without beraprost sodium. No non-exercise control group
Lee 2007	Non-randomised (clinical and cost-effectiveness)
Leon 2005	Focussed on cardiovascular disease, not intermittent claudication
Lepantalo 1984	Comparison of exercise plus flunarizine and exercise plus placebo. No non-exercise control group

Study	Reason for exclusion
LIFE Study	Original study is a multi-centred RCT that compared the ability of a structured physical activity intervention vs a successful ageing intervention to prevent mobility disability in older sedentary people, not specifically those with PAD.
Lundgren 1989	This study compared exercise vs surgical reconstruction.
Maejima 2005	Satisfied inclusion criteria, but no data available after attempt to contact trial authors. Reported improvement in walking time in exercise group at 12 weeks
Mannarino 1988	Controlled, not randomised
Mannarino 1989	This trial compared an exercise regimen with nothing (placebo tablets), but the report suggested that groups were not allocated by an acceptable randomisation method. Trialists have been approached to confirm the method of allocation; therefore it may become possible to include the trial at a later date.
Martinez 2009	3 groups with different durations of a walking programme
Mays 2015	Participants with PAD randomised to a community-based walking exercise programme compared to usual care advice. Control group received verbal advice to exercise but no formal training.
Mazari 2010	This trial compares PTA, a supervised exercise programme (SEP), and combined treatment (PTA plus SEP) for intermittent claudication.
McDermott 2004	Participants did not have intermittent claudication.
Nawaz 1999	Both groups received exercise advice. No non-exercise control group
Nawaz 2001	Comparison of upper limb and lower limb exercise. Separate non-randomised control group
NCT01065740	Outcome measure of 6-minute walking distance, not treadmill walking distance
NCT01241747	Treadmill training vs light resistance training without any walking
NCT02075502	Participants with PAD randomised to an exercise programme in the community setting incorporating training, monitoring, and coaching compared with participants who received the standard of care (exercise advice)
NCT02879019	Both groups use exercise.
Necker 2003	Training after angioplasty
Nicolai 2010	This paper compared 4 different graded treadmill protocols in random order.
Nielsen 1977	Both groups had exercise. No non-exercise control group
Nordanstig 2011	"All patients received verbal training advice and a written programme for IC. The patients were instructed to walk 1/hr day and up to MWD as often as possible". No arm received no exercise advice.
Parr 2009	All groups had exercise, no non-exercise control group. 3-armed trial of upper body strength training programme, conventional exercise programme, and walking advice for the control group
Patterson 1997	Comparison of supervised exercise programme plus lectures and home-based exercise plus lectures. No non-exercise control group

Study	Reason for exclusion
Pinto 1997	2 exercise regimens
Presern-Strukelj 200	Comparison of standard exercise alone and standard exercise plus electrostimulation in amputees with PAD
PROPEL study	Comparison between those with and without PAD. Looks at baseline data without exercise
Riccioni 2010	No mention of randomisation in methods
Richardson 1991	This compared rocker-bottomed shoes vs normal shoes with respect to walking distances.
Riebe 2001	Comparison of 2 progressive treadmill tests. No non-exercise control group
Ritti-Dias 2010	No control arm; strength training vs walking
Rodrigues 2014	Randomised cross-over of participants with PAD who performed 2 experimental sessions: control (C) and resistance exercise (R). Both sessions were identical (8 exercises, 3 × 10 reps), except that R session was performed with intensity between 5 and 7 on the OMNI-RES scale, and the C session was performed without any load.
Saleem 2011	Each group received exercise training. No non-exercise control group
Savage 2001	Compared supervised exercise vs home-based exercise. No non-exercise control group
Scheffler 1991	Each group received exercise training. No non-exercise control group
Schlager 2011a	Satisfied inclusion criteria but data not available in usable format despite attempt to contact trial author
Schoneberger 1994	Controlled study, not randomised
Silvestro 2002	Age-matched control group
Slordahl 2005	2 exercise groups
Snabl 1958	Not a randomised controlled trial
Sonaglia 2013	2 exercise groups. Eligible PAD La Fontaine IIa-IIb participants were randomised into 2 groups. Group A was treated with physical therapy plus oral pRLX, 20 ug b.i.d. for 12 weeks, and group B received physical therapy alone.
Spronk 2009	This study compared endovascular revascularisation vs supervised exercise.
Stewart 2008	Supervised exercise vs exercise advice. No control group
Streminski 1992	Satisfied inclusion criteria but data not available in usable format despite attempt to contact trial author. Actovegin vs exercise. Results given as mean change. Reported improvement in pain-free walking distance in exercise group
SUPER study	This study compared initial PTA vs initial supervised exercise therapy.
Taft 2004	This study was not truly randomised and did not contain any outcomes of relevance to this review.
Tebbutt 2011	Satisfied inclusion criteria but data not available in usable format despite attempt to contact trial author

Study	Reason for exclusion
Thomson 1999	2 exercise groups
Treat-Jacobson 2012	Participants with IC were assessed before and after a 12-week progressive, 3 times a week, supervised aerobic arm exercise training programme. No control arm
Ventura 1984	Satisfied inclusion criteria but data not available in usable format despite attempt to contact trial author
Walker 2000	The non-exercise group was not randomised.
Waller 1988	Comparison of exercise when participants had smoked immediately before treadmill test vs exercise when participants had not smoked before treadmill test
Wang 2008	Treatment: Plantar Flexion Ergometer, this occurred 3 times a week for 8 weeks. Each session lasted 40 minutes with equal time on each leg. Control: received advice in accordance with existing exercise guidelines for PAD patients provided by the AHA
Wang 2010	The study design was that of a cross-over from control to exercise, not a randomised controlled trial.
Winterfeld 1983	Not a randomised controlled trial
Zwierska 2004	Both groups had exercise. No non-exercise control group

ABI: ankle brachial index.

AHA: American Heart Association.

BP: blood pressure.

b.i.d.: twice daily.

CO₂: carbon dioxide.

IC: intermittent claudication.

MWD: maximum walking distance.

PAD: peripheral arterial disease.

PTA: percutaneous transluminal angioplasty.

RCT: randomised controlled trial.

SEP: supervised exercise programme.

WE: walking exercise.

Characteristics of ongoing studies *[ordered by study ID]*

NCT01231360

Trial name or title	The Effect of Exercise Training on Skeletal Muscle Metabolism in PAD
Methods	Allocation: randomised Intervention model: parallel assignment
Participants	Ages eligible for study: 40 years and older Genders eligible for study: both Accepts healthy volunteers: yes Inclusion criteria: Claudication symptoms

NCT01231360 (Continued)

- ABI $\leq$ 0.9 in the symptomatic leg

Exclusion criteria:

- Diabetes
- Impaired fasting glucose
- Peripheral vascular intervention within prior 6 months
- Recent unstable angina
- MI or stroke within prior 6 months
- Changes to HMG-CoA reductase inhibitor (statin) within past 3 months
- Changes to pentoxifylline and/or cilostazol regimen within past 3 months or anticipated to be necessary during the study
- On Coumadin
- Exercise limitations for reasons other than IC (such as congestive heart failure, angina, chronic lung disease, or other disorders affecting the limb such as arthritis or neuropathy)
- Rest pain or ulcers due to CLI
- Lower extremity amputation

Interventions	Active comparator: exercise training Participants randomised to exercise training will participate in a 3-month treadmill exercise programme in 1-h training sessions 3 times per week as previously described. After a 5-minute warm-up period, exercise is initiated at a low workload of 2 mph at 0% grade. Participants walk until moderate claudication severity develops, then rest until the discomfort resolves, repeating until the total exercise period is completed. The intensity of treadmill exercise is increased as tolerated by increasing walking speed by 0.5 to 1 mph and/or grade by 1% to 2%. Participants are encouraged to continue the walking programme at home for at least 30 minutes on 2 separate occasions each week. Active comparator: normal routine Participants randomised to the routine activity control group will be asked to keep a log of their daily activities and return to the Vascular Research Center at weeks 4, 8, and 12, at which time they will be asked to return their log and undergo repeat treadmill testing and complete the 6-minute walking test.
Outcomes	Specific aim 1: to test the hypothesis that subjects with PAD and IC have altered expression of genes that regulate skeletal muscle metabolism Specific aim 2: to test the hypothesis that exercise training improves calf skeletal muscle insulin resistance and genes that regulate skeletal muscle metabolic function in PAD patients with intermittent claudication
Starting date	Estimated enrolment: 75 Study start date: October 2010 Estimated study completion date: June 2013 Estimated primary completion date: December 2012 (final data collection date for primary outcome measure)
Contact information	Reena Pande, MD, Brigham and Women's Hospital skadivar@partners.org Contact: 617-732-6320
Notes	NCT01231360

NCT01822457

Trial name or title	Effect of Nike Fuel Band on Exercise and Function in Claudicants: A Randomised Controlled Trial
---------------------	---

Exercise for intermittent claudication (Review)

NCT01822457 (Continued)

Methods	Allocation: randomised Intervention model: parallel assignment Masking: single-blind (outcomes assessor) Primary purpose: supportive care - Experimental: Nike Fuel Band (NFB). Participants will receive a Nike Fuel Band to encourage exercise. - No intervention: control standard follow-up
Participants	Inclusion criteria: - Age 40 to 80 years - Referred to vascular rehabilitation service at St Mary's Hospital (UK) - IC involving the calf muscles - Clinical and duplex investigations indicate superficial femoral artery stenosis or occlusion Exclusion criteria: - Clinical and duplex investigations indicate iliac disease - Major joint disease in lower limb or lumbar spine/entrapment syndrome - Significant cardiopulmonary limitations (NYHA > 1) - MWD > 500 m - Hospital in-patient/living in a care home *Unfamiliarity with required technology - History of dementia - Inability to mobilise independently (does not include walking aids) - IC not a limiting factor of mobilisation, limited by other medical problems - Inability to attend supervised exercise programme - Patient owns or uses any type of activity monitor - Uses a walking frame *Participants should be able to use the NFB technology with minimal assistance. Gender: both Ages: 40 to 80 years
Interventions	Device: Nike Fuel Band (NFB). The NFB is a wrist-worn sensor with a built-in accelerometer for motion quantification. It is programmed to estimate the number of steps taken per day, and also to predict energy expenditure in units known as Nike Fuel. Accompanying software allows the user to set daily targets and monitor his or her activity through a graphical user interface.
Outcomes	All outcomes assessed at 3 months Primary: MWD - standardised treadmill test Secondary: - PFWD - standardised treadmill test - Disease-specific quality of life via the VasculQoL Questionnaire - Mood via the Hospital Anxiety and Depression Scale score
Starting date	August 2013
Contact information	Pasha Normahani: Pn106@imperial.ac.uk , Richard M Kwasnicki: rmk107@imperial.ac.uk
Notes	Estimated completion: February 2015 (final data collection date for primary outcome measure)


ABI: ankle brachial index.
CLI: critical limb ischaemia.
HMG-CoA: rate-controlling enzyme of the mevalonate pathway.
IC: intermittent claudication.
mph: miles per hour.
MI: myocardial infarction.
MWD: maximum walking distance.
NFB: Nike Fuel Band.
NYHA: New York Heart Association.
PAD: peripheral arterial disease.
PFWD: pain-free treadmill walking distance.

DATA AND ANALYSES


Comparison 1. Overall outcomes: exercise regimen compared with placebo or usual care

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maximum walking distance (m)	10	500	Mean Difference (IV, Random, 95% CI)	120.36 [50.79, 189.92]
2 Pain-free walking distance (m)	9	391	Mean Difference (IV, Fixed, 95% CI)	82.11 [71.73, 92.48]
3 Maximum walking time (min)	12	577	Mean Difference (IV, Random, 95% CI)	4.51 [3.11, 5.92]
3.1 Usual care	11	563	Mean Difference (IV, Random, 95% CI)	4.47 [3.00, 5.94]
3.2 Placebo	1	14	Mean Difference (IV, Random, 95% CI)	5.20 [0.88, 9.52]
4 Pain-free walking time (min)	11	534	Mean Difference (IV, Random, 95% CI)	2.93 [1.77, 4.09]
4.1 Usual care	10	520	Mean Difference (IV, Random, 95% CI)	3.08 [1.82, 4.34]
4.2 Placebo	1	14	Mean Difference (IV, Random, 95% CI)	1.77 [0.55, 2.99]
5 Change in MWD/T	15	656	Mean Difference (IV, Random, 95% CI)	40.25 [28.64, 51.86]
6 Change in ICD/T	15	703	Mean Difference (IV, Random, 95% CI)	58.42 [44.20, 72.64]
7 Ankle brachial index	13	570	Mean Difference (IV, Random, 95% CI)	0.04 [-0.00, 0.08]
8 Mortality	5	540	Risk Ratio (M-H, Fixed, 95% CI)	0.92 [0.39, 2.17]
9 Amputation	1	177	Risk Ratio (M-H, Fixed, 95% CI)	0.20 [0.01, 4.15]
10 Peak exercise calf blood flow (mL/100 mL/min)	4	103	Mean Difference (IV, Fixed, 95% CI)	0.94 [-0.81, 2.69]
10.1 Usual care	2	71	Mean Difference (IV, Fixed, 95% CI)	2.83 [0.18, 5.49]
10.2 Placebo	2	32	Mean Difference (IV, Fixed, 95% CI)	-0.52 [-2.85, 1.82]


Analysis 1.1. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 1 Maximum walking distance (m).




Analysis 1.2. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 2 Pain-free walking distance (m).




Analysis 1.3. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 3 Maximum walking time (min).




Analysis 1.4. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 4 Pain-free walking time (min).







Analysis 1.5. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 5 Change in MWD/T.




Analysis 1.6. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 6 Change in ICD/T.




Analysis 1.7. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 7 Ankle brachial index.




Analysis 1.8. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 8 Mortality.




Analysis 1.9. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 9 Amputation.



Analysis 1.10. Comparison 1 Overall outcomes: exercise regimen compared with placebo or usual care, Outcome 10 Peak exercise calf blood flow (mL/100 mL/min).






Comparison 2. Three-monthly outcomes: exercise regimen compared with placebo or usual care


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maximum walking distance (m)	3	116	Mean Difference (IV, Random, 95% CI)	104.46 [-64.33, 273.24]
2 Pain-free walking distance (m)	3	156	Mean Difference (IV, Fixed, 95% CI)	88.70 [58.25, 119.15]
3 Maximum walking time (min)	5	172	Mean Difference (IV, Fixed, 95% CI)	6.05 [5.47, 6.62]
3.1 Usual care	5	172	Mean Difference (IV, Fixed, 95% CI)	6.05 [5.47, 6.62]
4 Pain-free walking time (min)	3	132	Mean Difference (IV, Fixed, 95% CI)	4.95 [4.38, 5.53]
5 Ankle brachial index	4	130	Mean Difference (IV, Fixed, 95% CI)	0.06 [0.01, 0.11]
6 Mortality	2		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
7 Quality of Life SF-36	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
7.1 Physical function	2	82	Mean Difference (IV, Fixed, 95% CI)	6.60 [2.37, 10.83]
7.2 Bodily pain	2	82	Mean Difference (IV, Fixed, 95% CI)	3.89 [-1.91, 9.68]
7.3 General health	2	82	Mean Difference (IV, Fixed, 95% CI)	4.52 [-0.01, 9.04]
7.4 Mental health	2	82	Mean Difference (IV, Fixed, 95% CI)	1.44 [-0.93, 3.81]
7.5 Role emotional	2	82	Mean Difference (IV, Fixed, 95% CI)	1.26 [-4.84, 7.36]
7.6 Social function	2	82	Mean Difference (IV, Fixed, 95% CI)	1.49 [-4.16, 7.14]
7.7 Vitality	2	82	Mean Difference (IV, Fixed, 95% CI)	5.55 [1.54, 9.56]
7.8 Role physical	2	82	Mean Difference (IV, Fixed, 95% CI)	10.31 [3.64, 16.98]
7.9 Physical Summary Score	1	29	Mean Difference (IV, Fixed, 95% CI)	2.58 [-4.29, 9.45]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7.10 Mental Summary Score	1	29	Mean Difference (IV, Fixed, 95% CI)	2.05 [-4.73, 8.83]
8 Peak exercise calf blood flow (mL/100 mL/min)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected


Analysis 2.1. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 1 Maximum walking distance (m).




Analysis 2.2. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 2 Pain-free walking distance (m).




Analysis 2.3. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 3 Maximum walking time (min).






Analysis 2.4. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 4 Pain-free walking time (min).



Analysis 2.5. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 5 Ankle brachial index.




Analysis 2.6. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 6 Mortality.


Study or subgroup	Exercise		Placebo/usual care		Risk Ratio		Risk Ratio M-H, Fixed, 95% CI
	n/N		n/N		M-H, Fixed, 95% CI		
GOALS 2013	0/88		0/90				Not estimable
Tew 2009	0/27		1/24				0.3[0.01,6.98]
					Placebo/usual care	0.005 0.1 1 10 200	Exercise

Analysis 2.7. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 7 Quality of Life SF-36.

Study or subgroup	Exercise		Placebo/usual care		Mean Difference Fixed, 95% CI	Weight	Mean Difference Fixed, 95% CI
	N	Mean(SD)	N	Mean(SD)			
2.7.1 Physical function							
Guidon 2010	17	39.9 (9.2)	12	38.5 (9)		39.62%	1.42[-5.3,8.14]
Tsai 2002	27	58 (10.6)	26	48 (9.6)		60.38%	10[4.56,15.44]
Subtotal ***	44		38			100%	6.6[2.37,10.83]
Heterogeneity: Tau ² =0; Chi ² =3.78, df=1(P=0.05); I ² =73.58%							
Test for overall effect: Z=3.06(P=0)							
2.7.2 Bodily pain							
Guidon 2010	17	45.2 (9.8)	12	41.5 (9.8)		64.64%	3.72[-3.49,10.93]
Tsai 2002	27	81.5 (18.4)	26	77.3 (17.8)		35.36%	4.2[-5.55,13.95]
Subtotal ***	44		38			100%	3.89[-1.91,9.68]
Heterogeneity: Tau ² =0; Chi ² =0.01, df=1(P=0.94); I ² =0%							
Test for overall effect: Z=1.32(P=0.19)							
2.7.3 General health							
Guidon 2010	17	44.1 (9.6)	12	44.2 (8.6)		45.87%	-0.06[-6.74,6.62]
Tsai 2002	27	64.8 (0.1)	26	56.4 (16)		54.13%	8.4[2.25,14.55]
Subtotal ***	44		38			100%	4.52[-0.01,9.04]
Heterogeneity: Tau ² =0; Chi ² =3.33, df=1(P=0.07); I ² =70.01%							
Test for overall effect: Z=1.96(P=0.05)							
2.7.4 Mental health							
Guidon 2010	17	51.5 (9)	12	49.8 (8.9)		12.94%	1.72[-4.86,8.3]
Tsai 2002	27	80.1 (0.1)	26	78.7 (6.6)		87.06%	1.4[-1.14,3.94]
Subtotal ***	44		38			100%	1.44[-0.93,3.81]
Heterogeneity: Tau ² =0; Chi ² =0.01, df=1(P=0.93); I ² =0%							
Test for overall effect: Z=1.19(P=0.23)							
2.7.5 Role emotional							
Guidon 2010	17	49 (9.7)	12	49.2 (14.4)		42.57%	-0.14[-9.49,9.21]
Tsai 2002	27	93.3 (21.1)	26	91 (3.2)		57.43%	2.3[-5.75,10.35]
Subtotal ***	44		38			100%	1.26[-4.84,7.36]
Heterogeneity: Tau ² =0; Chi ² =0.15, df=1(P=0.7); I ² =0%							
Test for overall effect: Z=0.41(P=0.69)							
2.7.6 Social function							
Guidon 2010	17	49.8 (10.9)	12	49.1 (9.1)		59.67%	0.67[-6.64,7.98]
Tsai 2002	27	85 (17.5)	26	82.3 (15.5)		40.33%	2.7[-6.19,11.59]
Placebo/usual care -50 -25 0 25 50 Exercise							






Analysis 2.8. Comparison 2 Three-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 8 Peak exercise calf blood flow (mL/100 mL/min).



Comparison 3. Six-monthly outcomes: exercise regimen compared with placebo or usual care

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maximum walking distance (m)	3	156	Mean Difference (IV, Random, 95% CI)	138.36 [22.39, 254.34]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2 Pain-free walking distance (m)	2	116	Mean Difference (IV, Fixed, 95% CI)	52.14 [6.83, 97.45]
3 Maximum walking time (min)	4	295	Mean Difference (IV, Fixed, 95% CI)	3.20 [2.04, 4.36]
3.1 Usual care	3	281	Mean Difference (IV, Fixed, 95% CI)	3.05 [1.84, 4.25]
3.2 Placebo	1	14	Mean Difference (IV, Fixed, 95% CI)	5.20 [0.88, 9.52]
4 Pain-free walking time (min)	5	292	Mean Difference (IV, Random, 95% CI)	2.32 [0.91, 3.74]
4.1 Usual care	4	278	Mean Difference (IV, Random, 95% CI)	2.72 [0.67, 4.77]
4.2 Placebo	1	14	Mean Difference (IV, Random, 95% CI)	1.77 [0.55, 3.00]
5 Ankle brachial index	6	240	Mean Difference (IV, Random, 95% CI)	0.05 [-0.01, 0.11]
6 Mortality	1	178	Risk Ratio (M-H, Fixed, 95% CI)	0.51 [0.05, 5.54]
7 Quality of Life SF-36	5		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
7.1 Physical summary score	5	429	Mean Difference (IV, Fixed, 95% CI)	2.15 [1.26, 3.04]
7.2 Mental summary score	4	343	Mean Difference (IV, Fixed, 95% CI)	3.76 [2.70, 4.82]
7.3 Physical function	2	85	Mean Difference (IV, Fixed, 95% CI)	9.78 [0.82, 18.74]
7.4 Bodily pain	2	85	Mean Difference (IV, Fixed, 95% CI)	4.85 [-3.79, 13.50]
7.5 General health	2	85	Mean Difference (IV, Fixed, 95% CI)	10.19 [1.83, 18.55]
7.6 Mental health	2	85	Mean Difference (IV, Fixed, 95% CI)	1.82 [-5.28, 8.92]
7.7 Role emotional	2	85	Mean Difference (IV, Fixed, 95% CI)	4.90 [-7.15, 16.94]
7.8 Social function	2	85	Mean Difference (IV, Fixed, 95% CI)	3.48 [-6.74, 13.71]
7.9 Vitality	2	85	Mean Difference (IV, Fixed, 95% CI)	5.32 [-2.57, 13.22]
7.10 Role physical	2	85	Mean Difference (IV, Fixed, 95% CI)	8.55 [-2.69, 19.79]
8 Peak exercise calf blood flow (mL/100 mL/min)	2	66	Mean Difference (IV, Fixed, 95% CI)	3.79 [0.51, 7.07]
8.1 Usual care	1	52	Mean Difference (IV, Fixed, 95% CI)	3.10 [-0.46, 6.66]
8.2 Placebo	1	14	Mean Difference (IV, Fixed, 95% CI)	7.57 [-0.78, 15.92]

Analysis 3.1. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 1 Maximum walking distance (m).**Analysis 3.2. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 2 Pain-free walking distance (m).****Analysis 3.3. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 3 Maximum walking time (min).**

Study or subgroup	Exercise		Placebo/usual care		Mean Difference Fixed, 95% CI	Weight	Mean Difference Fixed, 95% CI
	N	Mean(SD)	N	Mean(SD)			

Test for overall effect: $Z=5.42$ ($P<0.0001$)
Test for subgroup differences: $\text{Chi}^2=0.89$, $df=1$ ($P=0.35$), $I^2=0\%$

Placebo/usual care -20 -10 0 10 20 Exercise

Analysis 3.4. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 4 Pain-free walking time (min).

Study or subgroup	Exercise		Placebo/usual care		Mean Difference Random, 95% CI	Weight	Mean Difference Random, 95% CI
	N	Mean(SD)	N	Mean(SD)			

3.4.1 Usual care

Collins 2005	25	15.1 (4.5)	24	10.3 (4.7)		15.15%	4.8[2.22,7.38]
Crowther 2012	10	7.8 (5.4)	6	2.3 (1.3)		10.54%	5.43[1.91,8.95]
GOALS 2013	77	4.2 (3.1)	68	3.5 (2.8)		26.9%	0.61[-0.34,1.56]
McDermott 2008	35	5.5 (3.2)	33	3.7 (3.2)		22.53%	1.86[0.33,3.39]
Subtotal ***	147		131			75.13%	2.72[0.67,4.77]

Heterogeneity: $\text{Tau}^2=3.19$; $\text{Chi}^2=14.67$, $df=3$ ($P=0$); $I^2=79.55\%$
Test for overall effect: $Z=2.6$ ($P=0.01$)

3.4.2 Placebo

Larsen 1966	7	3.5 (1.4)	7	1.7 (0.9)		24.87%	1.77[0.55,3]
Subtotal ***	7		7			24.87%	1.77[0.55,3]

Heterogeneity: $\text{Tau}^2=0$; $\text{Chi}^2=0$, $df=0$ ($P<0.0001$); $I^2=100\%$
Test for overall effect: $Z=2.83$ ($P=0$)

Total ***

	154		138			100%	2.32[0.91,3.74]
--	------------	--	------------	--	--	-------------	------------------------

Heterogeneity: $\text{Tau}^2=1.7$; $\text{Chi}^2=14.82$, $df=4$ ($P=0.01$); $I^2=73.02\%$
Test for overall effect: $Z=3.22$ ($P=0$)
Test for subgroup differences: $\text{Chi}^2=0.61$, $df=1$ ($P=0.44$), $I^2=0\%$

Placebo/usual care -10 -5 0 5 10 Exercise

Analysis 3.5. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 5 Ankle brachial index.


Study or subgroup	Exercise		Placebo/usual care		Mean Difference Random, 95% CI	Weight	Mean Difference Random, 95% CI
	N	Mean(SD)	N	Mean(SD)			

Castro-Sanchez 2013	32	0.9 (0)	31	0.9 (0.1)		25.92%	0.03[0.01,0.06]
Collins 2005	25	0.7 (0.2)	24	0.7 (0.2)		13.78%	0[-0.11,0.11]
Crowther 2012	10	0.7 (0.1)	6	0.5 (0.1)		18.86%	0.17[0.1,0.24]
Gardner 2002	28	0.7 (0.2)	24	0.7 (0.2)		11.97%	-0.05[-0.18,0.08]
McGuigan 2001	11	0.3 (0.1)	9	0.3 (0.1)		14.14%	0.02[-0.09,0.13]
Schlager 2011	20	0.7 (0.1)	20	0.6 (0.2)		15.32%	0.1[0,0.2]
Total ***	126		114			100%	0.05[-0.01,0.11]


Heterogeneity: $\text{Tau}^2=0$; $\text{Chi}^2=15.97$, $df=5$ ($P=0.01$); $I^2=68.69\%$
Test for overall effect: $Z=1.79$ ($P=0.07$)


Placebo/usual care -0.5 -0.25 0 0.25 0.5 Exercise

Analysis 3.6. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 6 Mortality.




Analysis 3.7. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 7 Quality of Life SF-36.






Analysis 3.8. Comparison 3 Six-monthly outcomes: exercise regimen compared with placebo or usual care, Outcome 8 Peak exercise calf blood flow (mL/100 mL/min).



Comparison 4. Exercise regimen compared with antiplatelet therapy

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maximum walking time (min)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2 Ankle brachial index	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
3 Peak exercise calf blood flow (mL/100 mL/min)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected


Analysis 4.1. Comparison 4 Exercise regimen compared with antiplatelet therapy, Outcome 1 Maximum walking time (min).



Analysis 4.2. Comparison 4 Exercise regimen compared with antiplatelet therapy, Outcome 2 Ankle brachial index.

Study or subgroup	Exercise		Antiplatelet therapy		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Mannarino 1991	10	0.7 (0.2)	10	0.7 (0.3)		0[-0.22,0.22]
					Antiplatelet -2 -1 0 1 2 Exercise	


Analysis 4.3. Comparison 4 Exercise regimen compared with antiplatelet therapy, Outcome 3 Peak exercise calf blood flow (mL/100 mL/min).

Study or subgroup	Exercise		Antiplatelet therapy		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Mannarino 1991	10	10.7 (0.3)	10	8.5 (4)		2.18[-0.28,4.64]
					Antiplatelet -10 -5 0 5 10 Exercise	

Comparison 5. Exercise regimen compared with pentoxifylline therapy

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maximum walking time (min)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected


Analysis 5.1. Comparison 5 Exercise regimen compared with pentoxifylline therapy, Outcome 1 Maximum walking time (min).

Study or subgroup	Exercise		Pentoxifylline		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Ciuffetti 1994	15	3.2 (0.4)	15	3.6 (0.1)		-0.45[-0.66,-0.24]
					Pentoxifylline -4 -2 0 2 4 Exercise	


Comparison 6. Exercise regimen compared with iloprost therapy

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maximum walking distance (m)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2 Pain-free walking distance (m)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected

Analysis 6.1. Comparison 6 Exercise regimen compared with iloprost therapy, Outcome 1 Maximum walking distance (m).

Study or subgroup	Exercise		Iloprost		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Arosio 2001	12	598 (375.4)	12	401.2 (324.1)		196.8[-83.8,477.4]


Analysis 6.2. Comparison 6 Exercise regimen compared with iloprost therapy, Outcome 2 Pain-free walking distance (m).

Study or subgroup	Exercise		Iloprost		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Arosio 2001	12	344.5 (293.6)	12	155.8 (87.4)		188.7[15.38,362.02]


Comparison 7. Exercise regimen compared with pneumatic foot and calf compression

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maximum walking distance (m)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2 Pain-free walking distance (m)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
3 Mortality	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected


Analysis 7.1. Comparison 7 Exercise regimen compared with pneumatic foot and calf compression, Outcome 1 Maximum walking distance (m).

Study or subgroup	Exercise		Pneumatic compression		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Kakkos 2005	12	417.1 (329.8)	13	479 (500)		-61.9[-391.59,267.79]

Analysis 7.2. Comparison 7 Exercise regimen compared with pneumatic foot and calf compression, Outcome 2 Pain-free walking distance (m).

Study or subgroup	Exercise		Pneumatic compression		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Kakkos 2005	12	92.9 (39.8)	13	253.2 (510.8)		-160.3[-438.88,118.28]


Analysis 7.3. Comparison 7 Exercise regimen compared with pneumatic foot and calf compression, Outcome 3 Mortality.




Comparison 8. Exercise regimen compared with vitamin E

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 3-monthly maximum walking time (min)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
2 6-monthly maximum walking time (min)	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
3 3-monthly ABI	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
4 6-monthly ABI	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected


Analysis 8.1. Comparison 8 Exercise regimen compared with vitamin E, Outcome 1 3-monthly maximum walking time (min).




Analysis 8.2. Comparison 8 Exercise regimen compared with vitamin E, Outcome 2 6-monthly maximum walking time (min).



Analysis 8.3. Comparison 8 Exercise regimen compared with vitamin E, Outcome 3 3-monthly ABI.



Analysis 8.4. Comparison 8 Exercise regimen compared with vitamin E, Outcome 4 6-monthly ABI.

Study or subgroup	Exercise		Vitamin E		Mean Difference	Mean Difference
	N	Mean(SD)	N	Mean(SD)	Fixed, 95% CI	Fixed, 95% CI
Collins 2005	11	0.7 (0.3)	13	0.6 (0.1)		0.1[-0.09,0.29]

ADDITIONAL TABLES

Table 1. Functional status and quality of life data (QoL) for all studies

Study	Measure reported	Effect reported
Collins 2005	Walking Impairment Questionnaire (WIQ) - perceived distance and walking speed	On the basis of analysis of change scores, the polestriding group reported significantly greater perceived ability to walk distance than the control group at 4 (P = 0.05), 12 (P = 0.001), and 24 (P = 0.002) weeks. Moreover, the polestriding group rated their perceived ability to walk faster to be significantly greater than the control group at 4 (P = 0.03), 12 (P = 0.19), and 24 (P = 0.02) weeks. The groups' ratings were equivalent at baseline (P > 0.05). In the polestriding group, polestriding aggregate scores for both distance and speed improved significantly between baseline and 12 weeks (P < 0.0001) and baseline and 24 weeks (P < 0.0001), but not between 12 and 24 weeks (P > 0.015). The perception of walking speed and distance did not improve in the control group.
	Rand Short Form-36 (SF-36) - perceived physical function	Exercise significantly improved the Physical Component Summary score (PSS) of SF-36 when compared with usual care. Difference using the change score between polestriding and control groups for the PSS was significant; P = 0.03. There was no difference in the change on the Mental Component Summary score between groups.
Gardner 2002	Health-related quality of life (QoL) assessed with the Medical Outcomes Study Short Form-36 (MOS SF-36)	Baseline measures of health-related QoL were comparable in both groups. The physical and mental health composite scores of the MOS SF-36 were similar between the 2 groups and did not change during the study. Consequently, no analyses were performed on the individual subscales.
	WIQ	No change in the WIQ was identified. Baseline scores on the 3 WIQ subscales ranged between 32% and 52%. Although the exercise group increased by 22% and 34% on the distance and speed subscales, respectively, these changes were not significant and did not differ from the changes in controls.
Gelin 2001	Intermittent Claudication-specific Sickness Impact Profile (SIP) scale, SIPIC	Supervised physical training produced significant improvements from baseline in only 2 health-related quality of life (HRQoL) domains (SIP Recreation and pastimes (P < 0.05) and the single-item rating scale Physical Condition) and an ambiguous pattern of positive and negative trends in others. Unexpectedly, deterioration was most striking in functional health, where reductions appeared in 10 of 15 SIP categories. Compared with no treatment, physical training produced significantly greater improvement in only 1 HRQoL category: SIP recreation and pastimes. Improvement in this category, however, may possibly be accounted for by the opportunities for increased leisure activity afforded by participation in the ongoing training programme.
	Health-related QoL, QoL overall	No significant improvement was observed between training and control groups. Training produced small SRMs (mean change between assessments divided by

Table 1. Functional status and quality of life data (QoL) for all studies (Continued)

the standard deviation of change) (0.2 to < 0.5) on 4 HRQoL dimensions, of which 2 represented improvement.

Guidon 2010	Medical Outcomes Study Short Form-36	No significant differences between groups were demonstrated for any of the MOS SF-36 scores over the 3 time points (baseline, 12 weeks, and 1 year). This study was not included in the meta-analysis, as the MOS SF-36 differs from the standard SF-36 Questionnaire.
	Disease-specific QoL (ICQ)	Results show a statistically significant decrease ($P = 0.003$) in ICQ scores from baseline to 12-week follow-up (mean difference -9.74, 95% CI -3.76 to -15.71) in the exercise group, indicating improved quality of life. No significant difference was demonstrated in the control group.
	WIQ	In the exercise group, increases were observed in all WIQ scores, with a statistically significant increase ($P = 0.015$) in the WIQ Distance score (mean difference 14.28, 95% CI 2.96 to 25.61). In the control group, scores for the WIQ Stair-climbing and Distance categories decreased, with a marginal increase in the WIQ Speed score. None of these changes were significant.
Kakkos 2005	Short Form-36 (SF-36)	Score for the general health domain of the SF-36 was significantly improved at 1 year in individuals who used intermittent pneumatic compression (IPC). This study was not included in the meta-analysis, as the full dataset was not available in the study and could not be obtained from the study author.
	WIQ	IPC improved speed score of WIQ significantly. WIQ scores for walking distance, walking speed, and stair climbing were reduced in the unsupervised exercise group, remained stable in the supervised exercise group, and were increased in the IPC group.
	Intermittent Claudication Questionnaire (ICQ)	Supervised exercise and IPC reduced (improved) the ICQ score, but this was significant only in the IPC group.
McDermott 2008	WIQ	Distance score improved ($P = 0.02$) in the treadmill group when compared to the control group. This was not apparent in the other 2 domains (speed and stair climbing).
	SF-36 physical functioning score	Improved ($P = 0.04$) in the treadmill group when compared to the control group.
GOALS 2013	WIQ scores	Participants in the intervention group, when compared with those in the control group, improved their WIQ distance score (35.3 to 47.4 vs 33.3 to 34.4; mean difference 11.1, 95% CI 3.9 to 18.1; $P = 0.003$) and their WIQ speed score (36.1 to 47.7 vs 35.3 to 36.6; mean difference 10.4, 95% CI 3.4 to 17.4; $P = 0.004$) but not their WIQ stair-climbing score (48.9 to 57.3 vs 47.9 to 48.5; mean difference 7.9, 95% CI 0.00 to 15.8; $P = 0.05$).
	Physical Health Composite Score (PCS) and Mental Health Composite Score (MCS) scales from the 12-item Medical Outcomes Study Short Form Health Survey (SF-12)	Results show no between-group differences in change in the SF-12 PCS or MCS subscales.
Tew 2015	Intermittent Claudication Questionnaire (ICD)	The intervention group demonstrated improvement in the ICD score of -10.6 (95% CI -18.9 to -2.3).

Table 1. Functional status and quality of life data (QoL) for all studies *(Continued)*

Tisi 1997	Nottingham Health Profile (NHP)	A daily home exercise programme, supervised weekly by a physiotherapist for the first month, achieves good compliance, increased walking distances, and improved quality of life as assessed by the Nottingham Health Profile Questionnaire.
Tsai 2002	WIQ	Improved speed ($P < 0.001$) and stairs ($P < 0.001$) in the exercise group when compared to the control group at 12 weeks. No significant difference in the distance domain
	SF-36 - version 1	Perception of QoL increased significantly in the exercise group compared to usual care for the domains of physical function, role limitation, bodily pain, general health, and vitality. This study was included in the 3-month meta-analysis; however the SF-36 version 1 differs from the standard SF-36 Questionnaire.
Zwierska 2005	WIQ	Improvement in all 3 domains was seen in the upper limb group at 24 and 48 weeks when compared to the control group. The lower limb group improved in stair and speed domains at 24 and 48 weeks only when compared to the control group.
	SF-36 - version 2	At 6 weeks, improvement in general health was seen in the lower-limb group when compared to the control group. At 24 weeks, a significant improvement was seen in general health and vitality in the lower limb group when compared to the control group. The upper limb group significantly improved in physical function and mental health when compared to the control group.

APPENDICES

Appendix 1. CENTRAL search strategy

#1	MESH DESCRIPTOR Arteriosclerosis	868
#2	MESH DESCRIPTOR Arteriolosclerosis EXPLODE ALL TREES	0
#3	MESH DESCRIPTOR Arteriosclerosis Obliterans	71
#4	MESH DESCRIPTOR Atherosclerosis	614
#5	MESH DESCRIPTOR Arterial Occlusive Diseases	722
#6	MESH DESCRIPTOR Intermittent Claudication	710
#7	MESH DESCRIPTOR Ischemia	786
#8	MESH DESCRIPTOR Peripheral Vascular Diseases EXPLODE ALL TREES	2195
#9	(atherosclero* or arteriosclero* or PVD or PAOD or PAD):TI,AB,KY	9000

(Continued)

#10	((arter* or vascular or vein* or veno* or peripher*) near (occlus* or reocclus* or re-occlus* or steno* or obstruct* or lesio* or block* or harden* or stiff-en*)) :TI,AB,KY	8590
#11	(peripheral near3 dis*) :TI,AB,KY	3324
#12	(claudic* or IC) :TI,AB,KY	3004
#13	(isch* or CLI) :TI,AB,KY	23368
#14	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13	42290
#15	MESH DESCRIPTOR exercise EXPLODE ALL TREES	16154
#16	MESH DESCRIPTOR Exercise Therapy EXPLODE ALL TREES	8362
#17	MESH DESCRIPTOR dancing EXPLODE ALL TREES	110
#18	MESH DESCRIPTOR Sports EXPLODE ALL TREES	11532
#19	MESH DESCRIPTOR Physical Exertion	3523
#20	MESH DESCRIPTOR Leisure Activities	214
#21	((fitness near3 (train* or intervention* or protocol* or program* or therap* or activit* or regim* or centre* or center*))) :TI,AB,KY	1051
#22	((training or conditioning) near3 (circuit or intervention* or protocol* or program* or activit* or regim*))) :TI,AB,KY	8527
#23	((walk* or run* or treadmill or aerobic or swim* or danc* or weight or squat* or lunge or bend* or raise or cycling or step)) :TI,AB,KY	94487
#24	kinesiotherap* :TI,AB,KY	1533
#25	((physical near3 (exertion or endurance or therap* or conditioning or activit* or fitness or train*))) :TI,AB,KY	25488
#26	exercise :TI,AB,KY	47402
#27	#15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26	132771
#28	#14 AND #27	7702
#29	NOT SR-PVD:CC AND 30/09/2013 TO 31/10/2016:DL	268200
#30	#28 AND #29	2165

Appendix 2. Trials registries searches

Clinicaltrials.gov

42 studies found for: intermittent claudication | exercise

WHO

62 records for 61 trials found for: exercise AND intermittent claudication

ISRCTN

22 results exercise AND intermittent claudication

WHAT'S NEW

Date	Event	Description
15 November 2016	New citation required but conclusions have not changed	Searches rerun, 2 new studies included, 18 additional studies excluded. New author (AH) joined the review team. Text updated to reflect current Cochrane standards. 'Summary of findings' table added. Conclusions not changed
15 November 2016	New search has been performed	Searches rerun, 2 new studies included, 18 additional studies excluded

HISTORY

Protocol first published: Issue 2, 1997

Review first published: Issue 1, 1998

Date	Event	Description
9 November 2013	New search has been performed	Searches rerun, 11 new studies included, 45 additional studies excluded
9 November 2013	New citation required but conclusions have not changed	New author (RAL) joined the review team. Searches rerun, 11 new studies included, 45 additional studies excluded. Risk of bias tables completed for all included studies In comparison with previous versions of this review, this version no longer focuses on supervised versus non-supervised exercise, or invasive interventions compared with exercise, as these are covered in other Cochrane reviews.
18 June 2008	New search has been performed	12 new trials considered. Conclusions confirm findings of the previous review.
29 April 2008	New citation required but conclusions have not changed	Two new review authors
8 April 2008	Amended	Converted to new review format
25 October 1999	New citation required but conclusions have not changed	Substantive updates made
17 June 1998	New citation required and minor changes	After consultation with GCL, original November 1997 version substituted for June 1998 'update', as latter was an incorrect version. Some phrasing changed in June 1998, no changes to figures, so earlier version more appropriate. Edited abstract added too (24.2.99)

CONTRIBUTIONS OF AUTHORS

Risha Lane (RL): selected trials, assessed quality, extracted data, and revised text for the third and fourth updates.

Amy Harwood (AH): selected trials, assessed quality, and extracted data for the fourth update.

Lorna Watson (LW): commented on content of the fourth update.

Gillian Leng (GCL): commented on content of the fourth update.

DECLARATIONS OF INTEREST

RL: none known.

AH: none known.

LW: none known.

GCL: I am responsible for the National Institute for Health and Care Excellence (NICE) Implementation programme and I am the Executive Director of NICE. I am a trustee of the Royal Society of Medicine and the Guidelines International Network.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- The Chief Scientist Office, Scottish Executive Health Directorates, The Scottish Government, UK.

The Cochrane Vascular editorial base is supported by the Chief Scientist Office.

- National Institute for Health Research, UK.

The Cochrane Vascular editorial base is supported by a programme grant from the NIHR.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Since the time of the original protocol, various changes have had to be instigated with regards to bias assessments. These have moved from the Schutlz assessment to Jadad and finally to the current Cochrane risk of bias assessment method ([Higgins 2011](#)). We have reassessed all trials to comply with the current bias scoring system.

The protocol itself has not been altered; however we have applied strict adherence to the protocol with regards to usual care. As walking advice has often been advocated as part of usual care, we have considered inclusion of only non-exercise control groups within this review. This has been clarified and searched for in each paper, as walking advice would constitute unsupervised exercise, which was not the focus of this review.

To avoid overlap with other existing Cochrane reviews ([Antoniou 2017](#); [Fowkes 1998](#)), we removed from this review the comparisons exercise versus angioplasty and exercise versus surgery.

In addition to planned treadmill walking distance and walking time, we have presented the results as a percentage change.

In keeping with current Cochrane policy, this version includes a 'Summary of findings' table and GRADE assessment of the quality of evidence. We have amended secondary outcomes to reflect clinical relevance.

INDEX TERMS

Medical Subject Headings (MeSH)

Amputation [statistics & numerical data]; Ankle Brachial Index; Exercise Therapy [*methods]; Intermittent Claudication [*therapy]; Pain Management [methods]; Quality of Life; Randomized Controlled Trials as Topic; Time Factors; Walking

MeSH check words

Female; Humans; Male