

## *Research Article*

# **Positive Approximation and Interpolation Using Compactly Supported Radial Basis Functions**

**Jinming Wu, Xiaolei Zhang, and Lihui Peng**

*College of Statistics and Mathematics, Zhejiang Gongshang University, Hangzhou 310018, China*

Correspondence should be addressed to Jinming Wu, wujm97@yahoo.com.cn

Received 17 July 2009; Revised 27 December 2009; Accepted 21 February 2010

Academic Editor: Victoria Vampa

Copyright © 2010 Jinming Wu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We discuss the problem of constrained approximation and interpolation of scattered data by using compactly supported radial basis functions, subjected to the constraint of preserving positivity. The approaches are presented to compute positive approximation and interpolation by solving the two corresponding optimization problems. Numerical experiments are provided to illustrate that the proposed method is flexible.

## **1. Introduction**

The problem of scattered data interpolation consists of constructing a function that interpolates data values which are known at some scattered points. However, we often have some additional information that we wish to confine to interpolation. For example, we know the quantity from which the data is sampled, positive, monotonic, or convex. Thus, it is important to construct a function which satisfies the underlying constraints.

In the past twenty years, the problem of positivity-preserving scattered data interpolation has been considered by many researchers. One class of methods for scattered data interpolation require the points to be triangulated mainly by using piecewise (rational) cubic interpolation and derived the conditions for preserving the shape of positive data. For related literature, we refer to the papers [1–6] and the references therein.

Another major class of methods for scattered data interpolation does not involve any prior triangulation, and can be viewed as meshless method. The two main types are radial basis functions (RBFs), which include multiquadrics and thin-plate splines, and so forth, and Shepard-type methods, which include the modified quadratic Shepard approach. Both types, RBF and Shepard, are widely used in practical fields.

However, there has been relatively little work done on the imposition of constraints for these meshless methods. For RBF, Utreras [7] considered how positivity can be imposed

as a constraint in case of thin plate spline interpolation for two-dimensional data. For Shepard, Asim and Brodlie [8, 9] discussed the modified quadratic Shepard method, which interpolates scattered data of any dimensionality and can be constrained to preserve positivity and more general constraint.

In this paper, we show how radial basis functions method, which interpolates or approximates scattered data, can be constrained to preserve positivity.

## 2. Radial Basis Functions

A radial basis function (RBF) is a relatively simple multivariate function generated by a univariate function. Due to its simple form and good approximation behavior, the radial basis function approach has become an effective tool for multivariate scattered data interpolation during the last two decades [10–14].

For any given scattered data  $(x_j, f_j) \in \mathbb{R}^s \times \mathbb{R}, j = 1, \dots, m$ , where points  $x_1, \dots, x_m \in \mathbb{R}^s$  are pairwise distinct, the so-called radial basis function interpolation is to use a function  $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}$  to construct the interpolant in the form of

$$s(x) = \sum_{j=1}^m \lambda_j \phi(\|x - x_j\|), \quad (2.1)$$

satisfying

$$s(x_i) = \sum_{j=1}^m \lambda_j \phi(\|x_i - x_j\|) = f_i, \quad i = 1, \dots, m. \quad (2.2)$$

The positive definiteness of  $\phi$  guarantees that the above interpolation problem (2.2) possesses a unique solution and refers to  $\phi$  as a classical radial basis function. If  $\phi$  has compact support, then the positive definite linear system is sparse and reduces computational cost greatly. Thus, we bypass this problem by restricting  $\phi$  to have compact support.

Compactly supported radial basis functions (CSRBFs) have only recently been constructed. Wu first constructed a broad variety of CSRBF [15]. Shortly after this, Wendland constructed these functions such that they possess the lowest degree among all CSRBFs which are positive definiteness for given space dimension and prescribed order of smoothness [16]. They are radial basis functions which are positive definite on  $\mathbb{R}^s$  for a given space dimension  $s$  ( $PD_s$ ), belong to a prescribed smoothness class ( $C^{2k}$ ), are compactly supported and easy to evaluate. Some examples of such radial basis functions are given in Table 1.

It is a useful property and provides a good selection of Wendland's functions with respect to the order of continuity and the dimension of space. Thus, CSRBFs have become a popular tool for multivariate interpolation of large scattered data [17, 18].

In order to adapt the interpolation to scattered data of different densities, it is necessary to be able to scale the support of  $\phi$ . So from now on we assume that the radius  $\alpha$  of support of  $\phi$  is one and replace  $\phi$  by

$$\phi_\alpha(\cdot) = \phi\left(\frac{\cdot}{\alpha}\right), \quad \text{for } \alpha > 0. \quad (2.3)$$

**Table 1:** Some of Wendland's CSRBF  $\phi_{s,k} \in PD_s \cap C^{2k}$ .

$\phi_{1,0} = (1-r)_+^r,$	$PD_1 \cap C^0$
$\phi_{3,0} = (1-r)_+^2,$	$PD_3 \cap C^0$
$\phi_{3,1} = (1-r)_+^4(4r+1),$	$PD_3 \cap C^2$

Meanwhile, in order to achieve both the best possible approximation behavior and best possible stability with respect to the support of CSRBF, we adopt the strategy to choose the radius  $\alpha$  as done in paper [17] throughout this paper.

### 3. Positive Approximation for Positive Data

The problem we are addressing is the following. Given a set of  $m$  scattered data  $x_i, x_i \in \mathbb{R}^s$  with associated data values  $f_i, i = 1, 2, \dots, m$ , where  $f_i \geq 0$ , we seek an optimal approximating function  $F(x)$  such that  $F(x) \geq 0$  in least-squares sense.

we first construct the function  $F(x)$  as follows:

$$F(x) = \sum_{i=1}^N \lambda_i \phi_\alpha(\|x - c_i\|), \quad (3.1)$$

where  $\phi_\alpha(\|x - c_i\|)$  is a compactly supported radial basis function centered on  $c_i$  with radius  $\alpha$  of support. Usually, the number of  $N$  is less than  $m$  greatly and centers  $c_i$  are different from data points  $x_i$ .

Thus, the so-called positive approximation for positive data is reduced to solve the following optimization problem

$$\begin{aligned} \min \quad & \sum_{j=1}^m (F(x_j) - f_j)^2, \\ \text{subject to} \quad & F(x) \geq 0. \end{aligned} \quad (3.2)$$

Note that all  $\phi_\alpha(\|x - c_i\|)$  are positive and  $F(x)$  is a linear combination of  $\phi_\alpha(\|x - c_i\|)$ . So, we can restrict each  $\lambda_i$  to be positive in order to guarantee the function to be everywhere positive. This sufficient condition is the kernel idea of our proposed method.


Therefore, the problem of positive approximation is transformed into a quadratic optimization problem subjected to linear constraints

$$\begin{aligned} \min \quad & \sum_{j=1}^m (F(x_j) - f_j)^2, \\ \text{subject to} \quad & \lambda_1 \geq 0, \dots, \lambda_N \geq 0. \end{aligned} \quad (3.3)$$

Without loss of generality, we illustrate it with a very simple example in 1D.

**Table 2:** One dimensional data for the velocity of wind.

Times (min)	0	0.25	0.5	1	1.2	1.8	2
Velocity (km/min)	2	0.8	0.5	0.1	1	0.5	1

**Figure 1:** Positive approximation curve for data of wind velocity.

*Example 3.1.* The set of 7 data points in Table 2 shows the velocity of wind. The velocity is inherently positive, and we therefore require the resulted approximating function to preserve this property.

Here, we choose the Wendland's function as  $\phi(r) = (1 - r)_+^4(4r + 1)$ . Figure 1 shows the positive approximation curve applied to the set of data for the velocity of wind using our proposed method (3.3) when  $c_i$  are chosen the same as  $x_i$ .

Table 3 shows the values of Error =  $\sum_{j=1}^7 (F(x_j) - f_j)^2$  when the number of centers  $N$  is varied. Of course, the total error will become less if the centers  $c_i$  are chosen properly.

*Remark 3.2.* The proposed method has a good chance to work well if the data are coming from a function  $f$  which is a convolution of the kernel with a positive function  $g$ , because then  $f$  can be recovered well by an integration formula which has exactly the form of (3.1).

#### 4. Positive Interpolation for Positive Data

The problem of positive interpolation we are addressing is the following. Given a set of  $m$  scattered data  $x_i, x_i \in \mathbb{R}^s$  with associated data values  $f_i, i = 1, 2, \dots, m$ , where  $f_i \geq 0$ , the so-called positive interpolation is to construct a function  $F(x)$  such that  $F(x_i) = f_i, i = 1, \dots, m$  and  $F(x) \geq 0$ , where  $F(x)$  is a linear combination of the CSRBF.

The basic idea is also to restrict each combination coefficient  $\lambda_i$  to be positive in order to guarantee that  $F(x)$  is positive everywhere. Obviously, classical RBF interpolation (2.2) cannot guarantee that all  $\lambda_i$  are positive.

One feasible way is to choose  $n$  new added data  $y_i, i = m + 1, \dots, m + n$  and then turn to find the following interpolant:

$$F(x) = \sum_{i=1}^m \lambda_i \phi_\alpha(\|x - x_i\|) + \sum_{j=m+1}^{m+n} \lambda_j \phi_\beta(\|x - y_j\|) \quad (4.1)$$

**Table 3:** Total error for different parametric values of  $N$ .

$N$	3	5	7	9
Error	$6.24 \times 10^{-1}$	$8.37 \times 10^{-2}$	$5.04 \times 10^{-2}$	$7.80 \times 10^{-3}$

such that

$$\begin{aligned} F(x_i) &= f_i, \quad i = 1, 2, \dots, m, \\ \lambda_i &\geq 0, \quad i = 1, 2, \dots, m + n. \end{aligned} \quad (4.2)$$

The above interpolant will naturally arise an inevitable problem: what makes the system of linear equations (4.1) have a unique solution  $F(x)$  satisfying the linear constraints. In order to solve it in special cases, we require the following lemma.

**Lemma 4.1** (see [19, Gordan's Theorem]). *Let  $A_1, A_2, \dots, A_m$  be  $n$ -dimensional vectors; there does not exist a vector  $P$  such that  $A_i^T P < 0$  if and only if there exist nonnegative real numbers  $\lambda_1, \lambda_2, \dots, \lambda_m$  such that  $\sum_{i=1}^m \lambda_i A_i = 0$ .*

Undoubtedly, it is hard to give a general result which can determine whether there exists a solution when the added centers  $y_j$  and the radius  $\beta_j$  are chosen randomly. Thus, we discuss the following special case to choose centers  $y_j$  and the radius  $\beta_j$ .

For each data  $x_i, i = 1, \dots, m$ , if we choose the new added data  $y_i$  in the neighborhood of  $x_i$  randomly and choose the radius  $\beta_i$  such that the influence domain of  $\phi_{\beta_i}(\|x - y_i\|)$  only contains the data  $x_i$ , then for this case we have the following.

**Theorem 4.2.** *If the new added data  $y_j$  and radius  $\beta_j$  are chosen as above, then there exists a solution subjected to the constraints (4.2).*


*Proof.* Let

$$\begin{aligned} A_j^T &= -\left\{ \phi_{\beta_j}(\|x_i - y_j\|) \right\}_{i=1}^m = (0, \dots, 0, -b_j, 0, \dots, 0), \quad j = m + 1, \dots, 2m, \\ A_j^T &= -\left\{ \phi_{\alpha}(\|x_i - x_j\|) \right\}_{i=1}^m, \quad j = 1, 2, \dots, m, \quad A_{2m+1}^T = (f_1, \dots, f_m), \quad \forall f_i \geq 0, \end{aligned} \quad (4.3)$$


where  $b_j, j = m + 1, \dots, 2m$ , are positive real numbers.

It is easy to see that there does not exist a vector  $P$  such that  $A_i^T P < 0, i = m + 1, \dots, 2m + 1$ . According to Gordon lemma, we know that there exist nonnegative real numbers  $\lambda_1, \lambda_2, \dots, \lambda_{2m+1}$  such that  $\sum_{i=1}^{2m+1} \lambda_i A_i = 0$  and  $\lambda_{2m+1} > 0$ .

We naturally wish that there are many zero coefficients  $\lambda_j, j = m + 1, \dots, m + n$ , in the formulas (4.1). That is to say, we add as few data as possible. For example, if the original interpolation function is positive, then we certainly need not add any new data. So from now on we hope to minimize the objective function  $\text{sgn}(\lambda_{m+1}) + \dots + \text{sgn}(\lambda_{2m})$ , where  $\text{sgn}(\cdot)$  denotes the sign function.



**Figure 2:** One-dimensional data for the velocity of wind—CSRBF interpolation loses positivity.



**Figure 3:** Positive interpolation curve for data of wind velocity using piecewise cubic.

Therefore, an approach is presented to compute the positive interpolation by solving the following optimization problem:

$$\begin{aligned} & \min \quad (\text{sgn}(\lambda_{m+1}) + \cdots + \text{sgn}(\lambda_{2m})), \\ & \text{subject to} \quad \begin{cases} F(x_i) = f_i, & i = 1, 2, \dots, m, \\ \lambda_i \geq 0, & i = 1, 2, \dots, 2m. \end{cases} \end{aligned} \quad (4.4)$$

We illustrate it with an example using the same data given in Table 2 and make comparisons with several existing methods.  $\square$

*Example 4.3* (Example 3.1 is continued). In Figure 2, we show the original interpolation curve by using CSRBF without the constraint of positivity. Generally, the curve has good approximation behavior, but the curve goes beyond the range of the data values which makes physical nonsense and indeed the positivity is violated.

In Figure 3, the positive curve applied to the same set of positive data using piecewise cubic method [8] is shown. Meanwhile, we reveal the positive curve using constrained modified quadratic Shepard method (CMQS) [9] in Figure 4.




Figure 4: Positive interpolation curve for data of wind velocity using CMQS.




Figure 5: Positive interpolation curve for positive data.

By contrast, in Figure 5, we show the positive interpolation curve applied to the same data of wind velocity using our proposed approach (4.1). From the experiment, the constructed curve possesses relatively good approximation behavior in contrast to piecewise cubic method and constrained modified quadratic Shepard method.

*Remark 4.4.* It is pointed out that we generally add corresponding new data  $y_{m+k}$  if  $f_k$  is the minimal data among all value  $f_i$ ,  $i = 1, \dots, m$  instead of adding  $m$  new data directly to solve the positive interpolation in practical use.

## 5. Numerical Example

The strong and sufficient condition that all  $\lambda_i$  are positive may degrade the quality of the interpolating function, in comparison with the original unconstrained CSRBF interpolation. We evaluate the quality of the new interpolant by calculating the variances between the exact and calculated values on a set of test points by the following experiment in 2D.




Figure 6: Interpolated function  $S(x, y)$ .




Figure 7: Positive interpolating surface for  $S(x, y)$ .

*Example 5.1.* We illustrate it with the following function  $S(x, y)$  [9] which is defined as:

$$S(x, y) = \begin{cases} 1, & \text{if } (y - x) \geq 0.5, \\ 2(y - x), & \text{if } 0.5 \geq (y - x) \geq 0, \\ \frac{\cos(4\pi r) + 1}{2}, & \text{if } r \leq \frac{1}{4}, \\ 0, & \text{otherwise,} \end{cases} \quad (5.1)$$

where  $r = \sqrt{(x - 1.5)^2 + (y - 0.5)^2}$ .

The interpolated function  $S(x, y)$  is shown in Figure 6. Meanwhile, we show the positive surface generated by our proposed method at a random set of 100 points in Figure 7.

In order to measure the quality of new interpolant, we carry out the following experiment. Firstly, we construct two interpolants by constrained modified quadratic Shepard method and our proposed method, based on a series of data sets where the number

**Table 4:** Variances for two interpolants for  $S(x, y)$ .

Number of data points	30	60	100	150	250
CMQS	1.54	1.20	0.97	0.54	0.11
Our proposed method	2.32	1.97	1.55	0.97	0.24

of randomly chosen points increases. Secondly, we evaluate the interpolants on a grid of  $25 \times 25$  points and calculate the variances between the exact and calculated values. The results are shown in Table 4.

## 6. Conclusion

Positivity preserving interpolation is an interesting work. In this paper, we discuss the approximation and interpolation problem of scattered data by using CSRBF under the constraint of positivity. The approaches are presented to compute positive approximation and interpolation by solving the two corresponding optimization problems.

However, we have not discussed how to select the optimal number and position of the new added data  $y_i$  in order to both achieve the existence and better possible approximation behavior of interpolation. These problems would be attractive and remain to be our future work.




## Acknowledgments

This work was supported by the Natural Science Foundation of Zhejiang Province (nos. Y7080068 and Y6090211), the Foundation of Department of Education of Zhejiang Province (nos. Y200802999 and Y200907579), and Tianyuan Foundation for Mathematics of China (no. 10926035). The authors are very grateful to the referees for their careful reading of the manuscript and many valuable comments and remarks, which greatly improved this paper.

## References

- [1] J. W. Schmidt and W. Hess, "Positivity of cubic polynomials on intervals and positive spline interpolation," *BIT*, vol. 28, no. 2, pp. 340–352, 1988.
- [2] S. Butt and K. W. Brodlie, "Preserving positivity using piecewise cubic interpolation," *Computers and Graphics*, vol. 17, no. 1, pp. 55–64, 1993.
- [3] B. H. Ong and H. C. Wong, "A  $C^1$  positivity preserving scattered data interpolation scheme," in *Advanced Topics in Multivariate Approximation*, vol. 8, pp. 259–274, World Scientific, Singapore, 1996.
- [4] M. Sarfraz, "Piecewise rational interpolation preserving positive data," in *Proceedings of the International Conference on Imaging Science, Systems, and Technology (CISST '00)*, pp. 103–109, Las Vegas, Nev, USA, 2000.
- [5] A. R. M. Piah, A. Saaban, and A. A. Majid, "Range restricted positivity-preserving scattered data interpolation," *Journal of Fundamental Sciences*, vol. 2, no. 1-2, pp. 63–75, 2006.
- [6] M. Z. Hussain and M. Sarfraz, "Positivity-preserving interpolation of positive data by rational cubics," *Journal of Computational and Applied Mathematics*, vol. 218, no. 2, pp. 446–458, 2008.
- [7] F. I. Utreras, "Positive thin plate splines," *Approximation Theory and Its Applications*, vol. 1, no. 3, pp. 77–108, 1985.
- [8] M. R. Asim and K. W. Brodlie, "Curve drawing subject to positivity and more general constraints," *Computers and Graphics*, vol. 27, no. 4, pp. 469–485, 2003.

- [9] K. W. Brodlie, M. R. Asim, and K. Unsworth, "Constrained visualization using the shepard interpolation family," *Computer Graphics Forum*, vol. 24, no. 4, pp. 809–820, 2005.
- [10] M. D. Buhmann, *Radial Basis Functions: Theory and Implementations*, Cambridge Monographs on Applied and Computational Mathematics, Cambridge University Press, Cambridge, UK, 2000.
- [11] R. Franke, "Scattered data interpolation: tests of some methods," *Mathematics of Computation*, vol. 38, no. 157, pp. 181–200, 1982.
- [12] M. J. D. Powell, "Radial basis function for multivariable interpolation: a review," in *Numerical Analysis*, D. Griffiths and G. Waston, Eds., pp. 223–241, Longman Scientific and Technical, Harlow, UK, 1987.
- [13] R. Schaback, "Creating surfaces from scattered data using radial basis functions," in *Mathematical Methods for Curves and Surfaces*, T. Lyche and L. Schumaker, Eds., pp. 477–496, Vanderbilt University, Nashville, Tenn, USA, 1995.
- [14] Z. M. Wu, "The radial basis functions-a survey," *Advances in Mathematics*, vol. 27, no. 3, pp. 202–208, 1998 (Chinese).
- [15] Z. M. Wu, "Compactly supported positive definite radial functions," *Advances in Computational Mathematics*, vol. 4, no. 3, pp. 283–292, 1995.
- [16] H. Wendland, "Piecewise polynomial, positive definite and compactly supported radial functions of minimal degree," *Advances in Computational Mathematics*, vol. 4, no. 4, pp. 389–396, 1995.
- [17] M. S. Floater and A. Iske, "Multistep scattered data interpolation using compactly supported radial basis functions," *Journal of Computational and Applied Mathematics*, vol. 73, no. 1–2, pp. 65–78, 1996.
- [18] D. Lazzaro and L. B. Montefusco, "Radial basis functions for the multivariate interpolation of large scattered data sets," *Journal of Computational and Applied Mathematics*, vol. 140, no. 1–2, pp. 521–536, 2002.
- [19] P. Gordan, "Über die Auflösung linearer Gleichungen mit reellen Coefficienten," *Mathematische Annalen*, vol. 6, no. 1, pp. 23–28, 1873.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

